

Critical Care Nephrology

64

Javier A. Neyra | J. Pedro Teixeira | Ashita J. Tolwani

CHAPTER OUTLINE

ACUTE KIDNEY INJURY IN THE CONTEXT OF CRITICAL ILLNESS, 1882
FLUID MANAGEMENT AND ITS IMPACT ON ACUTE KIDNEY INJURY, 1886

KIDNEY REPLACEMENT THERAPY, 1889
EXTRACORPOREAL BLOOD PURIFICATION, 1896

KEY POINTS

- Critical care nephrology is an evolving field that requires understanding of the pathobiology of critical illness as a multisystem condition characterized by crosstalk between multiple organs during injury, repair, and recovery.
- Critically ill in each patient involves the interplay of complex underlying comorbidities and specific acute challenges, leading to an individualized risk of incident acute kidney injury (AKI), AKI progression, requirement for kidney replacement therapy (KRT), and adverse post-AKI health care outcomes.
- There is no benefit to early goal-directed therapy for the management of sepsis, with regard to either mortality or prevention of AKI.
- Fluid therapy requires dynamic assessment of tolerance and responsiveness, incorporated into the dynamic timeline of phases of resuscitation, optimization, stabilization, and evacuation.
- The use of balanced crystalloids for resuscitation may reduce the risk of AKI in critical illness.
- In the context of AKI, accelerated (vs. standard) initiation of KRT is not recommended given the higher risk of complications including KRT dependence. Therefore the decision to initiate KRT should be individualized according to patient-specific solute and volume-control goals.
- The selection of KRT modality to support patients with AKI should consider available resources, cost, provider expertise, and patient needs.
- In the setting of acute liver failure or traumatic brain injury, continuous KRT (CRRT) is the preferred KRT modality to mitigate the risks of cerebral edema and herniation.
- The selection of CRRT modality to support critically ill patients with AKI should be considered according to available devices, protocol logistics, and provider expertise given that there is no objective evidence to favor one modality over the other.
- For critically ill patients with AKI receiving CRRT, key aspects of the CRRT prescription that should be standardized according to current evidence are 1. the prescription (on average) of a total effluent dose of 25 to 30 mL/kg/h; 2. the use of right internal jugular temporary catheters to minimize treatment interruptions; 3. the use of regional citrate anticoagulation—if feasible—to maximize filter life; and 4. the iterative monitoring of patient tolerance of fluid removal, fluid balance goals, and net ultrafiltration rate (UF_{NET}).
- Extracorporeal blood purification through adsorption-based technologies have gained prominence after the COVID-19 pandemic, although specific applications and protocols are the subject of ongoing investigation.

Studies now suggest that up to 60% of patients in intensive care units (ICUs) experience an episode of acute kidney injury (AKI).¹⁻³ In the most severe cases, kidney replacement therapy (KRT) is required. Short-term mortality among critically ill patients with AKI who require KRT is >50%⁴

and, among survivors, longevity and health-related quality of life are often poor.⁵ AKI and the resulting electrolyte and acid-base abnormalities occur on the background of multiple conditions that necessitate critical care. In this chapter, we first discuss AKI in the context of conditions

that are frequently encountered in the ICU including sepsis, cardiac surgery, liver failure, and acute respiratory distress syndrome (ARDS). This is followed by a discussion of fluid management and its relevance to the care of critically ill patients with AKI. We then review the principles of KRT prescription and delivery to critically ill patients with AKI. Finally, we discuss emerging technologies of blood purification for critically ill patients with AKI. We do not discuss acute brain injury in the ICU setting, which is associated with dysnatremias (including the syndrome of inappropriate secretion of antidiuretic hormone, cerebral salt wasting, and central diabetes insipidus). The reader is referred to [Chapter 14](#) for further discussion of these disorders.

ACUTE KIDNEY INJURY IN THE CONTEXT OF CRITICAL ILLNESS

A list of common etiologies and risk factors for AKI, specifically in ICU settings, is provided in [Table 64.1](#). Selected clinical contexts of AKI in critically ill adult patients are discussed as follows.

SEPSIS-ASSOCIATED ACUTE KIDNEY INJURY

Sepsis is the most common cause of AKI in the ICU, contributing to approximately 50% of cases.^{2,4,6} A large epidemiologic study found that sepsis-associated AKI (SA-AKI) was present in one of six ICU patients.⁷ SA-AKI is associated with substantially higher mortality than

sepsis without AKI or nonseptic AKI.^{6,9} Notably, the relationship between AKI and sepsis appears bidirectional, as accumulating data suggest that AKI is a risk factor for the subsequent development of secondary infections and de novo sepsis.¹⁰⁻¹² In 2023 a definition for SA-AKI was proposed.¹³ It combines the consensus criteria for both sepsis, using the sepsis-3 definition¹⁴ and the Kidney Disease: Improving Global Outcomes (KDIGO) AKI criteria.¹⁵ The traditional understanding of the pathophysiology of SA-AKI was that sepsis triggered acute tubular injury via global hypoperfusion.¹⁶ However, both animal and human studies have demonstrated that, while renal blood flow (RBF) appears to decline later in established SA-AKI, macrovascular RBF is increased in early SA-AKI, with relatively preserved histology with minimal or patchy tubular injury without necrosis.¹⁷⁻²⁵ The pathophysiology of SA-AKI is complex and heterogeneous, involving microcirculatory dysfunction, inflammation, oxidative stress, mitochondrial dysfunction, altered cellular bioenergetics, and endothelial dysfunction.²⁶⁻²⁸ In this context, one should recognize that while SA-AKI often coexists with ischemic acute tubular necrosis (ATN) in the setting of overt shock, sepsis without hypotension can also cause AKI.

Despite multiple prior and ongoing studies (NCT00711789, NCT02664753, NCT03901807, NCT05900284),²⁹ therapeutic trials specifically targeting SA-AKI have thus far failed to demonstrate clear benefits.³⁰⁻³² Thus management of SA-AKI remains largely supportive, with timely antibiotics, infectious source control, and hemodynamic optimization.

ACUTE KIDNEY INJURY IN THE SETTING OF HEART DISEASE

Since cardiorenal syndromes (CRS) are discussed in detail in [Chapter 42](#), we focus on AKI in the setting of cardiac surgery or mechanical cardiopulmonary support.

CARDIAC SURGERY-ASSOCIATED ACUTE KIDNEY INJURY

Cardiac surgery-associated AKI (CSA-AKI) is the second most common cause of AKI in the ICU and is associated with increased mortality.^{4,33-36} CSA-AKI is defined as AKI occurring 2 to 7 days after surgery and is reported, by KDIGO criteria, to complicate 20% to 30% of cardiac surgeries.³⁶⁻³⁸ The pathophysiology of CSA-AKI is multifactorial including hypoperfusion, inflammation, oxidative stress, atheroembolism, hemolysis, nephrotoxin exposure, and exposure to cardiopulmonary bypass (CPB).^{36,37} The duration of CPB is a significant predictor of AKI, but extensive randomized controlled trials (RCTs) aimed at avoiding or reducing CPB have yielded mixed results.³⁹⁻⁴¹ Similarly, RCTs of therapeutics to prevent or treat CSA-AKI have failed to result in benefits.⁴²⁻⁴⁵ Single-center RCTs^{46,47} have found that perioperative nitric oxide (NO) delivered via the CPB circuit or ventilator decreases the rate of CSA-AKI or chronic kidney disease (CKD), presumably via correction of hemolysis-induced depletion of renal NO. Though promising, the use of NO to prevent CSA-AKI remains investigational and additional RCTs are ongoing (NCT04216927, NCT05757557).^{37,48} The use of biomarkers to prevent CSA-AKI also holds promise. The multicenter PrevAKI study randomized 278 postoperative patients at

Table 64.1 Common Etiologies and Risk Factors for Acute Kidney Injury (AKI) in the Intensive Care Unit

Etiologies/Contributing Factors ^{2,4a}	Risk Factors/Predisposing Conditions ^{2,515}
- Sepsis/septic shock: 40%-50% - Major surgery: 35% - Volume depletion: 25%-35% - Cardiogenic shock: 15%-25% - Drug toxicity: 15%-20% - Hepatorenal syndrome: 3%-6% - Urinary tract obstruction: 1%-3%	- Chronic kidney disease - Prior AKI - Advanced age - Frailty⁵¹⁶ - Nonrenal comorbidities: - Diabetes - Heart disease, especially CHF - Liver disease - Malignancy - Treatment with diuretics, NSAIDs, or RAAS inhibitors

^aThe frequencies of these etiologies/factors are derived from Uchino S, Kellum JA, Bellomo R, et al. Acute renal failure in critically ill patients: a multinational, multicenter study. *JAMA*. 2005;294(7):813-818 and Hoste EA, Bagshaw SM, Bellomo R, et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med*. 2015;41(8):1411-1423 but are rounded for simplicity. CHF, Congestive heart failure; NSAID, nonsteroidal anti-inflammatory drug; RAAS, renin-angiotensin-aldosterone system.

high risk of CSA-AKI, identified on the basis of elevated urinary tissue inhibitor of metalloproteinases 2 (TIMP-2) * insulin-like growth factor binding protein 7 (IGFBP7), to standard care or the implementation of a KDIGO-based AKI bundle, which includes minimization of nephrotoxic medications, prevention of hyperglycemia, and volume optimization via hemodynamic monitoring.⁴⁹ Though the need for KRT and mortality were similar in both groups, the rate of stage 2-3 AKI was significantly reduced.⁴⁹

ACUTE KIDNEY INJURY IN PATIENTS REQUIRING MECHANICAL CARDIOPULMONARY SUPPORT

The use of extracorporeal membrane oxygenation (ECMO) in adults grew exponentially after the 2009 H1N1 influenza outbreak, with overall ECMO usage in adults more than tripling between 2008 and 2014.⁵⁰ Notably, 2023 guidelines recommend the use of venovenous (VV-) ECMO for refractory ARDS.⁵¹

Up to 85% of adults on ECMO develop AKI, with roughly 45% requiring KRT.⁵²⁻⁵⁷ As in other settings, AKI in patients requiring ECMO is independently associated with mortality.⁵²⁻⁵⁶ AKI develops more commonly in patients receiving venoarterial- (VA-) ECMO for cardiac failure than those on VV-ECMO for isolated respiratory failure.⁵⁶⁻⁵⁸ Observational data suggest that timely KRT may be beneficial in preventing or mitigating volume overload in ECMO patients.^{59,60} Fluid overload in ECMO patients is independently associated with increased mortality⁶¹⁻⁶⁴, and guidelines⁶⁵ suggest targeting euolemia with diuretics or KRT once feasible to facilitate ECMO weaning. However, experts recommend against *preemptive* KRT use in patients on ECMO, given a lack of data for benefit of KRT beyond the usual indications.⁶⁶

The pathogenesis of AKI on ECMO is complex. ECMO-specific factors that contribute to AKI include inflammation and disordered coagulation induced by the circuit, pigment injury from hemolysis, and nonpulsatile blood flow in VA-ECMO.⁶⁷ Excess circuit-related hemolysis may manifest with pink urine or KRT effluent and can be confirmed by measuring plasma-free hemoglobin levels.^{61,68-70}

Like ECMO, the use of left ventricular assist devices (LVADs) to treat end-stage heart failure has grown dramatically during the past 2 decades.^{71,72} On average, LVAD placement produces an initial improvement in kidney function, likely reflecting improved renal perfusion in patients with type 2 CRS (i.e., kidney dysfunction due to chronic cardiac dysfunction), followed by a gradual decline back toward baseline.⁷³⁻⁷⁶ However, kidney function may not improve in those with intrinsic kidney disease. Like patients receiving ECMO, patients managed with LVAD are at elevated risk for AKI throughout their course, and AKI (particularly AKI requiring KRT) is associated with poor outcomes.⁷⁵⁻⁷⁸ LVAD-specific causes of AKI include hemolysis, right ventricular failure, and possibly nonpulsatile blood flow.^{75,76}

ACUTE KIDNEY INJURY IN THE SETTING OF RESPIRATORY FAILURE AND ACUTE RESPIRATORY DISTRESS SYNDROME

Acute kidney injury is an independent risk factor for respiratory failure and vice versa, and the presence of

either condition complicating the other is associated with a greater than twofold increased risk of death.⁷⁹⁻⁸² Though traditionally attributed to fluid overload, the high mortality of combined respiratory and kidney failure may be mediated by kidney-lung crosstalk. This crosstalk involves a variety of complex pathophysiologic mechanisms including the inflammatory effects of AKI on the lung endothelium mediated by multiple cellular and cytokine mediators (e.g., interleukin-6), impaired alveolar fluid clearance due to downregulation of pulmonary sodium and water channels, and the deleterious hemodynamic effects of mechanical ventilation on RBF and microvascular flow.^{66,82,83}

ARDS is among the most severe forms of respiratory failure with an in-hospital mortality rate of approximately 40%.⁸⁴ ARDS is defined as acute-onset pulmonary edema not explained by cardiac failure or fluid overload resulting in hypoxemia with a PaO₂-to-FiO₂ ratio of ≤ 300 mm Hg.^{85,86} Treatment consists of addressing the underlying cause and providing lung protective mechanical ventilation (lung protective ventilation) with limitations in tidal volume and inspiratory pressures and tolerance of hypercapnia, a ventilatory strategy that was shown in the ARDS Network trial to reduce mortality.⁸⁷ Patients in the lung protective ventilation group had less renal dysfunction,⁸⁷ consistent with preclinical and epidemiologic data that injurious mechanical ventilation may induce AKI.⁸⁸⁻⁹⁰ Permissive hypercapnia normally triggers compensatory renal bicarbonate reabsorption.

Achieving permissive hypercapnia without causing severe acidemia in patients with AKI may necessitate supplemental bicarbonate or KRT,^{87,91} especially in severe ARDS when respiratory carbon dioxide (CO₂) elimination is severely impaired. However, the volume load of sodium bicarbonate may be prohibitive in patients with severe hypoxemia,⁹¹ and rapid administration of bicarbonate may abruptly worsen hypercarbia.⁹² If volume overload is present, correction of acidemia may be preferred through KRT. An alternative method to correct acidosis with lung protective ventilation is using an extracorporeal CO₂ removal (ECCO₂R) device, although trial data in ARDS are lacking.⁹³

Avoidance of fluid overload in ARDS is essential. A conservative fluid management strategy (i.e., less fluid/more diuretics) in ARDS was shown in a large multicenter RCT to result in shorter duration of mechanical ventilation and ICU stay without affecting mortality.⁹⁴ The conservative arm also had a trend toward decreased need for KRT (10 vs. 14%, $P = 0.06$)⁹⁴ and, when adjusted post hoc for the effect of fluid balance on serum creatinine, a decreased rate of AKI.⁹⁵ KRT in patients with ARDS without AKI to achieve negative or even fluid balance has been studied in some small trials and, though controversial, may be considered for refractory hypoxemia in some instances.⁹⁶⁻⁹⁸

ACUTE KIDNEY INJURY IN THE SETTING OF LIVER DISEASE

ACUTE KIDNEY INJURY IN THE SETTING OF CIRRHOSIS AND HEPATORENAL SYNDROME

Hepatorenal syndrome (HRS), particularly HRS-AKI, which was previously referred to as type 1 HRS, is a complication of decompensated cirrhosis associated with high mortality in which the vascular and neurohormonal disturbances of

advanced liver disease and portal hypertension result in profound renal hypoperfusion unresponsive to intravenous (IV) albumin or other fluids.⁹⁹⁻¹⁰² In modern cohorts, HRS-AKI accounts for 12% of AKI cases in the setting of cirrhosis, whereas prerenal AKI (44%) and ATN (30%) are significantly more common.¹⁰³ Notably, the mortality of HRS and ATN in the setting of cirrhosis is similar at approximately 50% at 90 days.^{103,104}

Developments in the diagnosis of HRS-AKI include adoption of the KDIGO criteria for AKI diagnosis.^{99,105} In addition, though these urinary indices tend to perform poorly in ICU patients in differentiating prerenal AKI from ATN,¹⁰⁶ cutoffs of fractional excretion of sodium (FENa) and urea (FeUrea) have been suggested to differentiate HRS from ATN or prerenal AKI (with FENa <0.2% and FeUrea <28.2% being consistent with HRS-AKI and <0.1% and <21%, respectively, being especially suggestive).¹⁰⁷⁻¹⁰⁹ Furthermore, though the absence of proteinuria >500 mg/g creatinine remains a traditional diagnostic criterion for HRS, urinary albumin has been shown in studies to be useful in differentiating ATN and HRS, with albuminuria >100 mg/g creatinine being suggestive of ATN.^{109,110} Likewise, studies suggest that the tubular injury biomarker neutrophil gelatinase-associated lipocalin (NGAL) may help differentiate HRS and ATN, with a urinary NGAL level >220 to 250 µg/g creatinine being consistent with ATN.¹⁰⁹⁻¹¹² Despite these diagnostic advances, the diagnosis of HRS-AKI in clinical practice remains a tremendous challenge. Though the diagnostic criteria imply that HRS is a diagnosis of exclusion, HRS can coexist with ATN, CRS from cirrhotic cardiomyopathy, abdominal compartment syndrome from tense ascites, or other unrelated causes of AKI or CKD.¹⁰⁰⁻¹⁰²

Terlipressin has been validated as an effective treatment for HRS-AKI in the Study to Confirm Efficacy and Safety of Terlipressin in Hepatorenal Syndrome (HRS) Type 1 (CONFIRM).¹¹³ Though CONFIRM demonstrated a significant increase in HRS reversal (29% vs. 16% in placebo arm), the terlipressin arm also experienced an increased risk of adverse effects including respiratory failure (10% with terlipressin vs. 3% in the placebo arm) and death due to respiratory failure (11% vs. 2%, respectively). This is thought to result from excessive albumin use and the regional hemodynamic effects of terlipressin including intense systemic vasoconstriction, pulmonary venoconstriction, lack of pulmonary arterial vasoconstriction, and absent inotropic effect.¹¹⁴ The positive primary outcome of the CONFIRM trial resulted in FDA approval of terlipressin in 2022 with a black box warning about the risk of fatal respiratory failure. Contraindications to terlipressin include any hypoxemia or need for supplemental oxygen, acute-on-chronic liver failure (ACLF) grade 3, and serum creatinine >5.0 mg/dL, as these patients are at increased risk of respiratory failure or treatment nonresponse.^{108,113,115,116} Norepinephrine is recommended as an alternative to terlipressin, as it appears to have similar efficacy in achieving HRS reversal, but it requires ICU admission in most centers.^{99,108} Octreotide and midodrine are not recommended because they are rarely effective and may cause harm by delaying effective therapy.^{99,108,113}

The results of CONFIRM and a second large RCT published in 2021 examining albumin use in hospitalized patients with cirrhosis¹¹⁷ highlight the risk of harmful

volume overload from excessive albumin administration in patients with liver disease. Similarly, the association between intraabdominal hypertension (IAH) and AKI in cirrhosis, as well as the potential for improved kidney function through cautious therapeutic paracentesis, is gaining recognition.¹¹⁸⁻¹²¹ Therefore while albumin remains the initial treatment for most cirrhosis patients with AKI, it is important to first assess their volume status thoughtfully. Furthermore, given data that delay in implementation of effective vasoconstrictor therapy increases the risk of treatment failure,^{99,108,113} those who felt likely to have HRS may benefit from initiation of vasoconstrictors before failing 48 hours of albumin therapy.^{100-102,122}

ACUTE LIVER FAILURE

Acute liver failure (ALF), previously known as fulminant hepatic failure, is defined by 1. hepatic encephalopathy, 2. coagulopathy (international normalized ratio [INR] ≥1.5), 3. onset of illness <26 weeks, and 4. no evidence of cirrhosis.^{123,124} The distinction between ALF and chronic liver disease is vitally important because some of the complications and management options apply specifically to ALF. Common causes of ALF include toxic, viral, and ischemic liver injury, with acetaminophen toxicity alone accounting for about 50% in modern cohorts.¹²³⁻¹²⁵ Though some cases will spontaneously resolve, the mortality of ALF is high with supportive care alone.¹²³ Complications of ALF overlap with decompensated cirrhosis and include hepatic encephalopathy AKI, coagulopathy and bleeding, hypoglycemia, infection and sepsis, and multiorgan failure.¹²⁵ However, in contrast to cirrhosis, in which clinically significant cerebral edema is characteristically absent,^{126,127} severe hepatic encephalopathy from ALF is frequently accompanied by cerebral edema with progressive intracranial hypertension and risk of lethal brainstem herniation. Cerebral edema is present in up to 35% of patients with ALF and grade 3 hepatic encephalopathy and 75% of patients with grade 4 hepatic encephalopathy.¹²³ First-line treatment for confirmed or suspected intracranial hypertension is hyperosmolar therapy with mannitol, though it requires monitoring of serum osmolality and its use may be limited by renal dysfunction in those not on KRT.^{123,128} Similarly, hypertonic saline to achieve serum sodium of 145 to 155 mEq/L is recommended in patients with or at high risk of cerebral edema.^{123,129}

AKI in the setting of ALF may be due to renal hypoperfusion (e.g., sepsis, bleeding, and HRS) or intrinsic injury (e.g., ischemic ATN, direct tubular toxicity of acetaminophen or *Amanita*) and is associated with increased risk of cerebral edema and overall poor prognosis.¹²³ For a variety of reasons, patients with ALF may require KRT relatively early in the course of their AKI.¹²⁵ Typically, continuous renal replacement therapy (CRRT) is preferred to intermittent hemodialysis (IHD) to minimize the risk of hypotension and increase in intracranial pressure (ICP), which can combine to severely reduce cerebral perfusion pressure. CRRT also offers more flexibility to manage volume status, clear ammonia, and precisely manage hyponatremia, which frequently accompanies ALF and may aggravate cerebral edema.^{123,130-132} In contrast to chronic liver disease, KRT does appear to have a role in specifically treating hyperammonemia in ALF in the absence of AKI,

particularly when ammonia is >250 $\mu\text{g/L}$, with higher-than-standard doses of CRRT typically required to effectively clear ammonia.^{131,133} Beyond the risk of hypotension and increased ICP, a preference for CRRT over IHD in ALF is supported by limited observational and trial data.^{124,130}

ABDOMINAL COMPARTMENT SYNDROME

Intraabdominal hypertension and abdominal compartment syndrome are increasingly recognized as common causes of decreased kidney function in ICU patients.¹³⁴⁻¹³⁷ Intraabdominal hypertension is defined as a sustained elevation in intraabdominal pressure (IAP) ≥ 12 mm Hg, and abdominal compartment syndrome is defined as sustained IAP >20 mm Hg associated with new organ dysfunction. Intraabdominal hypertension has been reported in 30% to 50% of mixed medical-surgical ICU patients, with abdominal compartment syndrome occurring in 5% to 12%.¹³⁵⁻¹³⁸ Risk factors for IAH include trauma, major burns, abdominal surgery, mechanical ventilation, obesity, ascites, hemoperitoneum, gastric or bowel distention, large-volume resuscitation, and pancreatitis.¹³⁸ Physical examination is unreliable for diagnosis.^{134,138,139} Instead, to estimate IAP, 25 to 30 mL of sterile water is instilled into the urinary bladder via a Foley catheter and pressure tubing is used to transduce bladder pressure at end-expiration with the patient relaxed in the supine position.¹³⁹

IAH can decrease the perfusion of any abdominal organ, and effects can be transmitted to other compartments leading to decreased cardiac output (CO), impaired ventilation, and increased ICP. The kidneys are particularly sensitive such that abdominal compartment syndrome is thought to be unlikely in the absence of oliguria. Decreased kidney function from IAH appears to be due to reduced perfusion mediated primarily by elevations in renal venous and parenchymal pressures (rather than by decreased CO or ureteral compression).^{138,139} Ultimately, a functional prerenal state develops with neurohormonal activation, renal vasoconstriction, renal sodium retention, and oliguria. IAH has been implicated in the pathogenesis of both CRS and HRS, with small studies in both settings suggesting that paracentesis may improve renal function.^{118-121,139,140}

In established abdominal compartment syndrome, decompressive laparotomy remains the treatment of choice despite its high morbidity.^{134,138,139} Less invasive measures to control IAH include sedation and potentially neuromuscular blockade to decrease abdominal muscle tone and decompression through paracentesis or nasogastric suction. Fluid management is complex in the setting of IAH. While decreased CO in abdominal compartment syndrome can worsen with hypovolemia, excessive fluid intake is associated with a higher risk of IAH. Any benefit from fluids following initial resuscitation is likely to be temporary, and giving too much fluid may trigger a harmful cycle where fluids are administered for a short-term improvement in CO but lead to increased IAH and subsequent reduced CO, prompting further fluid administration.¹³⁷ Animal model data suggest that IV fluids may restore CO but do not restore organ function in abdominal compartment syndrome.¹⁴¹ Traditional measures of volume status can be unreliable in abdominal compartment syndrome, as central venous and pulmonary artery occlusion pressures are increased, and

ultrasound may show a collapsed inferior vena cava, all due to IAH. Small case series suggest that CRRT can lower IAP, but the role of KRT for patients with IAH without kidney dysfunction remains unclear.^{139,142,143} When providing KRT to patients with IAH, the femoral vein should be avoided for dialysis catheter placement because IAH may lead to recirculation and impaired clearance.¹⁴⁴

KEY CONCEPTS IN NEPHROTOXIC ACUTE KIDNEY INJURY

In critically ill patients with AKI, medications must be reviewed frequently in order to 1. prevent worsening or prolongation of AKI due to avoidable nephrotoxic exposure, 2. prevent toxic accumulation of nonnephrotoxic drugs that may cause extrarenal toxicity in the presence of impaired kidney function (e.g., neurotoxicity from morphine, gabapentinoids, or cefepime),¹⁴⁵⁻¹⁴⁷ and 3. ensure proper drug dosing for patients receiving KRT. The topics of drug-induced acute interstitial nephritis (AIN), drug dosing in AKI, and kidney disease from anticancer drugs are reviewed in detail elsewhere (Chapters 37, 56, and 57, respectively). We focus on two evolving topics in drug-induced acute tubular injury in critically ill patients.

VANCOMYCIN-ASSOCIATED ACUTE KIDNEY INJURY

Though controversy persists given inherent limitations in observational data, vancomycin has been increasingly recognized as a contributor to AKI in critically ill patients.¹⁴⁸⁻¹⁵⁰ The nephrotoxicity of vancomycin was initially attributed to the impurities of early formulations, referred to as “Mississippi mud.”¹⁴⁸ Though current purified formulations are clearly less toxic than prior preparations, increasing rates of methicillin-resistant *Staphylococcus aureus* (MRSA) infections¹⁵¹ and the concomitant increase in vancomycin use have led to increasingly frequent reports of vancomycin-associated AKI (VA-AKI) with modern formulations.¹⁴⁸ In meta-analyses, the reported rates of VA-AKI ranged from 5% to 43% of vancomycin-treated patients with a relative risk of AKI with vancomycin of 2.45 compared with other antimicrobials (mostly linezolid).^{152,153} VA-AKI has a mean onset of 4 to 17 days after drug initiation and approximately 75% of cases improve or resolve by hospital discharge.^{148,152}

Vancomycin undergoes no metabolism and is entirely cleared unchanged by the kidneys.¹⁴⁸ The mechanism of VA-AKI appears to be toxic acute tubular injury caused by oxidative injury and obstructive tubular casts.^{148,154,155} The casts are composed of noncrystal nanospheric vancomycin aggregates embedded in uromodulin, as demonstrated by immunohistochemistry of human kidney biopsy and urinary sediment specimens, and reproduced in mouse models.^{154,155} Risk factors include both patient factors—baseline kidney function, obesity, severity of illness, concomitant nephrotoxin exposure (e.g., aminoglycosides), and age—and drug factors like higher target trough levels (i.e., >15 mg/L, as used for severe MRSA infections); higher total daily dose (i.e., >4 g per day); intermittent dosing (rather than continuous infusions); and longer duration of therapy (i.e., >7 days).^{148,152,153,156-159}

An additional reported risk factor for VA-AKI is concomitant exposure to piperacillin-tazobactam (P-TZ),

which alone is not generally considered nephrotoxic (except for rare cases of AIN). Though not found in all studies, a signal for increased rates of AKI with this combination compared with vancomycin alone has been repeatedly detected in the literature.¹⁶⁰⁻¹⁶⁸ However, with some exceptions,¹⁶⁰ the majority of the studies tying the combination of P-TZ and vancomycin to AKI have failed to detect a difference in need for KRT or mortality,¹⁶⁷⁻¹⁷² outcomes strongly linked to AKI. Moreover, the mechanism of this vancomycin-specific additive nephrotoxicity of P-TZ is unknown and multiple attempts to replicate this synergistic nephrotoxicity in animal models and in vitro studies have failed.¹⁷³⁻¹⁷⁵ One proposed explanation for these contradictory findings is that P-TZ, which is excreted via organic anion transporters,¹⁷⁶ inhibits tubular secretion of creatinine, leading to increased rates of creatinine-defined pseudo-AKI without any associated tubulopathy or morbidity. Support for this theory comes from a 2022 biomarker study of nearly 200 ICU patients treated with vancomycin and P-TZ or vancomycin and cefepime. The P-TZ-vancomycin combination was associated with higher rates of creatinine-defined AKI, but there was no association with AKI as defined by cystatin C, need for KRT, or mortality.¹⁷⁷ Subsequently, the 2023 Antibiotic Choice on Renal Outcomes (ACORN) study randomized more than 2500 adults hospitalized for acute infection to cefepime or P-TZ and, though 77% of the participants were also on vancomycin at enrollment, the rates of AKI or death were similar in both arms while the cefepime arm had increased rates of delirium or coma.¹⁷⁸ Considering that kidney dysfunction is a major risk factor for cefepime neurotoxicity,¹⁴⁷ these results suggest that cefepime should not be used preferentially to PT-Z in combination with vancomycin or in kidney dysfunction. However, as observational data suggest that synergistic nephrotoxicity is more likely with longer courses,^{160,179} caution may be warranted with prolonged courses of PT-Z and vancomycin.

ACUTE KIDNEY INJURY ASSOCIATED WITH INTRAVENOUS IODINATED CONTRAST

Emerging data suggest that IV iodinated contrast, as given with computed tomography (CT), is less nephrotoxic than previously thought. Although a few exceptions exist,^{180,181} more than a dozen studies and meta-analyses¹⁸²⁻¹⁹⁶ analyzing thousands of patients across a variety of settings have been published since 2013, suggesting that modern low- or iso-osmolar IV contrast exposure has little to no nephrotoxicity. Many of these analyses focused on patients at higher risk of contrast-associated AKI (CA-AKI) due to underlying CKD. A 2023 multicenter analysis of >14,000 patient encounters, using weighting by propensity scores and entropy balancing to mitigate confounding and biases, found no increased risk of persistent kidney dysfunction or need for KRT in patients with established AKI before IV contrast exposure.¹⁹¹ While the risk of CA-AKI and relevant AKI outcomes after IV contrast administration is far lower than previously thought, the risk may not be zero, and measures to reduce the risk, such as minimization of contrast exposure and provision of IV fluids when appropriate, should still be pursued.¹⁹⁷⁻²⁰⁰

Importantly, nearly all of these studies questioning the nephrotoxicity of iodinated contrast analyzed IV rather than

intraarterial contrast exposure. Both modern and historical data support the notion that intraarterial contrast exposure is more nephrotoxic, though many have speculated that this differential risk relates primarily to procedural aspects of angiography, such as the risk of subclinical atheroembolism, rather than enhanced toxicity related to the route of contrast administration.¹⁹⁹⁻²⁰⁵ Regardless of the mechanism, the risk of AKI after intraarterial contrast exposure needs to be carefully considered in at-risk patients, and implementation science to mitigate the risk of AKI after cardiac catheterization has shown promise.^{206,207}

Observational studies examining coronary angiography in advanced CKD patients with acute coronary syndromes reveal that although CKD patients face a higher risk of AKI or death compared to non-CKD patients, those with CKD who undergo angiography have lower mortality rates than CKD patients who do not receive indicated angiography.²⁰⁸⁻²¹¹ Though these observational studies are subject to residual confounding and biases, they suggest that withholding indicated diagnostic tests or interventions in patients with renal dysfunction due to excessive concern about AKI may inadvertently lead to worse outcomes, a concept termed “renalism.” Though data demonstrating that “renalism” exists in ICU populations are lacking, many experts^{199,200,203,205,212} and 2020 consensus guidelines²¹³ recommend performing studies using IV contrast regardless of kidney function when needed for definitive diagnosis or treatment of life-threatening conditions (e.g., sepsis or suspected pulmonary embolism).

FLUID MANAGEMENT AND ITS IMPACT ON ACUTE KIDNEY INJURY

VOLUME MANAGEMENT IN THE INTENSIVE CARE UNIT

The approach to fluid administration in the ICU has undergone an evolution from the adage that patients “must swell to get well”—driven by the influence of the landmark 2001 single-center early goal-directed therapy (EGDT) trial by Rivers et al.²¹⁴—to an increasing appreciation that volume overload is a marker and potentially a mediator of poor outcomes in critical illness. Three subsequent large multicenter RCTs, ProCESS (Protocol-Based Care for Early Septic Shock), ARISE (Australasian Resuscitation in Sepsis Evaluation), and ProMiSe (Protocolised Management in Sepsis), collectively randomized more than 4000 patients with septic shock and demonstrated no benefit of EGDT over standard care.²¹⁵⁻²¹⁷ Furthermore, observational studies and small trials have established that positive fluid balance, volume overload, and elevated filling pressures in patients with sepsis, AKI, or other critical illnesses are associated with poor outcomes including progressive kidney dysfunction, impaired AKI recovery, increased risk of subsequent sepsis, impaired wound healing, prolonged mechanical ventilation, and death.^{8,10,218-255} These data also suggest that fluid overload may dilute serum creatinine and thereby mask the diagnosis of AKI.^{95,256} These studies have led many to postulate that fluid overload directly impairs kidneys and other organ function,²⁵⁷⁻²⁵⁹ but whether fluid overload causes poor outcomes or this association is confounded by

severity of illness (i.e., sicker patients simply require larger volumes of fluid) remains unclear.

In this context, the concept of fluid restriction or deresuscitation has gained recognition in the critical care field. However, apart from possibly the setting of ARDS,⁹⁴ prospective data demonstrating the benefit of fluid restriction in the ICU are limited. Despite promising results of pilot studies and smaller trials,^{220,242,260-262} thus far large RCTs of fluid restriction in sepsis, major surgery, or other critical illness have not shown benefit. The 2018 Restrictive versus Liberal Fluid Therapy in Major Abdominal Surgery (RELIEF) trial randomized 3000 patients undergoing major abdominal surgery to two different fluid regimens and demonstrated that fluid restriction resulted in no benefit with increased rates of AKI and need for KRT.²⁶³ More recently, two large multicenter RCTs, the Conservative versus Liberal Approach to Fluid Therapy in Septic Shock (CLASSIC) and the Crystalloid Liberal or Vasopressors Early Resuscitation in Sepsis (CLOVERS) trials, analyzed the effects of fluid restriction in septic shock and demonstrated that fluid restriction was safe but failed to improve outcomes.^{264,265} Though additional large multicenter RCTs are ongoing (e.g., Australasian Resuscitation In Sepsis Evaluation: FLUID or vasopressors In emergency Department Sepsis [ARISE FLUIDS, NCT04569942] and Early Vasopressors in Sepsis [EVIS, NCT05179499] trials), high-quality evidence to guide the use of IV fluids in the ICU remains limited.

Nonetheless, guidelines based largely on observational data continue to emphasize the need for *early* resuscitation in septic shock.^{248,250,266-276} This has led to the conceptualization of four temporal phases of fluid therapy in sepsis or critical illness (Fig. 64.1).²⁷⁷⁻²⁷⁹ Consistent with this concept are trial data showing that fluid responsiveness disappears quickly, typically within hours, in patients with septic shock treated with modern resuscitation practices.²⁸⁰

Nephrologists are often tasked with helping achieve deresuscitation in patients with AKI requiring KRT. However, no large prospective studies exist to guide the optimal approach to the deresuscitation phase.²⁸¹⁻²⁸³ Observational data are conflicting, with some studies suggesting that the ability to achieve negative fluid balance with CRRT is associated with higher survival, primarily in patients with some degree of fluid overload at the time of CRRT initiation, and others finding that higher CRRT fluid removal rates are associated with increased mortality.^{233,284-289} Importantly, the gap between prescribed and achieved fluid removal with CRRT is an underrecognized parameter associated with increased mortality in patients with AKI and fluid overload.²⁸⁴ This lack of data has contributed to substantial worldwide practice variation in fluid management with KRT.²⁹⁰⁻²⁹³ Though pilot studies specifically analyzing approaches to ultrafiltration with CRRT are ongoing (NCT05306964, NCT05473143) and the use of point-of-care ultrasound and other novel technologies to guide fluid removal hold promise, additional data are clearly needed.^{294,295}

CHOICE OF INTRAVENOUS FLUIDS

CHOICE OF INTRAVENOUS FLUIDS: SALINE VERSUS BALANCED CRYSTALLOIDS

Observational²⁹⁶⁻³⁰⁰ and animal data^{301,302} suggest that 0.9% saline, when compared with balanced salt solutions such




Fig. 64.1 Four phases of fluid therapy in sepsis or other critical illness: “ROSE.” Optimal fluid resuscitation strategies should parallel the phases of septic shock or other critical illnesses. The first phase (e.g., first 6 hours) is **Resuscitation**, when initial aggressive resuscitation, recommended to be 30 mL/kg ideal body weight of crystalloid in most patients,²⁷⁶ should be provided. The second phase is **Optimization**, when additional fluids are given specifically targeting fluid responsiveness (i.e., increased stroke volume or cardiac output) or other specific hemodynamic endpoints to improve organ function. The third phase is **Stabilization**, also referred to as the “tone and time” phase, in which patients remain on vasopressors but vasopressor requirements are no longer increasing, patients are no longer fluid responsive, organ function is stable or improving, and further fluids are withheld. The fourth phase is **Evacuation**, also known as the deresuscitation or mobilization phase, when patients with volume overload are treated with diuretics or ultrafiltration to expedite the return to euvolemia in hopes of accelerating recovery from critical illness.²⁷⁷⁻²⁷⁹ The time at which a given patient transitions between phases varies, and repeated insults may substantially disrupt this sequence. (Reproduced with permission from ADQI (www.ADQI.org).)

as lactated Ringer (LR) or Plasma-Lyte (PL, Baxter), may increase the risk of AKI, need for KRT, and mortality. The pathophysiology underlying the nephrotoxicity of saline is thought to be related to the supraphysiologic chloride content and the resultant hyperchloremic acidosis that saline induces. Proposed mechanisms include reduced glomerular filtration rate via activation of the tubuloglomerular feedback triggered by increased chloride delivery to the macula densa, vasoconstriction caused by chloride-induced thromboxane release, and increased inflammatory cytokine expression induced by acidosis.³⁰¹⁻³⁰³ Imaging studies on human volunteers demonstrate that saline infusion reduces RBF.³⁰⁴

Since 2010, a series of large clinical trials have analyzed crystalloid choice in the ICU, starting with a 2012 before-and-after study and a 2015 cluster crossover RCT, which yielded conflicting results.^{305,306} In 2018, two large, single-center, pragmatic, unblinded, multiple-crossover cluster trials were published analyzing the use of LR or PL vs. saline, one in patients admitted from the ED to a non-ICU bed, the Saline against Lactated Ringer or Plasma-Lyte in the Emergency Department (SALT-ED) trial³⁰⁷ ($n > 13,000$), and another in ICU patients, the Isotonic Solutions and Major Adverse Renal Events Trial (SMART; $n > 15,000$).³⁰⁸ Both studies demonstrated that balanced solutions resulted in an approximate 1% reduction in the composite endpoint of major adverse kidney events (a composite of death, need for KRT, or persistent doubling of creatinine above baseline) by 30 days or discharge.

These two single-center studies were followed in 2021 and 2022 by two large multicenter RCTs. First, the Balanced Solutions in Intensive Care Study (BaSICS) in Brazil randomized >11,000 individual patients in 75 ICUs to the blinded use of PL or saline whenever possible throughout the entire ICU course and found no difference in 90-day mortality.³⁰⁹ There was also no difference in subgroups with sepsis or AKI, but the risk of death was significantly increased with LR in the subgroup with traumatic brain injury (TBI). Though methodologically rigorous, BaSICS was criticized for being underpowered despite the large sample size due to lower-than-predicted mortality, relatively low acuity of the subjects, administration of nonstudy fluids before-and-after randomization, and relatively low total volumes of fluids administered, which resulted in statistically significant but modest differences in serum chloride levels (e.g., 2–3 mEq/L higher in the saline arm at 48 hours).^{309,310} The subsequent Plasma-Lyte 148 versus Saline (PLUS) study, carried out in 53 ICUs in Australia and New Zealand, was similar in design, with the major exceptions being that the study excluded TBI and was aborted after enrollment of 5037 of the planned 8800 patients due to the COVID-19 pandemic.³¹¹ Otherwise, though PLUS also had some contamination by nonstudy fluid, the severity of acute illness and volumes of study fluid administered were somewhat higher than in the SMART or BaSICS trials. Like BaSICS, the primary outcome of 90-day mortality was not significantly different in the PL and saline arms. Soon after the publication of PLUS, a Bayesian meta-analysis of 13 RCTs including >35,000 ICU patients (of which >30,000 come from SMART, BaSICS, and PLUS) was published and concluded that the risk ratio for 90-day mortality with balanced crystalloids versus saline was 0.96 (95% confidence interval [CI], 0.91–1.01; $I^2=12.1\%$), with similar effects on risk of AKI and need for KRT.³¹² Large multicenter RCTs of PL versus saline in patients undergoing cadaveric renal transplantation ($n=808$) and in pediatric septic shock ($n=708$) have demonstrated benefits in kidney function with PL.^{313,314}

Though somewhat more expensive, there are no data (outside of TBI) to suggest harm from balanced solutions, and, as such, the use of balanced solutions seems reasonable in most ICU settings. Alternatively, one can reasonably interpret the BaSICS and PLUS trials as evidence that saline is largely safe in the ICU. Importantly, saline will continue to be the fluid of choice in patients with TBI or hypovolemic hypochloremic metabolic alkalosis.

CHOICE OF INTRAVENOUS FLUIDS: COLLOIDS VERSUS CRYSTALLOIDS

There are no compelling data to support the routine use of colloids over crystalloids as primary resuscitation fluid in ICU patients given a series of negative large RCTs^{315–319} and the increased expense of colloids. Two of these, the Saline versus Albumin Fluid Evaluation (SAFE)³¹⁵ and Albumin Italian Outcome Sepsis (ALBIOS)³¹⁹ trials, have shown albumin to be a safe alternative to crystalloids in ICU patients. Both trials suggested a possible benefit of lower mortality with albumin in subgroups with sepsis, via a nonsignificant trend in SAFE and a significant post hoc subgroup analysis in ALBIOS.^{315,319} Likewise, most^{320–322} (but not all³²³) meta-analyses have concluded that albumin use is associated with reduced mortality in sepsis. Additional RCTs evaluating the

role of albumin in the treatment of sepsis, such as albumin as endothelial rescue therapy, are ongoing (NCT03869385) or planned³²⁴ including the ALBIOS-BALANCED trial (NCT03654001).

On the basis of a statistically significant increase in mortality rate in the subgroup with TBI in SAFE,³¹⁵ albumin use should be avoided in TBI patients, for which saline is considered the fluid of choice. The use of artificial colloids, such as hydroxyethyl starch, gelatins, and dextrans, should be avoided in ICU patients due to an increased risk of AKI and a lack of evidence for their use.^{316–318,325,326}

USE OF INTRAVENOUS BICARBONATE IN THE INTENSIVE CARE UNIT

Metabolic acidosis is common in the ICU, present in 6% to 42% of patients, and, when persistent, is associated with a mortality of >50%.^{327,328} Using IV sodium bicarbonate to correct metabolic acidosis in the ICU, though commonly done, remains controversial.^{328–332} Despite the theoretical benefits of improved hemodynamic responsiveness to catechol vasopressors, animal models,^{333,334} human observational data,^{335,336} and small human trials^{337–339} suggest little to no benefit by attempting to correct acidemia with supplemental IV bicarbonate, with a possible signal in observational data for improved outcomes in patients with sepsis complicated by AKI.³³⁶ Potential harms of exogenous sodium bicarbonate administration include intracellular acidosis, cerebral spinal fluid acidosis, impaired oxygen offloading in tissues, stimulation of lactic acidosis, disinhibition of glycolysis, hypercarbia and/or depression of respiratory drive, hypokalemia, hypocalcemia, overshoot alkalosis, volume overload, and (when given in hypertonic solutions) hyponatremia.^{340,341}

The highest-quality data addressing the use of IV bicarbonate in the ICU available thus far come from the Sodium Bicarbonate to Treat Severe Acidosis in the Critically Ill (BICAR-ICU) trial.³⁴² The study randomized nearly 400 patients in 26 ICUs in France with severe metabolic or mixed acidosis ($\text{pH} \leq 7.20$, arterial $\text{pCO}_2 \leq 45$ mm Hg, serum bicarbonate ≤ 20 mmol/L) to unblinded treatment with 4.2% (500 mmol/L) sodium bicarbonate or standard care. Patients with pure respiratory acidosis, ketoacidosis, prior alkali therapy or KRT, stage IV CKD, and proven digestive or urinary tract loss of sodium bicarbonate were excluded, with the latter two criteria functionally excluding most non-anion gap metabolic acidosis. The primary outcome was the composite of death at 28 days or ≥ 1 organ failure at 7 days and was not statistically significant. However, patients in the subgroup with stage 2 or 3 AKI ($n=182$) had significant decreases in the primary outcome and both of its components, with a 28-day mortality of 46% vs. 63% ($P=0.017$) favoring sodium bicarbonate administration. In addition, there was a reduction in the need for KRT in the sodium bicarbonate group (35% vs. 52%). Though this subgroup analysis was prespecified and adjusted for multiple comparisons, the results remain hypothesis-generating. The findings of BICAR-ICU support the cautious use of sodium bicarbonate in critically ill patients with severe acidosis and AKI and the general avoidance of liberal sodium bicarbonate use in ICU patients without AKI.^{330–332} Larger studies are ongoing to better inform the use of sodium bicarbonate,

specifically in ICU patients with AKI including the BICAR-ICU-2 trial³⁴³ (NCT04010630) and the U.K. Multicentre Evaluation of Sodium Bicarbonate in Acute Kidney Injury in Critical Care (MOSAICC) trial (ISRCTN14027629).

KIDNEY REPLACEMENT THERAPY

The care of patients experiencing AKI is primarily supportive, with KRT required in about 5% to 10% of critically ill individuals with severe AKI.^{344,345} The goals of KRT in AKI include maintaining fluid balance, providing metabolic control, and preventing uremia-related complications, while simultaneously minimizing potential harm to the kidneys and facilitating renal recovery. KRT options include IHD, CRRT, hybrid therapies like prolonged intermittent renal replacement therapy (PIRRT), and peritoneal dialysis (PD). Despite these interventions, severe AKI requiring KRT is associated with a 50% to 60% mortality rate, prolonged hospital stay, and development of CKD and end-stage kidney disease.^{4,346-352}

TIMING OF KIDNEY REPLACEMENT THERAPY

Determining when to initiate KRT has been a subject of debate.³⁵³⁻³⁵⁵ In cases of severe AKI with medically refractory complications, initiating KRT is recommended to alleviate the deterioration of nonkidney organ function and reduce the risk of death. These urgent indications include severe hyperkalemia, severe metabolic acidosis, fluid overload resistant to diuretic therapy, signs of uremia, and specific drug intoxications.³⁵⁶⁻³⁵⁸ The specific criteria and thresholds for metabolic acidosis, hyperkalemia, or pulmonary edema vary widely in clinical practice.

Over the years, small, underpowered RCTs aimed to determine the optimal timing for initiating KRT, yielding mixed results.³⁵⁹⁻³⁶⁵ Since 2016, five well-designed RCTs addressed timing (Table 64.2). The trial Early versus Delayed Initiation of Renal Replacement Therapy on Mortality in Critically Ill Patients with Acute Kidney Injury (ELAIN) randomized 231 KDIGO stage 2 AKI patients to immediate or delayed KRT based on reaching KDIGO stage 3 AKI or developing life-threatening indications. The early KRT group had significantly lower 90-day mortality (39.3% vs. 54.7%) with a hazard ratio (HR) of 0.66 in favor of early KRT.³⁶⁶ In contrast, the Artificial Kidney Initiation in Kidney Injury (AKIKI),³⁶⁷ Initiation of Dialysis Early versus Delayed in the Intensive Care Unit (IDEAL-ICU),³⁶⁸ and Standard versus Accelerated Initiation of Renal Replacement Therapy in AKI (STARRT-AKI)³⁶⁹ trials failed to show an advantage to early initiation. The AKIKI trial enrolled 620 KDIGO stage 3 AKI patients, with no survival difference between early and delayed KRT.³⁶⁷ IDEAL-ICU, which included 448 AKI patients with septic shock, was terminated early due to no significant mortality difference.³⁶⁸ In the STARRT-AKI study, 2927 patients with KDIGO stage 2 or 3 AKI were randomized into an accelerated KRT group (initiation within 12 hours) or a standard group, where KRT was discouraged until AKI had persisted for more than 72 hours or an urgent indication developed (eTable 64.1).³⁶⁹ The study showed no 90-day mortality difference but more KRT-dependency and adverse events with early KRT.

The AKIKI2 study investigated whether adopting a strategy of delayed intervention, as initially defined in the first AKIKI trial, had inferior outcomes compared with an even more delayed approach. In this trial, 278 patients with oliguria for >72 hours or a BUN >112 mg/dL were randomized to immediate KRT initiation or a more delayed strategy, requiring a mandatory indication or a BUN >140 mg/dL.³⁷⁰ The primary outcome, days alive and free of KRT at day 28, did not differ significantly. However, multivariable analysis revealed a higher HR for death at 60 days with the more delayed strategy (1.65; 95% CI 1.09–2.50, $P = 0.018$).

The combined results from these trials indicate that, without specific indications, initiating KRT early does not offer clinical advantages and might impede renal recovery. Nonetheless, the AKIKI2 trial suggests there may be a threshold at which delaying KRT might result in adverse effects. The Acute Disease Quality Initiative workgroup on CRRT recommends a more personalized approach for acute KRT that does not rely solely on kidney function or AKI stage, but it also considers the patient's clinical condition (eTable 64.2).³⁵⁶ The workgroup proposes starting KRT when metabolic and fluid demands exceed kidney capacity, influenced by the severity of acute illness, solute load, fluid accumulation, and underlying chronic diseases.³⁵⁶ However, the specific components of the demand/capacity concept require further definition and validation.

KIDNEY REPLACEMENT THERAPY MODALITY

KRT modalities remove water and solutes by transport across a semipermeable membrane. Ultrafiltration, responsible for plasma water removal, operates through hydrostatic pressure (as in IHD, PIRRT, and CRRT) or an osmotic gradient (as in PD). Solute removal during KRT employs diffusion and/or convection. In extracorporeal treatments, certain solutes may also be eliminated by adsorption onto the membrane.

Convection, as occurs during ultrafiltration, involves the movement of solutes by bulk flow (Fig. 64.2A). When plasma water passes through the semipermeable membrane, it effectively “drags” small- and middle-molecular-weight solutes. Pore size and membrane charge determine the size of the solutes removed, and solute clearance is generally proportional to the ultrafiltration rate. In contrast, diffusion drives solutes across the membrane via concentration gradients between blood and dialysate (Fig. 64.2B). Running the dialysate countercurrent to blood flow maximizes this gradient. Smaller solutes like urea diffuse more effectively than larger ones such as β_2 -microglobulin, as solute diffusivity is inversely proportional to molecular weight. All KRT modalities utilize ultrafiltration for fluid removal and diffusion, convection, or both for solute clearance.

INTERMITTENT HEMODIALYSIS

IHD sessions are typically prescribed three times weekly, lasting 3 to 4 hours each, with blood flow rates of 300 to 500 mL/min and dialysate flow rates of 500 to 800 mL/min (Table 64.3). Ultrafiltration removes excess fluid, and solute clearance mainly relies on diffusion, with some larger solutes cleared via convection due to increased membrane porosity and internal filtration and backfiltration.³⁷¹ The frequency and duration of dialysis depend on patient metabolic control, volume status, and hemodynamic stability. Initial

Table 64.2 Randomized Controlled Trials Evaluating Timing of Kidney Replacement Therapy Initiation for Critically Ill Adult Patients With Acute Kidney Injury

	ELAIN (2016)³⁶⁶	AKIKI (2016)³⁶⁷	IDEAL-ICU (2018)³⁶⁸	STARRT-AKI (2020)³⁶⁹	AKIKI-2 (2021)³⁷⁰
Setting	Single center in Germany	31 Centers in France	29 Centers in France	168 Centers in 15 countries	39 Centers in France
N	231	620	488	2927	278
Inclusion criteria	KDIGO stage 2 AKI	KDIGO stage 3 AKI	RIFLE FAILURE stage AKI	KDIGO stage 2 or 3 AKI	KDIGO stage 3 with oliguria >72 h or BUN 112-140 mg/dL
Timing of earlier KRT	Within 8 h	Within 6 h	Within 12 h	Within 12 h	Within 12 h
Timing of later KRT	Within 12 h of KDIGO stage 3 AKI or “absolute indications” ^a	Life-threatening complications of AKI ^b , BUN >112 mg/dL, or oliguria >72 h	48 h post-randomization in the absence of kidney recovery	Life-threatening complications of AKI ^c or AKI persisting >72 h	Life-threatening complications of AKI ^b or BUN >140 mg/dL
Median time to KRT (earlier vs. later)	6 h vs. 25.5 h	2 h vs. 57 h	7.6 h vs. 51.5 h	6.1 h vs. 31.1 h	3 h vs. 33 h
% who received KRT (earlier vs. later)	100% vs. 91%	98% vs. 51%	97% vs. 62%	97% s. 62%	98% vs. 79%
Primary outcome (earlier vs. later)	90-day mortality: 39.3% vs. 54.7% (<i>P</i> = 0.03)	60-day mortality: 48.5% vs. 49.7% (<i>P</i> = 0.79)	90-day mortality: 58% vs. 54% (<i>P</i> = 0.38)	90-day mortality: 43.9% vs. 43.7% (<i>P</i> = 0.92)	KRT-free days within 28 days: 12 vs. 10 days (<i>P</i> = 0.93)
Other key outcomes	Shorter KRT duration and hospital length of stay in early arm	Spontaneous diuresis occurred earlier in delayed arm	No difference in ICU or hospital length of stay	KRT dependence at 90 days higher in early arm (10% vs. 6%, 95% CI 1.2-2.4)	In prespecified adjusted analysis, more delayed strategy was associated with mortality (<i>P</i> = 0.02)
KRT-related complications	No difference	More catheter-related blood stream infections in early arm	More hyperkalemia in delayed arm	More adverse events ^d (17% vs. 23%, <i>P</i> < 0.001) in early arm	No difference
Limitations	Limited generalizability (smaller sample, single center, mostly surgical)	55% of patients were treated initially with IHD	Stopped early due to futility	Decision for KRT initiation ultimately left to provider discretion	Small sample size

^aIndications for KRT initiation in the delayed arm in ELAIN included BUN >100 mg/dL, potassium >6 mEq/L or hyperkalemia-induced ECG changes, magnesium >9.7 mg/dL, anuria or oliguria with urine output <200 mL per 12 h, or organ edema resistant to ≥1 dose of loop diuretic.

^bIndications for KRT initiation in the delayed arm in AKIKI and AKIKI-2 included potassium >6 mEq/L, potassium >5.5 mEq/L despite medical therapy, metabolic acidosis with pH <7.15, and pulmonary edema resulting in hypoxemia despite diuretic therapy.

^cIndications for KRT initiation in the delayed arm in STARRT-AKI included potassium ≥6.0 mEq/L, pH ≤7.20 or serum bicarbonate ≤12 mEq/L, and volume overload with hypoxemia.

^dIn STARRT-AKI, the early arm experienced statistically significant increases in hypotension and hypophosphatemia, trends for more bloodstream infections and arrhythmias (*p* = 0.07 for both), but no significant difference in serious adverse events (*P* = 0.15).

BUN, Blood urea nitrogen; CI, confidence interval; IHD, intermittent hemodialysis; KDIGO, Kidney Disease: Improving Global Outcomes;

sessions typically use lower blood and dialysate flows for patients susceptible to dialysis disequilibrium.³⁷²

IHD is preferred for rapid correction of electrolyte imbalances and drug intoxications due to its rapid solute removal. However, IHD can lead to intradialytic hypotension in 10% to 70% of acute treatments.³⁷³⁻³⁷⁷

Proposed interventions for improving hemodynamic stability are mostly extrapolated from patients on maintenance IHD. These interventions include decreasing the ultrafiltration rate, lowering the blood flow rate, sodium modeling, cooling the dialysate, increasing the dialysate calcium concentration, using isolated ultrafiltration, using

eTable 64.1 Summary of Standard versus Accelerated Initiation of Renal Replacement Therapy in Acute Kidney Injury (STARRT-AKI) Trial³⁶⁹**Setting:**

- Multicenter: 168 hospitals in 15 countries in 5 continents
- October 2015 to September 2019

Key inclusion criteria:

- ICU admission
- Severe AKI (KDIGO stage 2 or 3)
- Need for clinical equipoise (i.e., the investigator felt that neither initiation nor deferral of KRT was mandated)^a

Study design and study arms:

- Open-label RCT with 1:1 allocation
- Accelerated strategy (KRT initiated within 12 hours of meeting eligibility): $n = 1462$
- Standard care (KRT discouraged unless specific criteria met): $n = 1465$

Patient characteristics at randomization:

- Mean age 65
- 32% female
- 44% with underlying chronic kidney disease
- 67% medical ICU admission
- 58% with sepsis
- 77% requiring mechanical ventilation
- 70% requiring vasoactive support
- Mean serum creatinine 3.5 mg/dL

Initial KRT modality (among KRT-treated patients)

- 69% CRRT
- 26% IHD
- 4.4% SLED

Primary outcome (accelerated arm vs. standard arm):

- No difference in 90-day mortality by modified intention-to-treat analysis (43.7% vs. 43.9%, RR 1.0, $P = 0.92$)

Key secondary outcomes (all reported as accelerated arm vs. standard arm):

- 96.8% vs. 61.8% initiated KRT
- Higher rate of KRT dependence at 90 days of 10.4% vs. 6.0% (RR 1.74, 95% CI 1.24-2.43)
- Higher rate of adverse events (AEs) of 23.0% vs. 16.5% ($P < 0.001$), with difference driven by the 2 most common AEs, hypotension and hypophosphatemia
- Higher rate of readmission at 90 days of 20.9% vs. 17.0% (RR 1.23, 95% CI 1.02-1.49)
- No difference in serious AEs (1.0% vs. 0.5%, $P = 0.15$)
- No difference in primary outcome in 6 prespecified subgroups: gender, baseline eGFR, severity of illness (i.e., SAPS II score), presence or absence of sepsis, surgical or medical ICU admission, and geographic region

^aOf >23,000 patients screened that otherwise met inclusion criteria, >7,800 were excluded on basis of lack of equipoise.

CI, Confidence interval; CRRT, continuous renal replacement therapy; eGFR, estimated glomerular filtration rate; ICU, intensive care unit; IHD, intermittent hemodialysis; KDIGO, Kidney Disease: Improving Global Outcomes; KRT, kidney replacement therapy; RCT, randomized controlled trial; RR, relative risk; SAPS II, Simplified Acute Physiology II score; SLED, sustained low-efficiency dialysis.

eTable 64.2 Factors to Consider When Deliberating Timing of Kidney Replacement Therapy (KRT) Initiation for Acute Kidney Injury (AKI) Patients in the Intensive Care Unit (ICU)

Illness Severity and Trajectory

- AKI severity and trend
- Fluid balance and symptoms of fluid overload
- Severity of electrolyte and acid-base disorders
- Presence of relevant nonrenal organ dysfunction impacted by AKI and/or fluid overload
- Anticipated need for enhanced metabolic support (e.g., rhabdomyolysis or tumor lysis syndrome)

Necessity of KRT

- Presence of oliguria, considering the effect of diuretics
- Likelihood of recovery of kidney function without KRT, considering the reversibility of the specific etiology of AKI, trends in kidney function and urine output, and baseline kidney function
- Underlying comorbidities impacted by AKI and/or fluid overload

Risks of KRT

- Hemodynamic instability from KRT
- Infection, including catheter-related bloodstream infection
- Other risks associated with vascular access (pneumothorax, procedural bleeding, catheter-associated deep venous thrombosis, etc.)
- Clearance of trace elements, water-soluble vitamins, phosphate, amino acids/small peptides, and drugs (especially antibiotics)
- Delayed renal recovery
- Increased risk of immobilization / impeding physical rehabilitation

Other Factors

- Patient and family wishes and overall goals of care
- Overall prognosis including likelihood of patient survival
- Availability of machines, disposable supplies, and nursing staff
- Health care costs

Adapted from Macedo E, Mehta RL. Continuous dialysis therapies: core curriculum 2016. *Am J Kidney Dis.* 2016;68(4):645–657 and Ostermann M, Joannidis M, Pani A, et al. Patient selection and timing of continuous renal replacement therapy. *Blood Purif.* 2016;42(3):224–237.

hypertonic infusions, extending the duration of dialysis or increasing the frequency of treatments in order to slow the ultrafiltration rate, and using midodrine.³⁷⁸⁻³⁹⁰ In general, these interventions lack robust evidence, especially in critically ill patients with AKI. Additionally, the rapid shifts of intracellular fluid and solutes during IHD can result in increased ICP and cerebral edema in patients with acute brain injury.³⁹¹

CONTINUOUS RENAL REPLACEMENT THERAPY

CRRT offers an alternative for managing hemodynamically unstable patients with AKI intolerant to IHD. CRRT uses diffusion, convection, or a combination to remove solutes and fluids gradually. This method theoretically allows for better hemodynamic tolerance compared with IHD by providing slower solute and fluid removal per unit time. CRRT is generally performed 24 hours a day with blood flow rates typically between 100 and 300 mL/min (Table 64.3). Advantages of CRRT include better hemodynamic tolerance, continuous solute clearance, and improved volume control (Fig. 64.3).²²³ Drawbacks include vascular access challenges, filter clotting, increased anticoagulation needs, reduced patient mobility, and higher costs and ICU nurse workload compared with IHD. Although rare, dialysis disequilibrium has been reported with CRRT.^{392,393}

CRRT modalities for solute removal are continuous venovenous hemofiltration (CVVH), continuous venovenous hemodialysis (CVVHD), and continuous venovenous hemodiafiltration (CVVHDF). CVVH relies on convection without dialysate, using a hemofiltration solution (often referred to as “replacement fluid” based on historical practices) infused before the filter (prefilter) or after the filter (postfilter) to restore volume and electrolytes. When using postfilter hemofiltration solution, the filtration fraction should be kept <0.25 to reduce the likelihood of clotting of the hemofilter.³⁹⁴ This can be done by increasing the blood flow rate³⁹⁵ or introducing some hemofiltration solution before the filter, although the latter will dilute the concentration of solutes entering the hemofilter, leading to a decrease in solute clearance.^{396,397} CVVHD primarily uses diffusion with slower dialysate flow rates, resulting in near-complete saturation of the dialysate. Ultrafiltration is primarily used for volume control. CVVHDF combines convection and diffusion, utilizing both dialysate and replacement fluid.

Despite enhanced middle-molecule clearance in convective therapies, meta-analyses comparing hemofiltration to hemodialysis for AKI found no significant differences in patient survival.³⁹⁸ Although high-volume hemofiltration (HVHF) with ultrafiltration rates >50 mL/kg/h has been used to enhance cytokine clearance in AKI patients with sepsis, controlled studies have not consistently demonstrated better outcomes.^{30,364,399,400} With no clear advantage of one CRRT modality over another, choice should be based on clinician preference and expertise, with attention to circuit survival.

PROLONGED INTERMITTENT RENAL REPLACEMENT THERAPY

Prolonged intermittent renal replacement therapy (PIRRT), an alternative to CRRT for critically ill patients

with AKI, offers a hybrid approach.⁴⁰¹⁻⁴⁰³ These techniques mitigate the hemodynamic impact of IHD by extending the duration of therapy and adapting conventional hemodialysis machines to operate at lower blood pump speeds (100–300 mL/min) and dialysate flow rates (100–300 mL/min) (Table 64.3). Alternatively, CRRT equipment can be adapted for PIRRT, with sessions lasting 6 to 18 hours. PIRRT practices vary substantially in technology, prescription, and anticoagulation.^{403,404} Diffusion-based PIRRT is described by terms like “sustained low-efficiency (daily) dialysis” (SLED or SLEDD), “extended daily dialysis” (EDD), “slow continuous dialysis” (SCD), and “go slow dialysis.” When combining convection and diffusion, these therapies are known as low-efficiency daily diafiltration (SLEDD-f) and sustained hemodiafiltration (S-HDF). “Accelerated venovenous hemofiltration” is used for exclusively convective therapy.⁴⁰⁴⁻⁴⁰⁷

PIRRT combines advantages of CRRT and IHD, offering improved hemodynamic stability by gradually removing solutes and fluids over an extended timeframe. It eliminates the need for 24-hour therapy and costly CRRT equipment and solutions. PIRRT provides high solute clearances similar to IHD, allowing for uninterrupted diagnostic and therapeutic procedures.^{402,404} It can also be administered overnight, promoting patient mobility during the day. Studies confirmed hemodynamic control comparable with CRRT.^{404,408-411} In some medical institutions, PIRRT is employed to transition patients from CRRT to conventional IHD when their hemodynamic stability improves.^{354,404}

PERITONEAL DIALYSIS

PD leverages the peritoneum as a semipermeable membrane for solute removal by diffusion. A catheter is used to instill a dialysate solution into the peritoneal cavity, where it dwells for a specified duration, allowing solutes to diffuse from the blood into the dialysate before drainage. The dialysate contains varying high-glucose concentrations to establish an osmotic gradient for fluid removal. Acute PD can be conducted intermittently or continuously and either manually or through an automated cycler. PD techniques for AKI encompass high-volume PD, acute intermittent peritoneal dialysis (AIPD), continuous equilibrated peritoneal dialysis (CEPD), tidal peritoneal dialysis (TPD), and continuous flow peritoneal dialysis (CFPD).

High-volume PD employs an automated cycler to deliver 18 to 22 exchanges of 2 L every 24 hours, offering the highest small-solute clearance and ultrafiltration rate among PD methods.^{412,413} AIPD employs brief dwell times with dialysate flows of 1 to 4 L/hour for 12 to 24 hours every other day. CEPD resembles continuous ambulatory PD for chronic outpatient PD but without patient ambulation.^{414,415} In TPD, a portion of dialysate remains in the peritoneal cavity, facilitating rapid exchanges through a cycler, and improving solute clearance.⁴¹⁶ CFPD involves simultaneous inflow and outflow of dialysate through two routes, maintaining a fixed intraperitoneal volume, with high dialysate flow rates and urea clearances of 30 to 50 mL/min.^{417,418}

PD offers advantages like technical simplicity, hemodynamic stability, reduced dialysis disequilibrium risk, no need for anticoagulation or vascular access, and cost-effectiveness.^{419,420} Gradual solute and fluid




Fig. 64.2 Modalities of kidney replacement therapy use primarily convective clearance (hemofiltration), diffusive clearance (hemodialysis), or both. (A) In hemofiltration, solute clearance occurs primarily by convection. In convection, solutes are transported across the hemofilter membrane along with plasma water as a result of hydrostatic pressure (i.e., transmembrane pressure) generated on the blood side of the membrane. Solute cleared by convection include urea and other small molecules along with larger “middle molecules.” (B) In hemodialysis (HD), solute clearance occurs primarily by diffusion, which is driven by a concentration gradient across the semipermeable membrane. Small solutes in high concentration in the blood diffuse across the membrane into the dialysate, which contains either little (e.g., potassium) or none (e.g., urea) of the solutes being cleared. Small solutes in higher concentration in the dialysate (e.g., bicarbonate) diffuse into the blood. Dialysate runs across the HD membrane countercurrent to the direction of blood flow to maintain a concentration gradient for removal of small solutes along the entire length of the semipermeable membrane. Modern hemodialyzers are virtually all “high-flux” dialyzers, which clear substances larger than historical low-flux dialyzers. However, unlike hemofiltration, HD does not effectively clear larger middle molecules. Ultrafiltration can be performed with HD by applying a transmembrane pressure across the membrane, but, in contrast to the high volume of ultrafiltration used to achieve significant solute clearance in hemofiltration, the volumes of ultrafiltration performed in HD are relatively small, contribute little to solute clearance, and are instead used only to achieve net volume removal. KRT, Kidney replacement therapy. (Reproduced with permission from Teixeira JP, Neyra JA, Tolwani A. Continuous KRT: a contemporary review. *Clin J Am Soc Nephrol.* 2023;18[2]:256–269.)

removal makes it safer for patients at risk of intracranial hypertension. Drawbacks include specialized catheter training, catheter placement complications, peritonitis risk, potential solute inadequacy, unpredictable ultrafiltration, hyperglycemia, protein loss, and respiratory compromise due to intraabdominal pressure.^{419,420} PD is contraindicated for individuals with recent abdominal surgery, abdominal drains, diaphragmatic disease, or ileus.^{419,420}

SELECTION OF KIDNEY REPLACEMENT THERAPY MODALITY

While CRRT is often considered standard of care for hemodynamically unstable patients needing KRT, no specific KRT modality has been proven to be superior in AKI.⁴²¹ Existing data from RCTs^{422–427} and meta-analyses^{428–434} generally regard CRRT and IHD as complementary therapies, with comparable outcomes including mortality and kidney recovery.^{15,435} Some trials suggest improved hemodynamic tolerance with CRRT, but others report no significant differences in hypotension rates or vasoactive support.^{422,424,427} PIRRT also exhibits comparable outcomes with CRRT regarding mortality, renal recovery, ICU length of stay, and fluid removal.^{409–411,436–438} However, phosphate

removal has been noted to be higher in PIRRT compared with CRRT.⁴³⁹

Small RCTs support PD as an effective AKI treatment with outcomes akin to IHD and CRRT in terms of efficacy and survival.^{440–442} A Cochrane review involving 484 patients with AKI requiring dialysis found no significant differences in mortality, recovery of kidney function, or infectious complications between PD and other KRT modalities.⁴⁴² In a subsequent trial, 125 critically ill patients with AKI were randomly assigned to high-volume tidal PD or CRRT. At 28 days, PD-treated patients had higher survival rates (70% vs. 47%, $P < 0.01$) and renal recovery rates (60% vs. 37%, $P < 0.01$), along with fewer infectious complications and shorter ICU stays.⁴⁴³ Overall, acute PD is valuable in patients with hemodynamic instability and vascular access challenges or in places with limited KRT resources.

In clinical practice, hemodynamically stable patients often receive IHD, while hemodynamically unstable, fluid-overloaded, or septic patients receive CRRT or PIRRT.^{15,444–446} CRRT might improve volume management more effectively than conventional IHD²²³ yet appears comparable with PIRRT.⁴³⁷ CRRT preserves cerebral perfusion, making it preferable for patients at risk of cerebral edema.^{447,448} Additionally, CRRT

Table 64.3 Comparison of Kidney Replacement Therapy Modalities

	Intermittent Hemodialysis (IHD)	Prolonged Intermittent Renal Replacement Therapy (PIRRT)^a	Continuous Renal Replacement Therapy (CRRT)
Typical parameters	- Duration: 3–4 h - Frequency: 3–4 times per week Q_b: 300–400 mL/min Q_d: N/A Q_r: 600–800 mL/min UF_{NET} rate: 0–10 mL/kg/h	- Duration: 6–12 h - Frequency: 3–6 times per week Q_b: 200–300 mL/min Q_d: N/A Q_r: 200–300 mL/min UF_{NET} rate: 0–8 mL/kg/h	- Duration/frequency: continuous Q_b: 100–300 mL/min Q_r: none or 16–50 mL/min Q_d: none or 16–50 mL/min UF_{NET} rate: 0–2 mL/kg/h
Advantages	- Widespread availability - Low cost of disposables - Delivery with no anticoagulation usually feasible - Most rapid instantaneous clearance - May be performed with catheter or arteriovenous fistula or graft - Lowest rate of hypophosphatemia - Time off KRT allows more time for physical therapy relative to CRRT	- Low cost of disposables - Delivery with no anticoagulation usually feasible - Likely improved hemodynamic tolerance and fluid removal relative to IHD - May be performed with catheter or arteriovenous fistula or graft - Time off KRT allows more time for physical therapy relative to CRRT	- Likely best hemodynamic stability - Most effective for fluid management - Least likely to exacerbate intracranial hypertension - Superior solute control (i.e., higher total daily or weekly dose) - Usually permits nutrition without restriction in protein, phosphate, or potassium - Facilitates gradual controlled correction of dysnatremia - Possibly increased renal recovery among survivors^b
Disadvantages	- Likely highest risk of hypotension - Likely highest risk of exacerbating intracranial hypertension - Limited fluid removal - Often requires limitation in dietary potassium or phosphate intake	- Least standardization of prescription - Lack of large randomized controlled trials validating optimal prescription or comparing efficacy to IHD or CRRT - Limited data on drug dosing, especially antimicrobials - Limited data on safety in acute brain injury	- High cost of disposables/highest overall cost - Lowest instantaneous solute clearance - Greatest need for anticoagulation - Highest risk of hypophosphatemia - Requires catheter placement^c - Increased risk of immobilization^d

^aPIRRT encompasses a variety of hybrid modalities such as sustained low-efficiency dialysis (SLED), accelerated venovenous hemofiltration (AVVH), and many others; note that the typical PIRRT parameters shown are for SLED and that the listed advantages and disadvantages may not apply equally to all types of PIRRT.

^bThe increased rate of renal recovery noted in survivors of AKI initially treated with CRRT relative to IHD is based on observational data only and therefore controversial.

^cAlthough one single-center observational study⁵¹⁷ suggests that using arteriovenous fistulae/grfts for CRRT may be safe and feasible, doing so is not standard of care in most institutions.

^dCRRT is not a contraindication to early mobilization per se as observational data⁵¹⁸ suggest that physical rehabilitation is feasible and safe in patients on CRRT, but many providers perceive CRRT to be a barrier to physical therapy.

Q_b , blood flow rate; Q_d , dialysate flow rate; Q_r , replacement fluid rate; UF_{NET} , net ultrafiltration rate.

offers the unique advantage of gradual and precise correction of dysnatremias.^{449,450} It is also suitable for conditions requiring ongoing solute control (e.g., severe rhabdomyolysis and tumor lysis syndrome). In contrast, IHD is preferred for rapid solute removal, as in severe hyperkalemia or drug intoxications.³⁹⁵ PIRRT and IHD allow intermittent treatment breaks, facilitating physical therapy and early mobilization. Ultimately, the selection of KRT modality should consider available resources, cost, provider expertise, and patient needs.

PRESCRIBING KIDNEY REPLACEMENT THERAPY

VASCULAR ACCESS

According to the 2012 KDIGO AKI guidelines, the preferred site for temporary dialysis catheter placement is the right internal jugular (IJ) vein, followed by the femoral vein,

followed by the left IJ, with the subclavian veins as a last resort.¹⁵ The catheter should be inserted with the tip in a large vein at an appropriate depth: 15 to 20 cm for right IJ catheters and 20 to 24 cm for left IJ catheters with a tip at the cavoatrial junction and at least 20 to 25 cm for femoral catheters with a tip in the inferior vena cava.^{451,452} Ultrasound guidance and strict aseptic technique are recommended to minimize complications.¹⁵ While there is consensus favoring the right IJ, some argue for left IJ over femoral placement, especially in obese patients at higher risk of femoral catheter complications.⁴⁵³ In an RCT comparing femoral and jugular catheters, infection rates were similar overall but higher in the femoral group for patients with a BMI >28.4 kg/m².⁴⁵³ In a post hoc analysis, right IJ placement had the lowest catheter malfunction rate, followed by femoral, with highest in the left IJ, though the malfunction of the left IJ catheters in this study may have been due to inadequate depth of




Fig. 64.3 Superior volume control with CRRT versus intermittent hemodialysis (IHD). (A) Fluid accumulation over time in patients on CRRT and on IHD. Data from the PICARD (Program to Improve Care in Acute Renal Disease) study. (B) Twenty-four-hour total fluid balance (I/Os; milliliters) during the first 3 days of kidney replacement therapy in patients on IHD and CVVHD therapy. Shown are median values with interquartile range (box borders) and extreme values (whiskers). CRRT, Continuous renal replacement therapy; CVVHD, continuous venovenous hemodialysis; IHD, intermittent hemodialysis. (A reproduced with permission from Bouchard J, Soroko SB, Chertow GM, et al. Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury. *Kidney Int.* 2009;76[4]:422–427. B reproduced with permission from Augustine JJ, Sandy D, Seifert TH, Paganini EP. A randomized controlled trial comparing intermittent with continuous dialysis in patients with ARF. *Am J Kidney Dis.* 2004;44[6]:1000–1007.)

insertion (16 cm).^{451,454} Observational studies suggest lower infection and thrombosis risks with tunneled catheters, along with better blood flow rates.^{377,455} Therefore considering a tunneled dialysis catheter over a nontunneled one should be considered for prolonged KRT requirements, but further research is needed before making it a standard practice for AKI in the ICU.^{377,456}

DOSE

KRT dosing in AKI is determined by Kt/V_{urea} and frequency in IHD and PIRRT, as well as effluent flow rate (mL/kg/h) in CRRT. Two large RCTs, the VA/NIH ATN study and the RENAL study, evaluated KRT dosing in AKI. In the ATN trial, intensive KRT included high-dose CVVHDF or PIRRT for unstable patients and IHD for stable patients. In the intensive approach, patients received CVVHDF with an effluent rate of 35 mL/kg/h or six PIRRT/IHD sessions per week, with each session targeting a Kt/V_{urea} of 1.2 to 1.4.⁴⁵⁷ The less intensive approach used CVVHDF with an effluent rate of 20 mL/kg/h or three PIRRT/IHD treatments per week with the same Kt/V_{urea} target. Notably, the intensive KRT strategy did not reduce 60-day mortality or improve kidney recovery. The RENAL trial compared CRRT at doses of 40 and 25 mL/kg/h and revealed no survival advantage with higher CRRT intensity at 90 days.⁴⁵⁸ Multiple meta-analyses also showed no survival benefit with increased KRT intensity.^{459–461} Studies of high-volume CRRT with doses up to 70 to 85 mL/kg/h have similarly shown no benefit.^{30,364}

On the basis of these studies, KDIGO 2012 AKI guidelines recommend a CRRT-delivered dose of 20 to 25 mL/kg/h and a Kt/V_{urea} of ≥ 1.2 per session for PIRRT or IHD.^{15,446} In IHD, continued patient assessment is needed to ascertain whether additional treatments are required to maintain electrolyte and acid-base balance and control volume status.⁴⁴⁶ For CRRT, doses of 25 to 30 mL/kg/h

can be considered to account for therapy interruptions.¹⁵ Temporary use of higher doses may be required for specific conditions including hyperkalemia, severe acidosis, or hyperammonemia, but dose should be adjusted back to standard ranges once solute control is achieved.³⁹⁵ CRRT dosing should remain adaptable and continuously adjusted to meet the dynamic needs of critically ill patients.

High-intensity KRT can have adverse consequences by removing beneficial solutes, such as essential nutrients, antiinflammatory cytokines, and antibiotics.^{15,462} Both the ATN and RENAL studies observed increased rates of hypophosphatemia with higher KRT intensity.^{457,458} In the ATN study, higher-intensity KRT was associated with lower rates of successful extubation and fewer ventilator-free days, especially in patients with lower serum phosphate levels.⁴⁶³ Higher KRT intensity also increased the risk of a $\geq 50\%$ decline in urine output (HR = 1.29; 95% CI, 1.1–1.51).⁴⁶⁴ Patients treated only with IHD in the ATN study had more KRT-free days with lower-intensity KRT.⁴⁶⁵ A meta-analysis of six trials involving 1926 subjects found that higher-intensity KRT was associated with longer KRT duration and a higher likelihood of KRT dependence at day 28 (HR = 1.15, 95% CI 1–1.33).⁴⁶¹ Additionally, high-intensity KRT is costlier due to expenses for disposable items and nursing labor.^{462,466} The threshold for harmful underdosing in KRT intensity is uncertain. Some experts suggest a potential mortality risk below 20 mL/kg/h, but observational data from Japan suggest that doses as low as 15 mL/kg/h are safe.⁴⁶⁷

ANTICOAGULATION

KDIGO 2012 guidelines for AKI recommend using regional citrate anticoagulation (RCA) over heparin in CRRT unless contraindicated.¹⁵ Citrate chelates ionized calcium (iCa) in the extracorporeal circuit, and optimal anticoagulation is achieved when hemofilter iCa is <0.4 mmol/L.⁴⁶⁸ A portion

of the calcium-citrate complex is lost across the hemofilter while the rest enters the systemic circulation, where the liver metabolizes citrate to bicarbonate and calcium is released into the circulation.⁴⁶⁹ Calcium is infused into the patient to replace the calcium lost across the hemofilter.

Proactive management of RCA is essential to mitigate potential side effects including metabolic alkalosis and acidosis, hypocalcemia and hypercalcemia, and hypomagnesemia.⁴⁶⁹⁻⁴⁷² Patients with severe shock liver and lactic acidosis may not effectively metabolize citrate.^{469,473,474} Citrate accumulation is characterized by low systemic iCa, elevated total calcium, total calcium to systemic iCa ratio >2.5, anion gap metabolic acidosis, and escalating calcium infusion requirements (Table 64.4).^{472,475-477} Nonetheless, RCA has been safely used in advanced liver disease and perioperative liver transplant patients.⁴⁷⁸⁻⁴⁸⁰

Multiple RCTs including the multicenter Effect of Regional Citrate Anticoagulation versus Systemic Heparin Anticoagulation During Continuous KRT on Dialysis Filter Life Span and Mortality Among Critically Ill Patients With Acute Kidney Injury (RICH) trial support RCA's superiority over systemic heparin in extending filter life and reducing bleeding risk.⁴⁸¹⁻⁴⁸³ In the RICH trial, exploratory secondary outcomes demonstrated a potential association between citrate and an increased risk of infection.^{482,483} However, a post hoc analysis found that filter lifespan exceeding 48 h, but not anticoagulation choice, was associated with higher infection rates.⁴⁸⁴ Further research is needed to clarify this observation.

Anticoagulation-free KRT sessions are feasible with IHD and PIRRT due to the higher blood flow and shorter durations. CRRT without anticoagulation may require more circuit changes due to clotting, but strategies such as higher blood flow rates and prefilter replacement solution may

mitigate clotting risk.⁴⁸⁵⁻⁴⁸⁸ In one RCT, different blood flow rates (150 vs. 250 mL/min) in CRRT had no significant impact on filter clotting, possibly due to higher flow rates triggering more pressure alarms and reductions or stoppages in blood pump speed, with consequent increased clotting risk.⁴⁸⁹

ULTRAFILTRATION

Achieving euvolemia is crucial in critical illness, especially due to the harms of fluid overload.^{223,254,490} Ultrafiltration and rapid osmotic shifts during IHD can cause intradialytic hypotension from inadequate plasma refill and impaired hemodynamic compensatory mechanisms.³⁷⁷ Additionally, IHD can induce hypotension due to "myocardial stunning," unrelated to ultrafiltration, observed in both patients on maintenance IHD and those with AKI undergoing IHD.^{491,492} Surprisingly, myocardial stunning was also seen in one study of patients with AKI on CRRT, suggesting microcirculatory factors might play a role.⁴⁹³

Data on prescribing net ultrafiltration (UF_{NET}) in CRRT are limited. UF_{NET} is the net volume of fluid removed per hour adjusted for patient body weight (mL/kg/h).⁴⁹⁴ In a secondary analysis of the RENAL trial, UF_{NET} rates above 1.75 mL/kg/h were associated with lower survival and a higher risk of hypophosphatemia.²⁸⁵ Accumulating observational data suggest a nonlinear relationship between UF_{NET} and mortality, with the lowest mortality at UF_{NET} rates of 1.0 to 1.75 mL/kg/h (Fig. 64.4).^{285,494} Iterative assessment of patient's fluid balance (desired vs. achieved) and tolerance to fluid removal is key for risk stratification.^{284,495} Prospective studies are needed to validate the proactive targeting of UF_{NET} within the 1.0 to 1.75 mL/kg/h range and/or other methods to achieve personalized fluid balance goals.³⁹⁵

Table 64.4 Metabolic Complications of Regional Citrate Anticoagulation Related to Excess or Inadequate Citrate

Complication	Mechanism	Diagnostic Features	Management Options
Excess buffer from citrate	Excess citrate supply, resulting in excessive metabolic conversion of citrate to bicarbonate and excess buffer	- Metabolic alkalosis $tCa/iCa < 2.5$	- Decrease blood flow rate/citrate rate^a - Increase dialysate rate - Decrease buffer concentration in other CRRT solutions
Citrate accumulation ^b	Impaired metabolic conversion of citrate resulting in accumulation of citrate-calcium complexes in blood	- Anion gap metabolic acidosis $tCa/iCa > 2.5$ - Escalating calcium infusion requirements	- Decrease blood flow rate/citrate rate^a - Increase dialysate rate - Discontinue regional citrate anticoagulation
Insufficient buffer from citrate	Inadequate citrate supply, resulting in insufficient conversion of citrate to bicarbonate and insufficient buffer	- Metabolic acidosis $tCa/iCa < 2.5$	- Increase blood flow rate/citrate rate^a - Decrease dialysate rate - Increase buffer concentration in other CRRT solutions

^aThese changes in blood flow and citrate rate are done proportionally, to increase or decrease the delivery of citrate while maintaining the same ratio of citrate infusion rate to blood flow rate and thus similar intrafilter ionized calcium concentration.

^bCitrate accumulation is also known as citrate toxicity or citrate lock.

CRRT, Continuous renal replacement therapy; iCa, ionized calcium; tCa, total calcium.

Adapted with permission from Neyra JA, Yessayan L, Thompson Bastin ML, et al. How to prescribe and troubleshoot continuous renal replacement therapy: a case-based review. *Kidney360*. 2021;2(2):371–384.




Fig. 64.4 Net ultrafiltration rate is independently associated with mortality in patients treated with continuous renal replacement therapy (CRRT) in a nonlinear fashion. The relationship between mortality and net ultrafiltration (UF_{NET}) in patients in the intensive care unit with acute kidney injury (AKI) is J shaped. In a post hoc analysis of >1400 patients from the Randomized Evaluation of Normal versus Augmented Level Replacement Therapy (RENAL) study comparing high versus low dose of CRRT for AKI in the intensive care unit,⁴⁵⁸ low UF_{NET} rates <1.01 mL/kg per hour, and high UF_{NET} rates >1.75 mL/kg per hour were associated with higher risk-adjusted 90-day mortality compared with UF_{NET} rates in a middle range of 1.01–1.75 mL/kg per hour. It has been proposed that these effects are mediated by harms of organ edema in those treated with low UF_{NET} rates and by organ ischemia in those treated with high UF_{NET} rates, but no prospective trial data yet exist to demonstrate that targeting a moderate rate of UF_{NET} improves outcomes. (Data from Murugan R, Kerti SJ, Chang CH, et al. Association of net ultrafiltration rate with mortality among critically ill adults with acute kidney injury receiving continuous venovenous hemodiafiltration: a secondary analysis of the randomized evaluation of normal vs augmented level [RENAL] of renal replacement therapy trial. *JAMA Netw Open*. 2019;2[6]:e195418. Reprinted with permission from Murugan R, Bellomo R, Palevsky PM, Kellum JA. Ultrafiltration in critically ill patients treated with kidney replacement therapy. *Nat Rev Nephrol*. 2021;17[4]:262–276.)

DRUG-DOSING CONSIDERATIONS

Specific drug-dosing considerations in critically ill patients with AKI include attention to drugs with long half-lives and those that rapidly achieve therapeutic concentrations. Importantly, loading doses do not typically need to be adjusted in patients with AKI or those requiring KRT. In contrast, maintenance doses require adjustments in patients with AKI or those requiring KRT. When KRT is initiated, dosing strategies must further account for the extracorporeal clearance of medications. Drug characteristics (e.g., volume of distribution, molecular weight, and extent of protein binding), KRT prescriptions (e.g., blood flow, session duration, proportion of convective and diffusive clearances), and dialyzer features (e.g., porosity or flux, surface area or efficiency) are important determinants of medication clearance. In some cases, KRT may have a more subtle impact on drug clearance by enhancing extrarenal drug metabolism, perhaps through the removal of uremic toxins.⁴⁹⁶ For recipients of IHD, medications should be administered after dialysis or a postdialysis supplemental dose should be considered. With the predominant use of high-flux filters, which enhance drug clearance, dosing

recommendations that were derived for use with low-flux filters typically need to be increased by 25% to 50%.⁴⁹⁶

The advent of PIRRT poses a further challenge because extracorporeal clearance is augmented by this modality as compared with IHD. This may result in increased drug removal and inadequate blood concentrations of vital medications including antibiotics. To date, drug-dosing guidance with PIRRT is available for only a few agents.⁴⁹⁷ As a general rule, drug doses should probably be higher than those administered to patients receiving IHD and, where feasible, their use should be accompanied by careful drug level monitoring.⁴⁹⁸ A “top-up” dose following the conclusion of a PIRRT session is advisable for agents that are likely to be dialyzable.

In patients receiving CRRT, total effluent dose and the relative breakdown of convective and diffusive therapy will affect drug removal and subsequent dosing changes. CRRT at usual doses, assuming no interruptions to therapy, begins to approximate endogenous kidney function (~30 to 60 mL/min), and more frequent drug dosing is generally required than with IHD or PIRRT. For a more detailed discussion of this topic, the reader is referred to [Chapter 56](#).

EXTRACORPOREAL BLOOD PURIFICATION

Extracorporeal blood purification (EBP) entails removal of pathogens, endotoxins, and inflammatory mediators from the blood and has evolving applications in critically ill patients with multiorgan failure and sepsis.⁴⁹⁹ Among the various EBP technologies, adsorptive and immunomodulatory EBP devices have gained interest.

In the setting of infection, immune cells recognize pathogen-associated membrane proteins (PAMPs) that activate the production of antiinflammatory and proinflammatory cytokines. Injured host cells also express damage-associated molecular patterns, which cause leukocyte activation and cytokine production.^{499,500} EBP technologies using adsorption provide means to modulate a dysregulated immune system by removal of inflammatory mediators and highly protein-bound toxins that are not effectively cleared by diffusion or convection. Adsorption consists of removal of molecules via attachment to various sorbents given chemical affinity to ion exchange resins and chemisorbents.^{501,502}

Traditionally, adsorptive-based hemoperfusion has been used for the treatment of intoxications (e.g., paraquat poisoning) and extracorporeal liver support devices.⁵⁰³ Liver support devices, such as the molecular adsorbent recirculating system, remove albumin-bound solutes and have been used for refractory cholestatic pruritus or drug intoxications, especially when accompanied by severe liver injury and for acute or acute-on-chronic liver failure as a bridge to liver transplantation.⁵⁰³⁻⁵⁰⁶

The use of novel adsorptive-based EBP devices was accelerated during the COVID-19 pandemic. These new technologies include filters for direct pathogen removal (Seraph 100 Microbind, GARNET, Hemopurifier),^{507,508} endotoxin removal (Toraymyxin),⁴⁹⁹ cytokine removal (CytoSorb, oXiris, SepXiris),^{499,509} and immunomodulatory devices such as the selective cytopheretic device (SCD).⁵¹⁰ SCD is a synthetic membrane cartridge that binds and

deactivates activated leukocytes and therefore mitigates cytokine production.

While scientific progress in EBP is promising, outstanding questions related to clinical utility, patient selection, timing of initiation, duration of treatment, heterogeneity of treatment effect, and others remain unanswered.^{13,511} Importantly, the application of adsorptive-based nonselective clearance to remove both proinflammatory and antiinflammatory mediators may not be beneficial if applied at the wrong time, given that some patients may develop immune dysfunction or immunoparalysis characterized by impaired cytokine production and subsequent increased risk of infection.^{10,11,512,513} Nonetheless, extensive research is underway to address key questions about how various extracorporeal therapies and the interactions among the host, pathogen/insult, and medical devices affect patient-centered outcomes in critical illness and multiorgan failure.⁵¹⁴

 Visit Elsevier eBooks+ (eBooks.Health.Elsevier.com) for a complete set of references and Board Review Questions.

KEY REFERENCES

2. Hoste EA, Bagshaw SM, Bellomo R, et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med.* 2015;41(8):1411–1423.
7. White KC, Serpa-Neto A, Hurford R, et al. Sepsis-associated acute kidney injury in the intensive care unit: incidence, patient characteristics, timing, trajectory, treatment, and associated outcomes. A multicenter, observational study. *Intensive Care Med.* 2023;49(9):1079–1089.
13. Zarbock A, Nadim MK, Pickkers P, et al. Sepsis-associated acute kidney injury: consensus report of the 28th Acute Disease Quality Initiative workgroup. *Nat Rev Nephrol.* 2023;19(6):401–417.
15. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl.* 2012;2(1):1–138.
26. Gomez H, Ince C, De Backer D, et al. A unified theory of sepsis-induced acute kidney injury: inflammation, microcirculatory dysfunction, bioenergetics, and the tubular cell adaptation to injury. *Shock.* 2014;41(1):3–11.
36. Wang Y, Bellomo R. Cardiac surgery-associated acute kidney injury: risk factors, pathophysiology and treatment. *Nat Rev Nephrol.* 2017;13(11):697–711.
38. Zarbock A, Weiss R, Albert F, et al. Epidemiology of surgery associated acute kidney injury (EPIS-AKI): a prospective international observational multicenter clinical study. *Intensive Care Med.* 2023;49(12):1441–1455.
49. Zarbock A, Kullmar M, Ostermann M, et al. Prevention of cardiac surgery-associated acute kidney injury by implementing the KDIGO guidelines in high-risk patients identified by biomarkers: the PrevAKI-multicenter randomized controlled trial. *Anesth Analg.* 2021;133(2):292–302.
66. Joannidis M, Forni LG, Klein SJ, et al. Lung-kidney interactions in critically ill patients: consensus report of the Acute Disease Quality Initiative (ADQI) 21 Workgroup. *Intensive Care Med.* 2020;46(4):654–672.
67. Villa G, Katz N, Ronco C. Extracorporeal membrane oxygenation and the kidney. *Cardiorenal Med.* 2015;6(1):50–60.
75. Roehm B, Vest AR, Weiner DE. Left ventricular assist devices, kidney disease, and dialysis. *Am J Kidney Dis.* 2018;71(2):257–266.
82. Teixeira JP, Ambruso S, Griffin BR, Faubel S. Pulmonary consequences of acute kidney injury. *Semin Nephrol.* 2019;39(1):3–16.
84. Bellani G, Laffey JG, Pham T, et al. Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA.* 2016;315(8):788–800.
87. Brower RG, Matthay MA, Acute Respiratory Distress Syndrome Network, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med.* 2000;342(18):1301–1308.
94. Wiedemann HP, Wheeler AP, National Heart Lung and Blood Institute Acute Respiratory Distress Syndrome Clinical Trials Network, et al. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med.* 2006;354(24):2564–2575.
100. Velez JCQ, Therapondos G, Juncos LA. Reappraising the spectrum of AKI and hepatorenal syndrome in patients with cirrhosis. *Nat Rev Nephrol.* 2020;16(3):137–155.
113. Wong F, Pappas SC, Curry MP, et al. Terlipressin plus albumin for the treatment of type 1 hepatorenal syndrome. *N Engl J Med.* 2021;384(9):818–828.
114. Pichler RH, Swenson ER, Leary PJ, Paine CH. Terlipressin: hopes fulfilled or dashed? *Clin J Am Soc Nephrol.* 2022;17(1):140–142.
117. China L, Freemantle N, Forrest E, et al. A randomized trial of albumin infusions in hospitalized patients with cirrhosis. *N Engl J Med.* 2021;384(9):808–817.
131. Gupta S, Fennes AZ, Hootkins R. The role of RRT in hyperammonemic patients. *Clin J Am Soc Nephrol.* 2016;11(10):1872–1878.
134. Kirkpatrick AW, Roberts DJ, De Waele J, et al. Intra-abdominal hypertension and the abdominal compartment syndrome: updated consensus definitions and clinical practice guidelines from the World Society of the Abdominal Compartment Syndrome. *Intensive Care Med.* 2013;39(7):1190–1206.
153. Sinha Ray A, Haikal A, Hammoud KA, Yu ASL. Vancomycin and the risk of AKI: a systematic review and meta-analysis. *Clin J Am Soc Nephrol.* 2016;11(12):2132–2140.
178. Qian ET, Casey JD, Wright A, et al. Cefepime vs piperacillin-tazobactam in adults hospitalized with acute infection: the ACORN randomized clinical trial. *JAMA.* 2023;330(16):1557–1567.
200. Hiremath S, Velez JCQ. Preventing a nonexistent entity: the curious case of contrast and acute kidney injury. *Curr Opin Nephrol Hypertens.* 2020;29(1):152–160.
258. Perner A, Prowle J, Joannidis M, Young P, Hjortrup PB, Pettila V. Fluid management in acute kidney injury. *Intensive Care Med.* 2017;43(6):807–815.
264. Meyhoff TS, Hjortrup PB, Wetterslev J, et al. Restriction of intravenous fluid in ICU patients with septic shock. *N Engl J Med.* 2022;386(26):2459–2470.
265. Shapiro NI, Douglas IS, National Heart Lung and Blood Institute Prevention and Early Treatment of Acute Lung Injury Clinical Trials Network, et al. Early restrictive or liberal fluid management for sepsis-induced hypotension. *N Engl J Med.* 2023;388(6):499–510.
266. Murphy CV, Schramm GE, Doherty JA, et al. The importance of fluid management in acute lung injury secondary to septic shock. *Chest.* 2009;136(1):102–109.
278. Hoste EA, Maitland K, Brudney CS, et al. Four phases of intravenous fluid therapy: a conceptual model. *Br J Anaesth.* 2014;113(5):740–747.
307. Self WH, Semler MW, Wanderer JP, et al. Balanced crystalloids versus saline in noncritically ill adults. *N Engl J Med.* 2018;378(9):819–828.
308. Semler MW, Self WH, Wanderer JP, et al. Balanced crystalloids versus saline in critically ill adults. *N Engl J Med.* 2018;378(9):829–839.
309. Zampieri FG, Machado FR, Biondi RS, et al. Effect of intravenous fluid treatment with a balanced solution vs 0.9% saline solution on mortality in critically ill patients: the BaSICS randomized clinical trial. *JAMA.* 2021;326(9):1–12.
311. Finfer S, Micallef S, Hammond N, et al. Balanced multielectrolyte solution versus saline in critically ill adults. *N Engl J Med.* 2022;386(9):815–826.
315. Finfer S, Bellomo R, Boyce N, et al. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med.* 2004;350(22):2247–2256.
319. Caironi P, Tognoni G, Masson S, et al. Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med.* 2014;370(15):1412–1421.
342. Jaber S, Paugam C, Futier E, et al. Sodium bicarbonate therapy for patients with severe metabolic acidemia in the intensive care unit (BICAR-ICU): a multicentre, open-label, randomised controlled, phase 3 trial. *Lancet.* 2018;392(10141):31–40.
366. Zarbock A, Kellum JA, Schmidt C, et al. Effect of early vs delayed initiation of renal replacement therapy on mortality in critically ill patients with acute kidney injury: the ELAIN randomized clinical trial. *JAMA.* 2016;315(20):2190–2199.
367. Gaudry S, Hajage D, Schortgen F, et al. Initiation strategies for renal-replacement therapy in the intensive care unit. *N Engl J Med.* 2016;375(2):122–133.

368. Barbar SD, Clere-Jehl R, Bourredjem A, et al. Timing of renal-replacement therapy in patients with acute kidney injury and sepsis. *N Engl J Med*. 2018;379(15):1431–1442.
369. STARRT-AKI Investigators, Canadian Critical Care Trials Group, Australian New Zealand Intensive Care Society Clinical Trials Group, et al. Timing of initiation of renal-replacement therapy in acute kidney injury. *N Engl J Med*. 2020;383(3):240–251.
370. Gaudry S, Hajage D, Martin-Lefevre L, et al. Comparison of two delayed strategies for renal replacement therapy initiation for severe acute kidney injury (AKIKI 2): a multicentre, open-label, randomised, controlled trial. *Lancet*. 2021;397(10281):1293–1300.
395. Teixeira JP, Neyra JA, Tolwani A. Continuous KRT: a contemporary review. *Clin J Am Soc Nephrol*. 2023;18(2):256–269.
453. Parienti JJ, Thirion M, Megarbane B, et al. Femoral vs jugular venous catheterization and risk of nosocomial events in adults requiring acute renal replacement therapy: a randomized controlled trial. *JAMA*. 2008;299(20):2413–2422.
457. Palevsky PM, Zhang JH, VA/NIH Acute Renal Failure Trial Network, et al. Intensity of renal support in critically ill patients with acute kidney injury. *N Engl J Med*. 2008;359(1):7–20.
458. Bellomo R, Cass A, RENAL Replacement Therapy Study Investigators, et al. Intensity of continuous renal-replacement therapy in critically ill patients. *N Engl J Med*. 2009;361(17):1627–1638.
486. Neyra JA, Yessayan L, Thompson Bastin ML, Wille KM, Tolwani AJ. How to prescribe and troubleshoot continuous renal replacement therapy: a case-based review. *Kidney*. 2021;2(2):371–384.
482. Zarbock A, Kullmar M, Kindgen-Milles D, et al. Effect of regional citrate anticoagulation vs systemic heparin anticoagulation during continuous kidney replacement therapy on dialysis filter life span and mortality among critically ill patients with acute kidney injury: a randomized clinical trial. *JAMA*. 2020;324(16):1629–1639.
494. Murugan R, Bellomo R, Palevsky PM, Kellum JA. Ultrafiltration in critically ill patients treated with kidney replacement therapy. *Nat Rev Nephrol*. 2021;17(4):262–276.
499. Monard C, Rimmele T, Ronco C. Extracorporeal blood purification therapies for sepsis. *Blood Purif*. 2019;47(suppl 3):1–14.
518. Mayer KP, Joseph-Isang E, Robinson LE, Parry SM, Morris PE, Neyra JA. Safety and feasibility of physical rehabilitation and active mobilization in patients requiring continuous renal replacement therapy: a systematic review. *Crit Care Med*. 2020;48(11):e1112–e1120.

REFERENCES

1. Odutayo A, Adhikari NK, Barton J, et al. Epidemiology of acute kidney injury in Canadian critical care units: a prospective cohort study. *Can J Anaesth*. 2012;59(10):934–942.
2. Hoste EA, Bagshaw SM, Bellomo R, et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med*. 2015;41(8):1411–1423.
3. Nisula S, Kaukonen KM, Vaara ST, et al. Incidence, risk factors and 90-day mortality of patients with acute kidney injury in Finnish intensive care units: the FINNAKI study. *Intensive Care Med*. 2013;39(3):420–428.
4. Uchino S, Kellum JA, Bellomo R, et al. Acute renal failure in critically ill patients: a multinational, multicenter study. *JAMA*. 2005;294(7):813–818.
5. Johansen KL, Smith MW, Unruh ML, et al. Predictors of health utility among 60-day survivors of acute kidney injury in the veterans Affairs/National Institutes of Health Acute Renal Failure Trial Network Study. *Clin J Am Soc Nephrol*. 2010;5(8):1366–1372.
6. Bouchard J, Acharya A, Cerda J, et al. A prospective international multicenter study of AKI in the intensive care unit. *Clin J Am Soc Nephrol*. 2015;10(8):1324–1331.
7. White KC, Serpa-Neto A, Hurford R, et al. Sepsis-associated acute kidney injury in the intensive care unit: incidence, patient characteristics, timing, trajectory, treatment, and associated outcomes. A multicenter, observational study. *Intensive Care Med*. 2023;49(9):1079–1089.
8. Yegenaga I, Hoste E, Van Biesen W, et al. Clinical characteristics of patients developing ARF due to sepsis/systemic inflammatory response syndrome: results of a prospective study. *Am J Kidney Dis*. 2004;43(5):817–824.
9. Bagshaw SM, Uchino S, Bellomo R, et al. Septic acute kidney injury in critically ill patients: clinical characteristics and outcomes. *Clin J Am Soc Nephrol*. 2007;2(3):431–439.
10. Mehta RL, Bouchard J, Soroko SB, et al. Sepsis as a cause and consequence of acute kidney injury: program to improve care in acute renal disease. *Intensive Care Med*. 2011;37(2):241–248.
11. Griffin BR, Teixeira JP, Ambruso S, et al. Stage 1 acute kidney injury is independently associated with infection following cardiac surgery. *J Thorac Cardiovasc Surg*. 2021;161(4):1346–1355 e3.
12. Formeck CL, Feldman R, Althouse AD, Kellum JA. Risk and timing of de novo sepsis in critically ill children after acute kidney injury. *Kidney*. 2023;4(3):308–315.
13. Zarbock A, Nadim MK, Pickkers P, et al. Sepsis-associated acute kidney injury: consensus report of the 28th Acute Disease Quality Initiative workgroup. *Nat Rev Nephrol*. 2023;19(6):401–417.
14. Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*. 2016;315(8):801–810.
15. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl*. 2012;2(1):1–138.
16. Schrier RW, Wang W. Acute renal failure and sepsis. *N Engl J Med*. 2004;351(2):159–169.
17. Brenner M, Schaer GL, Mallory DL, Suffredini AF, Parrillo JE. Detection of renal blood flow abnormalities in septic and critically ill patients using a newly designed indwelling thermodilution renal vein catheter. *Chest*. 1990;98(1):170–179.
18. Di Giamtomasso D, May CN, Bellomo R. Vital organ blood flow during hyperdynamic sepsis. *Chest*. 2003;124(3):1053–1059.
19. Langenberg C, Wan L, Egi M, May CN, Bellomo R. Renal blood flow in experimental septic acute renal failure. *Kidney Int*. 2006;69(11):1996–2002.
20. Prowle JR, Molan MP, Hornsey E, Bellomo R. Measurement of renal blood flow by phase-contrast magnetic resonance imaging during septic acute kidney injury: a pilot investigation. *Crit Care Med*. 2012;40(6):1768–1776.
21. Calzavacca P, Evans RG, Bailey M, Bellomo R, May CN. Cortical and medullary tissue perfusion and oxygenation in experimental septic acute kidney injury. *Crit Care Med*. 2015;43(10):e431–e439.
22. Lerolle N, Nochy D, Guerot E, et al. Histopathology of septic shock induced acute kidney injury: apoptosis and leukocytic infiltration. *Intensive Care Med*. 2010;36(3):471–478.
23. Takasu O, Gaut JP, Watanabe E, et al. Mechanisms of cardiac and renal dysfunction in patients dying of sepsis. *Am J Respir Crit Care Med*. 2013;187(5):509–517.
24. Langenberg C, Gobe G, Hood S, May CN, Bellomo R. Renal histopathology during experimental septic acute kidney injury and recovery. *Crit Care Med*. 2014;42(1):e58–e67.
25. Maiden MJ, Otto S, Brealey JK, et al. Structure and function of the kidney in septic shock. A prospective controlled experimental study. *Am J Respir Crit Care Med*. 2016;194(6):692–700.
26. Gomez H, Ince C, De Backer D, et al. A unified theory of sepsis-induced acute kidney injury: inflammation, microcirculatory dysfunction, bioenergetics, and the tubular cell adaptation to injury. *Shock*. 2014;41(1):3–11.
27. Peerapornratana S, Manrique-Caballero CL, Gomez H, Kellum JA. Acute kidney injury from sepsis: current concepts, epidemiology, pathophysiology, prevention and treatment. *Kidney Int*. 2019;96(5):1083–1099.
28. Sun J, Zhang J, Tian J, et al. Mitochondria in sepsis-induced AKI. *J Am Soc Nephrol*. 2019;30(7):1151–1161.
29. Pickkers P, Angus DC, Arend J, et al. Study protocol of a randomised, double-blind, placebo-controlled, two-arm parallel-group, multi-centre phase 3 pivotal trial to investigate the efficacy and safety of recombinant human alkaline phosphatase for treatment of patients with sepsis-associated acute kidney injury. *BMJ Open*. 2023;13(4):e065613.
30. Joannes-Boyau O, Honore PM, Perez P, et al. High-volume versus standard-volume haemofiltration for septic shock patients with acute kidney injury (IVOIRE study): a multicentre randomized controlled trial. *Intensive Care Med*. 2013;39(9):1535–1546.
31. Pickkers P, Mehta RL, Murray PT, et al. Effect of human recombinant alkaline phosphatase on 7-day creatinine clearance in patients with sepsis-associated acute kidney injury: a randomized clinical trial. *JAMA*. 2018;320(19):1998–2009.
32. Dellinger RP, Bagshaw SM, Antonelli M, et al. Effect of targeted polymyxin B hemoperfusion on 28-day mortality in patients with septic shock and elevated endotoxin level: the EUPHRATES randomized clinical trial. *JAMA*. 2018;320(14):1455–1463.
33. Chertow GM, Levy EM, Hammermeister KE, Grover F, Daley J. Independent association between acute renal failure and mortality following cardiac surgery. *Am J Med*. 1998;104(4):343–348.
34. Hobson CE, Yavas S, Segal MS, et al. Acute kidney injury is associated with increased long-term mortality after cardiothoracic surgery. *Circulation*. 2009;119(18):2444–2453.
35. Hu J, Chen R, Liu S, Yu X, Zou J, Ding X. Global incidence and outcomes of adult patients with acute kidney injury after cardiac surgery: a systematic review and meta-analysis. *J Cardiothorac Vasc Anesth*. 2016;30(1):82–89.
36. Wang Y, Bellomo R. Cardiac surgery-associated acute kidney injury: risk factors, pathophysiology and treatment. *Nat Rev Nephrol*. 2017;13(11):697–711.
37. Cheruku SR, Raphael J, Neyra JA, Fox AA. Acute kidney injury after cardiac surgery: prediction, prevention, and management. *Anesthesiology*. 2023.
38. Zarbock A, Weiss R, Albert F, et al. Epidemiology of surgery associated acute kidney injury (EPIS-AKI): a prospective international observational multi-center clinical study. *Intensive Care Med*. 2023;49(12):1441–1455.
39. Garg AX, Devereaux PJ, Yusuf S, et al. Kidney function after off-pump or on-pump coronary artery bypass graft surgery: a randomized clinical trial. *JAMA*. 2014;311(21):2191–2198.
40. Lamy A, Devereaux PJ, Prabhakaran D, et al. Five-year outcomes after off-pump or on-pump coronary-artery bypass grafting. *N Engl J Med*. 2016;375(24):2359–2368.
41. Shroyer AL, Hattler B, Wagner TH, et al. Five-year outcomes after on-pump and off-pump coronary-artery bypass. *N Engl J Med*. 2017;377(7):623–632.
42. Bove T, Zangrillo A, Guarracino F, et al. Effect of fenoldopam on use of renal replacement therapy among patients with acute kidney injury after cardiac surgery: a randomized clinical trial. *JAMA*. 2014;312(21):2244–2253.
43. Hausenloy DJ, Candilio L, Evans R, et al. Remote ischemic preconditioning and outcomes of cardiac surgery. *N Engl J Med*. 2015;373(15):1408–1417.
44. Meybohm P, Bein B, Brosteanu O, et al. A multicenter trial of remote ischemic preconditioning for heart surgery. *N Engl J Med*. 2015;373(15):1397–1407.
45. Billings FT, Hendricks PA, Schildcrout JS, et al. High-dose perioperative atorvastatin and acute kidney injury following cardiac surgery: a randomized clinical trial. *JAMA*. 2016;315(9):877–888.

46. Lei C, Berra L, Rezoagli E, et al. Nitric oxide decreases acute kidney injury and stage 3 chronic kidney disease after cardiac surgery. *Am J Respir Crit Care Med*. 2018;198(10):1279–1287.
47. Kamenshchikov NO, Anfimenova YJ, Kozlov BN, et al. Nitric oxide delivery during cardiopulmonary bypass reduces acute kidney injury: a randomized trial. *J Thorac Cardiovasc Surg*. 2022;163(4):1393–1403.
48. Marrazzo F, Spina S, Zadek F, et al. Protocol of a randomised controlled trial in cardiac surgical patients with endothelial dysfunction aimed to prevent postoperative acute kidney injury by administering nitric oxide gas. *BMJ Open*. 2019;9(7):e026848.
49. Zarbock A, Kullmar M, Ostermann M, et al. Prevention of cardiac surgery-associated acute kidney injury by implementing the KDIGO guidelines in high-risk patients identified by biomarkers: the PrevAKI-multicenter randomized controlled trial. *Anesth Analg*. 2021;133(2):292–302.
50. Sanaiha Y, Bailey K, Downey P, et al. Trends in mortality and resource utilization for extracorporeal membrane oxygenation in the United States: 2008–2014. *Surgery*. 2019;165(2):381–388.
51. Grasselli G, Calfee CS, Camporota L, et al. ESICM guidelines on acute respiratory distress syndrome: definition, phenotyping and respiratory support strategies. *Intensive Care Med*. 2023;49(7):727–759.
52. Brogan TV, Thiagarajan RR, Rycus PT, Bartlett RH, Bratton SL. Extracorporeal membrane oxygenation in adults with severe respiratory failure: a multi-center database. *Intensive Care Med*. 2009;35(12):2105–2114.
53. Yan X, Jia S, Meng X, et al. Acute kidney injury in adult postcardiotomy patients with extracorporeal membrane oxygenation: evaluation of the RIFLE classification and the Acute Kidney Injury Network criteria. *Eur J Cardio Thorac Surg*. 2010;37(2):334–338.
54. Chen YC, Tsai FC, Chang CH, et al. Prognosis of patients on extracorporeal membrane oxygenation: the impact of acute kidney injury on mortality. *Ann Thorac Surg*. 2011;91(1):137–142.
55. Han SS, Kim HJ, Lee SJ, et al. Effects of renal replacement therapy in patients receiving extracorporeal membrane oxygenation: a meta-analysis. *Ann Thorac Surg*. 2015;100(4):1485–1495.
56. Kielstein JT, Heiden AM, Beutel G, et al. Renal function and survival in 200 patients undergoing ECMO therapy. *Nephrol Dial Transplant*. 2013;28(1):86–90.
57. Aubron C, Cheng AC, Pilcher D, et al. Factors associated with outcomes of patients on extracorporeal membrane oxygenation support: a 5-year cohort study. *Crit Care*. 2013;17(2):R73.
58. Seczynska B, Krolikowski W, Nowak I, Jankowski M, Szuldrzynski K, Szczeklik W. Continuous renal replacement therapy during extracorporeal membrane oxygenation in patients treated in medical intensive care unit: technical considerations. *Ther Apher Dial*. 2014;18(6):523–534.
59. Hoover NG, Heard M, Reid C, et al. Enhanced fluid management with continuous venovenous hemofiltration in pediatric respiratory failure patients receiving extracorporeal membrane oxygenation support. *Intensive Care Med*. 2008;34(12):2241–2247.
60. Selewski DT, Cornell TT, Blatt NB, et al. Fluid overload and fluid removal in pediatric patients on extracorporeal membrane oxygenation requiring continuous renal replacement therapy. *Crit Care Med*. 2012;40(9):2694–2699.
61. Gbadegesin R, Zhao S, Charpie J, Brophy PD, Smoyer WE, Lin JJ. Significance of hemolysis on extracorporeal life support after cardiac surgery in children. *Pediatr Nephrol*. 2009;24(3):589–595.
62. Schmidt M, Bailey M, Kelly J, et al. Impact of fluid balance on outcome of adult patients treated with extracorporeal membrane oxygenation. *Intensive Care Med*. 2014;40(9):1256–1266.
63. Dado DN, Ainsworth CR, Thomas SB, et al. Outcomes among patients treated with renal replacement therapy during extracorporeal membrane oxygenation: a single-center retrospective study. *Blood Purif*. 2020;49(3):341–347.
64. Gunning S, Kutub F, Rose R, Trevino S, Song T, Koyner JL. Fluid overload and mortality in patients with severe acute kidney injury and extracorporeal membrane oxygenation. *Kidney*. 2020;1(4):232–240.
65. *Extracorporeal Life Support Organization Guidelines for Cardiopulmonary Extracorporeal Life Support, Version 1.4*. 2017. Ann Arbor, MI, USA. <https://www.elseo.org/Resources/Guidelines.aspx>.
66. Joannidis M, Forni LG, Klein SJ, et al. Lung-kidney interactions in critically ill patients: consensus report of the Acute Disease Quality Initiative (ADQI) 21 Workgroup. *Intensive Care Med*. 2020;46(4):654–672.
67. Villa G, Katz N, Ronco C. Extracorporeal membrane oxygenation and the kidney. *Cardiorenal Med*. 2015;6(1):50–60.
68. Lyu L, Long C, Hei F, et al. Plasma free hemoglobin is a predictor of acute renal failure during adult venous-arterial extracorporeal membrane oxygenation support. *J Cardiothorac Vasc Anesth*. 2016;30(4):891–895.
69. Borasino S, Kalra Y, Elam AR, et al. Impact of hemolysis on acute kidney injury and mortality in children supported with cardiac extracorporeal membrane oxygenation. *J Extra Corpor Technol*. 2018;50(4):217–224. <https://www.ncbi.nlm.nih.gov/pubmed/30581228>.
70. Lou S, MacLaren G, Best D, Delzoppo C, Butt W. Hemolysis in pediatric patients receiving centrifugal-pump extracorporeal membrane oxygenation: prevalence, risk factors, and outcomes. *Crit Care Med*. 2014;42(5):1213–1220.
71. Kirklin JK, Naftel DC, Pagani FD, et al. Seventh INTERMACS annual report: 15,000 patients and counting. *J Heart Lung Transplant*. 2015;34(12):1495–1504.
72. Varshney AS, DeFilippis EM, Cowger JA, Netuka I, Pinney SP, Givertz MM. Trends and outcomes of left ventricular assist device therapy: JACC focus seminar. *J Am Coll Cardiol*. 2022;79(11):1092–1107.
73. Brisco MA, Kimmel SE, Coca SG, et al. Prevalence and prognostic importance of changes in renal function after mechanical circulatory support. *Circ Heart Fail*. 2014;7(1):68–75.
74. Raichlin E, Baibhav B, Lowes BD, et al. Outcomes in patients with severe preexisting renal dysfunction after continuous-flow left ventricular assist device implantation. *ASAIO J*. 2016;62(3):261–267.
75. Roehm B, Vest AR, Weiner DE. Left ventricular assist devices, kidney disease, and dialysis. *Am J Kidney Dis*. 2018;71(2):257–266.
76. Ross DW, Stevens GR, Wanchoo R, et al. Left ventricular assist devices and the kidney. *Clin J Am Soc Nephrol*. 2018;13(2):348–355.
77. Borgi J, Tsiouris A, Hodari A, Cogan CM, Paone G, Morgan JA. Significance of postoperative acute renal failure after continuous-flow left ventricular assist device implantation. *Ann Thorac Surg*. 2013;95(1):163–169.
78. Muslem R, Caliskan K, Akin S, et al. Acute kidney injury and 1-year mortality after left ventricular assist device implantation. *J Heart Lung Transplant*. 2018;37(1):116–123.
79. Liu KD, Glidden DV, Eisner MD, et al. Predictive and pathogenetic value of plasma biomarkers for acute kidney injury in patients with acute lung injury. *Crit Care Med*. 2007;35(12):2755–2761. <https://www.ncbi.nlm.nih.gov/pubmed/18074478>.
80. Demirjian S, Chertow GM, Zhang JH, et al. Model to predict mortality in critically ill adults with acute kidney injury. *Clin J Am Soc Nephrol*. 2011;6(9):2114–2120.
81. McNicholas BA, Rezoagli E, Pham T, et al. Impact of early acute kidney injury on management and outcome in patients with acute respiratory distress syndrome: a secondary analysis of a multicenter observational study. *Crit Care Med*. 2019;47(9):1216–1225.
82. Teixeira JP, Ambruso S, Griffin BR, Faubel S. Pulmonary consequences of acute kidney injury. *Semin Nephrol*. 2019;39(1):3–16.
83. Faubel S, Edelstein CL. Mechanisms and mediators of lung injury after acute kidney injury. *Nat Rev Nephrol*. 2016;12(1):48–60.
84. Bellani G, Laffey JG, Pham T, et al. Epidemiology, patterns of care, and mortality for acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA*. 2016;315(8):788–800.
85. Ranieri VM, Rubenfeld GD, ARDS Definition Task Force, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA*. 2012;307(2):2526–2533.
86. Matthay MA, Arabi Y, Arroliga AC, et al. A new global definition of acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2023.
87. Brower RG, Matthay MA, Acute Respiratory Distress Syndrome Network, et al. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342(18):1301–1308.
88. Imai Y, Parodo J, Kajikawa O, et al. Injurious mechanical ventilation and end-organ epithelial cell apoptosis and organ dysfunction in an experimental model of acute respiratory distress syndrome. *JAMA*. 2003;289(16):2104–2112.

89. Choi WI, Quinn DA, Park KM, et al. Systemic microvascular leak in an in vivo rat model of ventilator-induced lung injury. *Am J Respir Crit Care Med*. 2003;167(12):1627–1632.
90. van den Akker JP, Egal M, Groeneveld AB. Invasive mechanical ventilation as a risk factor for acute kidney injury in the critically ill: a systematic review and meta-analysis. *Crit Care*. 2013;17(3):R98.
91. Liu KD, Matthay MA, Chertow GM. Evolving practices in critical care and potential implications for management of acute kidney injury. *Clin J Am Soc Nephrol*. 2006;1(4):869–873.
92. Kallet RH, Jasmer RM, Luce JM, Lin LH, Marks JD. The treatment of acidosis in acute lung injury with tris-hydroxymethyl aminomethane (THAM). *Am J Respir Crit Care Med*. 2000;161(4 Pt 1):1149–1153.
93. McNamee JJ, Gillies MA, Barrett NA, et al. Effect of lower tidal volume ventilation facilitated by extracorporeal carbon dioxide removal vs standard care ventilation on 90-day mortality in patients with acute hypoxemic respiratory failure: the REST randomized clinical trial. *JAMA*. 2021;326(11):1013–1023.
94. Wiedemann HP, Wheeler AP, National Heart Lung and Blood Institute Acute Respiratory Distress Syndrome Clinical Trials Network, et al. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med*. 2006;354(24):2564–2575.
95. Liu KD, Thompson BT, Ancukiewicz M, et al. Acute kidney injury in patients with acute lung injury: impact of fluid accumulation on classification of acute kidney injury and associated outcomes. *Crit Care Med*. 2011;39(12):2665–2671.
96. Han F, Sun R, Ni Y, et al. Early initiation of continuous renal replacement therapy improves clinical outcomes in patients with acute respiratory distress syndrome. *Am J Med Sci*. 2015;349(3):199–205.
97. Putzu A, Fang MX, Boscolo BM, et al. Blood purification with continuous veno-venous hemofiltration in patients with sepsis or ARDS: a systematic review and meta-analysis. *Minerva Anestesiol*. 2017;83(8):867–877.
98. Fielding-Singh V, Matthay MA, Calfee CS. Beyond low tidal volume ventilation: treatment adjuncts for severe respiratory failure in acute respiratory distress syndrome. *Crit Care Med*. 2018;46(11):1820–1831.
99. Biggins SW, Angeli P, Garcia-Tsao G, et al. Diagnosis, evaluation, and management of ascites, spontaneous bacterial peritonitis and hepatorenal syndrome: 2021 practice guidance by the American association for the study of liver diseases. *Hepatology*. 2021;74(2):1014–1048.
100. Velez JCQ, Therapondos G, Juncos LA. Reappraising the spectrum of AKI and hepatorenal syndrome in patients with cirrhosis. *Nat Rev Nephrol*. 2020;16(3):137–155.
101. Cullaro G, Kanduri SR, Velez JCQ. Acute kidney injury in patients with liver disease. *Clin J Am Soc Nephrol*. 2022;17(11):1674–1684.
102. Velez JCQ. Hepatorenal syndrome type 1: from diagnosis ascertainment to goal-oriented pharmacologic therapy. *Kidney*. 2022;3(2):382–395.
103. Patidar KR, Belcher JM, Regner KR, et al. Incidence and outcomes of acute kidney injury including hepatorenal syndrome in hospitalized patients with cirrhosis in the US. *J Hepatol*. 2023;79(6):1408–1417.
104. Allegretti AS, Parada XV, Eneanya ND, et al. Prognosis of patients with cirrhosis and AKI who initiate RRT. *Clin J Am Soc Nephrol*. 2018;13(1):16–25.
105. Angeli P, Gines P, Wong F, et al. Diagnosis and management of acute kidney injury in patients with cirrhosis: revised consensus recommendations of the International Club of Ascites. *J Hepatol*. 2015;62(4):968–974.
106. Aron AW, Amatruda JG. Fractional excretion of sodium and urea are useful tools in the evaluation of AKI: CON. *Kidney*. 2023;4(6):e728–e730.
107. Patidar KR, Kang L, Bajaj JS, Carl D, Sanyal AJ. Fractional excretion of urea: a simple tool for the differential diagnosis of acute kidney injury in cirrhosis. *Hepatology*. 2018;68(1):224–233.
108. Patidar KR, Piano S, Cullaro G, Belcher JM, Allegretti AS, Consortium HR-H. Recent advances in the management of hepatorenal syndrome: a us perspective. *Clin Gastroenterol Hepatol*. 2023;21(4):897–901.
109. Belcher JM, Sanyal AJ, Peixoto AJ, et al. Kidney biomarkers and differential diagnosis of patients with cirrhosis and acute kidney injury. *Hepatology*. 2014;60(2):622–632.
110. Ariza X, Sola E, Elia C, et al. Analysis of a urinary biomarker panel for clinical outcomes assessment in cirrhosis. *PLoS One*. 2015;10(6):e0128145.
111. Gambino C, Piano S, Stenico M, et al. Diagnostic and prognostic performance of urinary neutrophil gelatinase-associated lipocalin in patients with cirrhosis and acute kidney injury. *Hepatology*. 2023;77(5):1630–1638.
112. Huelin P, Sola E, Elia C, et al. Neutrophil gelatinase-associated lipocalin for assessment of acute kidney injury in cirrhosis: a prospective study. *Hepatology*. 2019;70(1):319–333.
113. Wong F, Pappas SC, Curry MP, et al. Terlipressin plus albumin for the treatment of type 1 hepatorenal syndrome. *N Engl J Med*. 2021;384(9):818–828.
114. Pichler RH, Swenson ER, Leary PJ, Paine CH. Terlipressin: hopes fulfilled or dashed? *Clin J Am Soc Nephrol*. 2022;17(1):140–142.
115. TERLIVAZ [package insert], Bedminster NJ: Mallinckrodt Hospital Products. Inc.; 2022.
116. Moreau R, Jalan R, Gines P, et al. Acute-on-chronic liver failure is a distinct syndrome that develops in patients with acute decompensation of cirrhosis. *Gastroenterology*. 2013;144(7):1426–1437, 1437 e1–9.
117. China L, Freemantle N, Forrest E, et al. A randomized trial of albumin infusions in hospitalized patients with cirrhosis. *N Engl J Med*. 2021;384(9):808–817.
118. Cade R, Wagemaker H, Vogel S, et al. Hepatorenal syndrome. Studies of the effect of vascular volume and intraperitoneal pressure on renal and hepatic function. *Am J Med*. 1987;82(3):427–438.
119. Savino JA, Cerabona T, Agarwal N, Byrne D. Manipulation of ascitic fluid pressure in cirrhotics to optimize hemodynamic and renal function. *Ann Surg*. 1988;208(4):504–511.
120. Umgelter A, Reindl W, Franzen M, Lenhardt C, Huber W, Schmid RM. Renal resistive index and renal function before and after paracentesis in patients with hepatorenal syndrome and tense ascites. *Intensive Care Med*. 2009;35(1):152–156.
121. Seethapathy H, Sharma S, Zhao S, et al. Acute kidney injury following paracentesis among inpatients with cirrhosis. *Kidney Int Rep*. 2020;5(8):1305–1308.
122. Velez JCQ, Petkovich B, Karakala N, Huggins JT. Point-of-Care echocardiography unveils misclassification of acute kidney injury as hepatorenal syndrome. *Am J Nephrol*. 2019;50(3):204–211.
123. Lee WM, Stravitz RT, Larson AM. Introduction to the revised American association for the study of liver diseases position paper on acute liver failure 2011. *Hepatology*. 2012;55(3):965–967.
124. Cardoso FS, Gottfried M, Tuijss S, Olson JC, Karvellas CJ, Group USALFS. Continuous renal replacement therapy is associated with reduced serum ammonia levels and mortality in acute liver failure. *Hepatology*. 2018;67(2):711–720.
125. Stravitz RT, Kramer DJ. Management of acute liver failure. *Nat Rev Gastroenterol Hepatol*. 2009;6(9):542–553.
126. Poveda MJ, Bernabeu A, Concepcion L, et al. Brain edema dynamics in patients with overt hepatic encephalopathy A magnetic resonance imaging study. *Neuroimage*. 2010;52(2):481–487.
127. Haussinger D. Low grade cerebral edema and the pathogenesis of hepatic encephalopathy in cirrhosis. *Hepatology*. 2006;43(6):1187–1190.
128. Canalese J, Gimson AE, Davis C, Mellon PJ, Davis M, Williams R. Controlled trial of dexamethasone and mannitol for the cerebral oedema of fulminant hepatic failure. *Gut*. 1982;23(7):625–629. <https://www.ncbi.nlm.nih.gov/pubmed/6806155>.
129. Murphy N, Auzinger G, Bernel W, Wendon J. The effect of hypertonic sodium chloride on intracranial pressure in patients with acute liver failure. *Hepatology*. 2004;39(2):464–470.
130. Davenport A, Will EJ, Davidson AM. Improved cardiovascular stability during continuous modes of renal replacement therapy in critically ill patients with acute hepatic and renal failure. *Crit Care Med*. 1993;21(3):328–338. <https://www.ncbi.nlm.nih.gov/pubmed/8440100>.
131. Gupta S, Fennes AZ, Hootkins R. The role of RRT in hyperammonemic patients. *Clin J Am Soc Nephrol*. 2016;11(10):1872–1878.
132. Gonwa TA, Wadei HM. The challenges of providing renal replacement therapy in decompensated liver cirrhosis. *Blood Purif*. 2012;33(1–3):144–148.
133. Paine C, Pichler R. How we treat hyperammonemia in acute liver failure. *Clin J Am Soc Nephrol*. 2024;19(2):254–256.
134. Kirkpatrick AW, Roberts DJ, De Waele J, et al. Intra-abdominal hypertension and the abdominal compartment syndrome: updated consensus definitions and clinical practice guidelines from the

- World Society of the Abdominal Compartment Syndrome. *Intensive Care Med.* 2013;39(7):1190–1206.
135. Malbrain ML, Chiumello D, Pelosi P, et al. Incidence and prognosis of intraabdominal hypertension in a mixed population of critically ill patients: a multiple-center epidemiological study. *Crit Care Med.* 2005;33(2):315–322. <https://www.ncbi.nlm.nih.gov/pubmed/15699833>.
 136. Vidal MG, Ruiz Weissner J, Gonzalez F, et al. Incidence and clinical effects of intra-abdominal hypertension in critically ill patients. *Crit Care Med.* 2008;36(6):1823–1831.
 137. Dalfino L, Tullo L, Donadio I, Malcangi V, Brienza N. Intra-abdominal hypertension and acute renal failure in critically ill patients. *Intensive Care Med.* 2008;34(4):707–713.
 138. De Waele JJ, De Laet I, Kirkpatrick AW, Hoste E. Intra-abdominal hypertension and abdominal compartment syndrome. *Am J Kidney Dis.* 2011;57(1):159–169.
 139. Mohmand H, Goldfarb S. Renal dysfunction associated with intra-abdominal hypertension and the abdominal compartment syndrome. *J Am Soc Nephrol.* 2011;22(4):615–621.
 140. Mullens W, Abrahams Z, Francis GS, Taylor DO, Starling RC, Tang WH. Prompt reduction in intra-abdominal pressure following large-volume mechanical fluid removal improves renal insufficiency in refractory decompensated heart failure. *J Card Fail.* 2008;14(6):508–514.
 141. Schachtrupp A, Lawong G, Afify M, Graf J, Toens C, Schumpelick V. Fluid resuscitation preserves cardiac output but cannot prevent organ damage in a porcine model during 24 h of intraabdominal hypertension. *Shock.* 2005;24(2):153–158. <https://www.ncbi.nlm.nih.gov/pubmed/16044086>.
 142. Kula R, Szturz P, Sklienka P, Neiser J, Jahoda J. A role for negative fluid balance in septic patients with abdominal compartment syndrome? *Intensive Care Med.* 2004;30(11):2138–2139.
 143. Oda S, Hirasawa H, Shiga H, et al. Management of intra-abdominal hypertension in patients with severe acute pancreatitis with continuous hemodiafiltration using a polymethyl methacrylate membrane hemofilter. *Ther Apher Dial.* 2005;9(4):355–361.
 144. Sood P, Dass B, Bakuzonis C, Ross EA. Intra-abdominal hypertension can be monitored with femoral vein catheters during CRRT and may cause access recirculation. *Clin Nephrol.* 2010;74(3):223–228. <https://www.ncbi.nlm.nih.gov/pubmed/20860907>.
 145. Trainor D, Borthwick E, Ferguson A. Perioperative management of the hemodialysis patient. *Semin Dial.* 2011;24(3):314–326.
 146. Miller A, Price G. Gabapentin toxicity in renal failure: the importance of dose adjustment. *Pain Med.* 2009;10(1):190–192.
 147. Payne LE, Gagnon DJ, Riker RR, et al. Cefepime-induced neurotoxicity: a systematic review. *Crit Care.* 2017;21(1):276.
 148. Filippone EJ, Kraft WK, Farber JL. The nephrotoxicity of vancomycin. *Clin Pharmacol Ther.* 2017;102(3):459–469.
 149. Murphy ME, Barreto EF. Vancomycin should Be considered a nephrotoxic antimicrobial agent: PRO. *Kidney.* 2022;3(9):1484–1487.
 150. Mullaney SR. Vancomycin should Be considered a nephrotoxic antimicrobial agent: CON. *Kidney.* 2022;3(9):1488–1490.
 151. Chambers HF, Deleo FR. Waves of resistance: Staphylococcus aureus in the antibiotic era. *Nat Rev Microbiol.* 2009;7(9):629–641.
 152. van Hal SJ, Paterson DL, Lodise TP. Systematic review and meta-analysis of vancomycin-induced nephrotoxicity associated with dosing schedules that maintain troughs between 15 and 20 milligrams per liter. *Antimicrob Agents Chemother.* 2013;57(2):734–744.
 153. Sinha Ray A, Haikal A, Hammoud KA, Yu ASL. Vancomycin and the risk of AKI: a systematic review and meta-analysis. *Clin J Am Soc Nephrol.* 2016;11(12):2132–2140.
 154. Luque Y, Louis K, Jouanneau C, et al. Vancomycin-associated cast nephropathy. *J Am Soc Nephrol.* 2017;28(6):1723–1728.
 155. Rafat C, Zaworski J, Luque Y, et al. Noninvasive screening of vancomycin-associated cast nephropathy. *Kidney Int.* 2022;101(2):425.
 156. Rybak MJ, Albrecht LM, Boike SC, Chandrasekar PH. Nephrotoxicity of vancomycin, alone and with an aminoglycoside. *J Antimicrob Chemother.* 1990;25(4):679–687.
 157. Lodise TP, Lomaestro B, Graves J, Drusano GL. Larger vancomycin doses (at least four grams per day) are associated with an increased incidence of nephrotoxicity. *Antimicrob Agents Chemother.* 2008;52(4):1330–1336.
 158. Bosso JA, Nappi J, Rudisill C, et al. Relationship between vancomycin trough concentrations and nephrotoxicity: a prospective multicenter trial. *Antimicrob Agents Chemother.* 2011;55(12):5475–5479.
 159. Flannery AH, Bissell BD, Bastin MT, Morris PE, Neyra JA. Continuous versus intermittent infusion of vancomycin and the risk of acute kidney injury in critically ill adults: a systematic review and meta-analysis. *Crit Care Med.* 2020;48(6):912–918.
 160. Chen AY, Deng CY, Calvachi-Prieto P, et al. A large-scale multicenter retrospective study on nephrotoxicity associated with empiric broad-spectrum antibiotics in critically ill patients. *Chest.* 2023;164(2):355–368.
 161. Moenster RP, Linneman TW, Finnegan PM, Hand S, Thomas Z, McDonald JR. Acute renal failure associated with vancomycin and beta-lactams for the treatment of osteomyelitis in diabetics: piperacillin-tazobactam as compared with cefepime. *Clin Microbiol Infect.* 2014;20(6):O384–O389.
 162. Giuliano CA, Patel CR, Kale-Pradhan PB. Is the combination of piperacillin-tazobactam and vancomycin associated with development of acute kidney injury? A meta-analysis. *Pharmacotherapy.* 2016;36(12):1217–1228.
 163. Hammond DA, Smith MN, Painter JT, Meena NK, Lusardi K. Comparative incidence of acute kidney injury in critically ill patients receiving vancomycin with concomitant piperacillin-tazobactam or cefepime: a retrospective cohort study. *Pharmacotherapy.* 2016;36(5):463–471.
 164. Hammond DA, Smith MN, Li C, Hayes SM, Lusardi K, Bookstaver PB. Systematic review and meta-analysis of acute kidney injury associated with concomitant vancomycin and piperacillin/tazobactam. *Clin Infect Dis.* 2017;64(5):666–674.
 165. Rutter WC, Cox JN, Martin CA, Burgess DR, Burgess DS. Nephrotoxicity during vancomycin therapy in combination with piperacillin-tazobactam or cefepime. *Antimicrob Agents Chemother.* 2017;61(2).
 166. Luther MK, Timbrook TT, Caffrey AR, Dosa D, Lodise TP, LaPlante KL. Vancomycin plus piperacillin-tazobactam and acute kidney injury in adults: a systematic review and meta-analysis. *Crit Care Med.* 2018;46(1):12–20.
 167. Schreier DJ, Kashani KB, Sakhuja A, et al. Incidence of acute kidney injury among critically ill patients with brief empiric use of antipseudomonal beta-lactams with vancomycin. *Clin Infect Dis.* 2019;68(9):1456–1462.
 168. Alshehri AM, Alzahrani MY, Abujamal MA, et al. Comparative risk of acute kidney injury following concurrent administration of vancomycin with piperacillin/tazobactam or meropenem: a systematic review and meta-analysis of observational studies. *Antibiotics (Basel).* 2022;11(4).
 169. Blevins AM, Lashinsky JN, McCammon C, Kollef M, Micek S, Juang P. Incidence of acute kidney injury in critically ill patients receiving vancomycin with concomitant piperacillin-tazobactam, cefepime, or meropenem. *Antimicrob Agents Chemother.* 2019;63(5).
 170. Avedissian SN, Pais GM, Liu J, Rhodes NJ, Scheetz MH. Piperacillin-tazobactam added to vancomycin increases risk for acute kidney injury: fact or fiction? *Clin Infect Dis.* 2020;71(2):426–432.
 171. Bellos I, Karageorgiou V, Pergialiotis V, Perrea DN. Acute kidney injury following the concurrent administration of antipseudomonal beta-lactams and vancomycin: a network meta-analysis. *Clin Microbiol Infect.* 2020;26(6):696–705.
 172. O'Callaghan K, Hay K, Lavana J, McNamara JF. Acute kidney injury with combination vancomycin and piperacillin-tazobactam therapy in the ICU: a retrospective cohort study. *Int J Antimicrob Agents.* 2020;56(1):106010.
 173. Pais GM, Liu J, Avedissian SN, et al. Lack of synergistic nephrotoxicity between vancomycin and piperacillin/tazobactam in a rat model and a confirmatory cellular model. *J Antimicrob Chemother.* 2020;75(5):1228–1236.
 174. He M, Souza E, Matvekas A, Crass RL, Pai MP. Alteration in acute kidney injury potential with the combination of vancomycin and imipenem-cilastatin/relebactam or piperacillin/tazobactam in a preclinical model. *Antimicrob Agents Chemother.* 2021;65(4).
 175. Chang J, Pais GM, Valdez K, Marianski S, Barreto EF, Scheetz MH. Glomerular function and urinary biomarker changes between vancomycin and vancomycin plus piperacillin-tazobactam in a translational rat model. *Antimicrob Agents Chemother.* 2022;66(3):e0213221.

176. Wen S, Wang C, Duan Y, et al. OAT1 and OAT3 also mediate the drug-drug interaction between piperacillin and tazobactam. *Int J Pharm.* 2018;537(1–2):172–182.
177. Miano TA, Hennessy S, Yang W, et al. Association of vancomycin plus piperacillin-tazobactam with early changes in creatinine versus cystatin C in critically ill adults: a prospective cohort study. *Intensive Care Med.* 2022;48(9):1144–1155.
178. Qian ET, Casey JD, Wright A, et al. Cefepime vs piperacillin-tazobactam in adults hospitalized with acute infection: the ACORN randomized clinical trial. *JAMA.* 2023;330(16):1557–1567.
179. Kollef MH, Juang P, Micek ST. Vancomycin/Piperacillin-tazobactam nephrotoxicity in the critically ill. *Clin Infect Dis.* 2020;70(7):1520–1521.
180. Davenport MS, Khalatbari S, Cohan RH, Dillman JR, Myles JD, Ellis JH. Contrast material-induced nephrotoxicity and intravenous low-osmolality iodinated contrast material: risk stratification by using estimated glomerular filtration rate. *Radiology.* 2013;268(3):719–728.
181. Kene M, Arasu VA, Mahapatra AK, Huang J, Reed ME. Acute kidney injury after CT in emergency patients with chronic kidney disease: a propensity score-matched analysis. *West J Emerg Med.* 2021;22(3):614–622.
182. McDonald RJ, McDonald JS, Carter RE, et al. Intravenous contrast material exposure is not an independent risk factor for dialysis or mortality. *Radiology.* 2014;273(3):714–725.
183. Heller M, Krieger P, Finefrock D, Nguyen T, Akhtar S. Contrast CT scans in the emergency department do not increase risk of adverse renal outcomes. *West J Emerg Med.* 2016;17(4):404–408.
184. Caspi O, Habib M, Cohen Y, et al. Acute kidney injury after primary angioplasty: is contrast-induced nephropathy the culprit? *J Am Heart Assoc.* 2017;6(6).
185. Hinson JS, Ehmann MR, Fine DM, et al. Risk of acute kidney injury after intravenous contrast media administration. *Ann Emerg Med.* 2017;69(5):577–586.e4.
186. Wilhelm-Leen E, Montez-Rath ME, Chertow G. Estimating the risk of radiocontrast-associated nephropathy. *J Am Soc Nephrol.* 2017;28(2):653–659.
187. Hinson JS, Al Jalbout N, Ehmann MR, Klein EY. Acute kidney injury following contrast media administration in the septic patient: a retrospective propensity-matched analysis. *J Crit Care.* 2019;51:111–116.
188. Hsu YC, Su HY, Sun CK, Liang CY, Chen TB, Hsu CW. Risk of post-contrast acute kidney injury in emergency department patients with sepsis. *Hong Kong Med J.* 2019;25(6):429–437.
189. Elias A, Aronson D. Risk of acute kidney injury after intravenous contrast media administration in patients with suspected pulmonary embolism: a propensity-matched study. *Thromb Haemost.* 2021;121(6):800–807.
190. Su TH, Hsieh CH, Chan YL, et al. Intravenous CT contrast media and acute kidney injury: a multicenter emergency department-based study. *Radiology.* 2021;301(3):571–581.
191. Ehmann MR, Mitchell J, Levin S, et al. Renal outcomes following intravenous contrast administration in patients with acute kidney injury: a multi-site retrospective propensity-adjusted analysis. *Intensive Care Med.* 2023;49(2):205–215.
192. Tergast TL, Schulte B, Griemsmann M, et al. Application of CT contrast medium is not associated with an increased risk for acute kidney injury in patients with decompensated cirrhosis. *Aliment Pharmacol Ther.* 2023;57(1):136–145.
193. McDonald JS, McDonald RJ, Comin J, et al. Frequency of acute kidney injury following intravenous contrast medium administration: a systematic review and meta-analysis. *Radiology.* 2013;267(1):119–128.
194. Ehrmann S, Quartin A, Hobbs BP, et al. Contrast-associated acute kidney injury in the critically ill: systematic review and Bayesian meta-analysis. *Intensive Care Med.* 2017;43(6):785–794.
195. Aycok RD, Westafer LM, Boxen JL, Majlesi N, Schoenfeld EM, Bannuru RR. Acute kidney injury after computed tomography: a meta-analysis. *Ann Emerg Med.* 2018;71(1):44–53.e4.
196. Oloko A, Talreja H, Davis A, et al. Does iodinated contrast affect residual renal function in dialysis patients? A systematic review and meta-analysis. *Nephron.* 2020;144(4):176–184.
197. Lopez-Ruiz A, Chandrashekar K, Juncos LA. Changing paradigms in contrast nephropathy. *J Am Soc Nephrol.* 2017;28(2):397–399.
198. Kashani K, Levin A, Schetz M. Contrast-associated acute kidney injury is a myth: we are not sure. *Intensive Care Med.* 2018;44(1):110–114.
199. Lakhali K, Ehrmann S, Robert-Edan V. Iodinated contrast medium: is there a re(n)al problem? A clinical vignette-based review. *Crit Care.* 2020;24(1):641.
200. Hiremath S, Velez JCQ. Preventing a nonexistent entity: the curious case of contrast and acute kidney injury. *Curr Opin Nephrol Hypertens.* 2020;29(1):152–160.
201. Gomes AS, Baker JD, Martin-Paredero V, et al. Acute renal dysfunction after major arteriography. *AJR Am J Roentgenol.* 1985;145(6):1249–1253.
202. Ando G, Cortese B, Russo F, et al. Acute kidney injury after radial or femoral access for invasive acute coronary syndrome management: AKI-MATRIX. *J Am Coll Cardiol.* 2017.
203. Luk L, Steinman J, Newhouse JH. Intravenous contrast-induced nephropathy—the Rise and fall of a threatening idea. *Adv Chron Kidney Dis.* 2017;24(3):169–175.
204. Schonenberger E, Martus P, Bosserdt M, et al. Kidney injury after intravenous versus intra-arterial contrast agent in patients suspected of having coronary artery disease: a randomized trial. *Radiology.* 2019;292(3):664–672.
205. Neilipovitz J, Rosenberg H, Hiremath S, et al. CJEM Debate Series: contrast-enhanced imaging should not be withheld for emergency department patients as contrast-induced acute kidney injury is very uncommon. *CJEM.* 2021;23(4):432–436.
206. Brown JR, Solomon R, Stabler ME, et al. Team-based coaching intervention to improve contrast-associated acute kidney injury: a cluster-randomized trial. *Clin J Am Soc Nephrol.* 2023;18(3):315–326.
207. James MT, Har BJ, Tyrrell BD, et al. Effect of clinical decision support with audit and feedback on prevention of acute kidney injury in patients undergoing coronary angiography: a randomized clinical trial. *JAMA.* 2022;328(9):839–849.
208. Chertow GM, Normand SL, McNeil BJ. “Renalism”: inappropriately low rates of coronary angiography in elderly individuals with renal insufficiency. *J Am Soc Nephrol.* 2004;15(9):2462–2468.
209. Szummer K, Lundman P, Jacobson SH, et al. Influence of renal function on the effects of early revascularization in non-ST-elevation myocardial infarction: data from the Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). *Circulation.* 2009;120(10):851–858.
210. Goldenberg I, Subirana I, Boyko V, et al. Relation between renal function and outcomes in patients with non-ST-segment elevation acute coronary syndrome: real-world data from the European Public Health Outcome Research and Indicators Collection Project. *Arch Intern Med.* 2010;170(10):888–895.
211. Weisbord SD, Mor MK, Hochheiser H, et al. Utilization and outcomes of clinically indicated invasive cardiac care in veterans with acute coronary syndrome and chronic kidney disease. *J Am Soc Nephrol.* 2023;34(4):694–705.
212. Ehrmann S, Aronson D, Hinson JS. Contrast-associated acute kidney injury is a myth: yes. *Intensive Care Med.* 2018;44(1):104–106.
213. Davenport MS, Perazella MA, Yee J, et al. Use of intravenous iodinated contrast media in patients with kidney disease: consensus statements from the American college of radiology and the national kidney foundation. *Kidney Med.* 2020;2(1):85–93.
214. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med.* 2001;345(19):1368–1377.
215. Yealy DM, Kellum JA, ProCESS Investigators, et al. A randomized trial of protocol-based care for early septic shock. *N Engl J Med.* 2014;370(18):1683–1693.
216. Peake SL, ARISE Investigators, ANZICS Clinical Trials Group, et al. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med.* 2014;371(16):1496–1506.
217. Mouncey PR, Osborn TM, Power GS, et al. Trial of early, goal-directed resuscitation for septic shock. *N Engl J Med.* 2015;372(14):1301–1311.
218. Humphrey H, Hall J, Sznajder I, Silverstein M, Wood L. Improved survival in ARDS patients associated with a reduction in pulmonary capillary wedge pressure. *Chest.* 1990;97(5):1176–1180.
219. Alsous F, Khamiees M, DeGirolamo A, Amoateng-Adjepong Y, Manthous CA. Negative fluid balance predicts survival in patients with

- septic shock: a retrospective pilot study. *Chest*. 2000;117(6):1749–1754.
220. Brandstrup B, Tonnesen H, Beier-Holgersen R, et al. Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens: a randomized assessor-blinded multicenter trial. *Ann Surg*. 2003;238(5):641–648.
 221. Van Biesen W, Yegenaga I, Vanholder R, et al. Relationship between fluid status and its management on acute renal failure (ARF) in intensive care unit (ICU) patients with sepsis: a prospective analysis. *J Nephrol*. 2005;18(1):54–60. <https://www.ncbi.nlm.nih.gov/pubmed/15772923>.
 222. Payen D, de Pont AC, Sakr Y, et al. A positive fluid balance is associated with a worse outcome in patients with acute renal failure. *Crit Care*. 2008;12(3):R74.
 223. Bouchard J, Soroko SB, Chertow GM, et al. Fluid accumulation, survival and recovery of kidney function in critically ill patients with acute kidney injury. *Kidney Int*. 2009;76(4):422–427.
 224. Mullens W, Abrahams Z, Francis GS, et al. Importance of venous congestion for worsening of renal function in advanced decompensated heart failure. *J Am Coll Cardiol*. 2009;53(7):589–596.
 225. Fulop T, Pathak MB, Schmidt DW, et al. Volume-related weight gain and subsequent mortality in acute renal failure patients treated with continuous renal replacement therapy. *ASAIO J*. 2010;56(4):333–337.
 226. Sutherland SM, Zappitelli M, Alexander SR, et al. Fluid overload and mortality in children receiving continuous renal replacement therapy: the prospective pediatric continuous renal replacement therapy registry. *Am J Kidney Dis*. 2010;55(2):316–325.
 227. Boyd JH, Forbes J, Nakada TA, Walley KR, Russell JA. Fluid resuscitation in septic shock: a positive fluid balance and elevated central venous pressure are associated with increased mortality. *Crit Care Med*. 2011;39(2):259–265.
 228. Grams ME, Estrella MM, Coresh J, Brower RG, Liu KD. National heart lung and blood institute acute respiratory distress syndrome network. Fluid balance, diuretic use, and mortality in acute kidney injury. *Clin J Am Soc Nephrol*. 2011;6(5):966–973.
 229. Bagshaw SM, Wald R, Barton J, et al. Clinical factors associated with initiation of renal replacement therapy in critically ill patients with acute kidney injury—a prospective multicenter observational study. *J Crit Care*. 2012;27(3):268–275.
 230. Cordemans C, De Laet I, Van Regenmortel N, et al. Aiming for a negative fluid balance in patients with acute lung injury and increased intra-abdominal pressure: a pilot study looking at the effects of PAL-treatment. *Ann Intensive Care*. 2012;2(suppl 1):S15.
 231. Cordemans C, De Laet I, Van Regenmortel N, et al. Fluid management in critically ill patients: the role of extravascular lung water, abdominal hypertension, capillary leak, and fluid balance. *Ann Intensive Care*. 2012;2(suppl 1 Diagnosis and management of intra-abdominal hypertension):S1.
 232. Heung M, Wolfgram DF, Kommareddi M, Hu Y, Song PX, Ojo AO. Fluid overload at initiation of renal replacement therapy is associated with lack of renal recovery in patients with acute kidney injury. *Nephrol Dial Transplant*. 2012;27(3):956–961.
 233. Bellomo R, Cass A, RENAL Replacement Therapy Study Investigators, et al. An observational study fluid balance and patient outcomes in the Randomized Evaluation of Normal vs. Augmented Level of Replacement Therapy trial. *Crit Care Med*. 2012;40(6):1753–1760.
 234. Vaara ST, Korhonen AM, Kaukonen KM, et al. Fluid overload is associated with an increased risk for 90-day mortality in critically ill patients with renal replacement therapy: data from the prospective FINNAKI study. *Crit Care*. 2012;16(5):R197.
 235. Legrand M, Dupuis C, Simon C, et al. Association between systemic hemodynamics and septic acute kidney injury in critically ill patients: a retrospective observational study. *Crit Care*. 2013;17(6):R278.
 236. Micek ST, McEvoy C, McKenzie M, Hampton N, Doherty JA, Kollef MH. Fluid balance and cardiac function in septic shock as predictors of hospital mortality. *Crit Care*. 2013;17(5):R246.
 237. Teixeira C, Garzotto F, Piccinni P, et al. Fluid balance and urine volume are independent predictors of mortality in acute kidney injury. *Crit Care*. 2013;17(1):R14.
 238. Acheampong A, Vincent JL. A positive fluid balance is an independent prognostic factor in patients with sepsis. *Crit Care*. 2015;19:251.
 239. Koonrangsomboon W, Khwannimit B. Impact of positive fluid balance on mortality and length of stay in septic shock patients. *Indian J Crit Care Med*. 2015;19(12):708–713.
 240. Raimundo M, Crichton S, Martin JR, et al. Increased fluid administration after early acute kidney injury is associated with less renal recovery. *Shock*. 2015;44(5):431–437.
 241. Chen KP, Cavender S, Lee J, et al. Peripheral edema, central venous pressure, and risk of AKI in critical illness. *Clin J Am Soc Nephrol*. 2016;11(4):602–608.
 242. Hjortrup PB, Haase N, Bundgaard H, et al. Restricting volumes of resuscitation fluid in adults with septic shock after initial management: the CLASSIC randomised, parallel-group, multicentre feasibility trial. *Intensive Care Med*. 2016;42(11):1695–1705.
 243. Neyra JA, Li X, Canepa-Escaro F, et al. Cumulative fluid balance and mortality in septic patients with or without acute kidney injury and chronic kidney disease. *Crit Care Med*. 2016;44(10):1891–1900.
 244. Samoni S, Vigo V, Resendiz LI, et al. Impact of hyperhydration on the mortality risk in critically ill patients admitted in intensive care units: comparison between bioelectrical impedance vector analysis and cumulative fluid balance recording. *Crit Care*. 2016;20:95.
 245. Semler MW, Wheeler AP, Thompson BT, et al. Impact of initial central venous pressure on outcomes of conservative versus liberal fluid management in acute respiratory distress syndrome. *Crit Care Med*. 2016;44(4):782–789.
 246. Balakumar V, Murugan R, Sileanu FE, Palevsky P, Clermont G, Kellum JA. Both positive and negative fluid balance may be associated with reduced long-term survival in the critically ill. *Crit Care Med*. 2017;45(8):e749–e757.
 247. Marik PE, Linde-Zwirble WT, Bittner EA, Sahatjian J, Hansell D. Fluid administration in severe sepsis and septic shock, patterns and outcomes: an analysis of a large national database. *Intensive Care Med*. 2017;43(5):625–632.
 248. Sakr Y, Rubatto Birri PN, Kotfis K, et al. Higher fluid balance increases the risk of death from sepsis: results from a large international audit. *Crit Care Med*. 2017;45(3):386–394.
 249. Silversides JA, Major E, Ferguson AJ, et al. Conservative fluid management or deresuscitation for patients with sepsis or acute respiratory distress syndrome following the resuscitation phase of critical illness: a systematic review and meta-analysis. *Intensive Care Med*. 2017;43(2):155–170.
 250. Tigabu BM, Davari M, Kebriaeezadeh A, Mojtahedzadeh M. Fluid volume, fluid balance and patient outcome in severe sepsis and septic shock: a systematic review. *J Crit Care*. 2018;48:153–159.
 251. Silversides JA, Fitzgerald E, Manickavasagam US, et al. Dere-suscitation of patients with iatrogenic fluid overload is associated with reduced mortality in critical illness. *Crit Care Med*. 2018;46(10):1600–1607.
 252. Huang AC, Lee TY, Ko MC, et al. Fluid balance correlates with clinical course of multiple organ dysfunction syndrome and mortality in patients with septic shock. *PLoS One*. 2019;14(12):e0225423.
 253. Woodward CW, Lambert J, Ortiz-Soriano V, et al. Fluid overload associates with major adverse kidney events in critically ill patients with acute kidney injury requiring continuous renal replacement therapy. *Crit Care Med*. 2019;47(9):e753–e760.
 254. Messmer AS, Zingg C, Muller M, Gerber JL, Schefold JC, Pfortmueller CA. Fluid overload and mortality in adult critical care patients—A systematic review and meta-analysis of observational studies. *Crit Care Med*. 2020;48(12):1862–1870.
 255. Cleymaet R, Scheinok TJ, Maes H, et al. Prognostic value of bioelectrical impedance analysis for assessment of fluid overload in ICU patients: a pilot study. *Anaesthesiol Intensive Ther*. 2021;53(1):10–17.
 256. Macedo E, Bouchard J, Soroko SH, et al. Fluid accumulation, recognition and staging of acute kidney injury in critically-ill patients. *Crit Care*. 2010;14(3):R82.
 257. Prowle JR, Kirwan CJ, Bellomo R. Fluid management for the prevention and attenuation of acute kidney injury. *Nat Rev Nephrol*. 2014;10(1):37–47.
 258. Perner A, Prowle J, Joannidis M, Young P, Hjortrup PB, Pettila V. Fluid management in acute kidney injury. *Intensive Care Med*. 2017;43(6):807–815.
 259. Prowle JR, Echeverri JE, Ligabo EV, Ronco C, Bellomo R. Fluid balance and acute kidney injury. *Nat Rev Nephrol*. 2010;6(2):107–115.
 260. Nisanevich V, Felsenstein I, Almog G, Weissman C, Einav S, Matot I. Effect of intraoperative fluid management on outcome after intraabdominal surgery. *Anesthesiology*. 2005;103(1):25–32.

261. Permpikul C, Tongyoo S, Viarasilpa T, Trainarongsakul T, Chakorn T, Udompanturak S. Early use of norepinephrine in septic shock resuscitation (CENSER). A randomized trial. *Am J Respir Crit Care Med*. 2019;199(9):1097–1105.
262. Vaara ST, Ostermann M, Bitker L, et al. Restrictive fluid management versus usual care in acute kidney injury (REVERSE-AKI): a pilot randomized controlled feasibility trial. *Intensive Care Med*. 2021;47(6):665–673. (In eng).
263. Myles PS, Bellomo R, Corcoran T, et al. Restrictive versus liberal fluid therapy for major abdominal surgery. *N Engl J Med*. 2018;378(24):2263–2274.
264. Meyhoff TS, Hjortrup PB, Wetterslev J, et al. Restriction of intravenous fluid in ICU patients with septic shock. *N Engl J Med*. 2022;386(26):2459–2470.
265. Shapiro NI, Douglas IS, National Heart Lung and Blood Institute Prevention and Early Treatment of Acute Lung Injury Clinical Trials Network, et al. Early restrictive or liberal fluid management for sepsis-induced hypotension. *N Engl J Med*. 2023;388(6):499–510.
266. Murphy CV, Schramm GE, Doherty JA, et al. The importance of fluid management in acute lung injury secondary to septic shock. *Chest*. 2009;136(1):102–109.
267. Lee SJ, Ramar K, Park JG, Gajic O, Li G, Kashyap R. Increased fluid administration in the first three hours of sepsis resuscitation is associated with reduced mortality: a retrospective cohort study. *Chest*. 2014;146(4):908–915.
268. Seymour CW, Cooke CR, Heckbert SR, et al. Prehospital intravenous access and fluid resuscitation in severe sepsis: an observational cohort study. *Crit Care*. 2014;18(5):533.
269. Leisman D, Wie B, Doerfler M, et al. Association of fluid resuscitation initiation within 30 minutes of severe sepsis and septic shock recognition with reduced mortality and length of stay. *Ann Emerg Med*. 2016;68(3):298–311.
270. Leisman DE, Goldman C, Doerfler ME, et al. Patterns and outcomes associated with timeliness of initial crystalloid resuscitation in a prospective sepsis and septic shock cohort. *Crit Care Med*. 2017;45(10):1596–1606.
271. Lane DJ, Wunsch H, Saskin R, et al. Association between early intravenous fluids provided by paramedics and subsequent in-hospital mortality among patients with sepsis. *JAMA Netw Open*. 2018;1(8):e185845.
272. Pruinelli L, Westra BL, Yadav P, et al. Delay within the 3-hour surviving sepsis campaign guideline on mortality for patients with severe sepsis and septic shock. *Crit Care Med*. 2018;46(4):500–505.
273. Kuttub HI, Lykins JD, Hughes MD, et al. Evaluation and predictors of fluid resuscitation in patients with severe sepsis and septic shock. *Crit Care Med*. 2019;47(11):1582–1590.
274. Taenzer AH, Patel SJ, Allen TL, et al. Improvement in mortality with early fluid bolus in sepsis patients with a history of congestive heart failure. *Mayo Clin Proc Innov Qual Outcomes*. 2020;4(5):537–541.
275. Ward MA, Kuttub HI, Tuck N, Taleb A, Okut H, Badgett RG. The effect of fluid initiation timing on sepsis mortality: a meta-analysis. *J Intensive Care Med*. 2022;37(11):1504–1511.
276. Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med*. 2021;47(11):1181–1247.
277. Vincent JL, De Backer D. Circulatory shock. *N Engl J Med*. 2013;369(18):1726–1734.
278. Hoste EA, Maitland K, Brudney CS, et al. Four phases of intravenous fluid therapy: a conceptual model. *Br J Anaesth*. 2014;113(5):740–747.
279. Malbrain M, Van Regenmortel N, Sauge B, et al. Principles of fluid management and stewardship in septic shock: it is time to consider the four D's and the four phases of fluid therapy. *Ann Intensive Care*. 2018;8(1):66.
280. Kattan E, Ospina-Tascon GA, Teboul JL, et al. Systematic assessment of fluid responsiveness during early septic shock resuscitation: secondary analysis of the ANDROMEDA-SHOCK trial. *Crit Care*. 2020;24(1):23.
281. Bissell BD, Laine ME, Thompson Bastin ML, et al. Impact of protocolized diuresis for de-resuscitation in the intensive care unit. *Crit Care*. 2020;24(1):70.
282. Silversides JA, McMullan R, Emerson LM, et al. Feasibility of conservative fluid administration and deresuscitation compared with usual care in critical illness: the Role of Active Deresuscitation after Resuscitation-2 (RADAR-2) randomised clinical trial. *Intensive Care Med*. 2022;48(2):190–200.
283. Wichmann S, Barbateskovic M, Liang N, et al. Loop diuretics in adult intensive care patients with fluid overload: a systematic review of randomised clinical trials with meta-analysis and trial sequential analysis. *Ann Intensive Care*. 2022;12(1):52.
284. Neyra JA, Lambert J, Ortiz-Soriano V, et al. Assessment of prescribed vs. achieved fluid balance during continuous renal replacement therapy and mortality outcome. *PLoS One*. 2022;17(8):e0272913.
285. Murugan R, Kerti SJ, Chang CH, et al. Association of net ultrafiltration rate with mortality among critically ill adults with acute kidney injury receiving continuous venovenous hemodiafiltration: a secondary analysis of the randomized evaluation of normal vs augmented level (renal) of renal replacement therapy trial. *JAMA Netw Open*. 2019;2(6):e195418.
286. Naorungroj T, Neto AS, Zwakman-Hessels L, et al. Mediators of the impact of hourly net ultrafiltration rate on mortality in critically ill patients receiving continuous renal replacement therapy. *Crit Care Med*. 2020;48(10):e934–e942.
287. Naorungroj T, Neto AS, Zwakman-Hessels L, et al. Early net ultrafiltration rate and mortality in critically ill patients receiving continuous renal replacement therapy. *Nephrol Dial Transplant*. 2021;36(6):1112–1119.
288. Naorungroj T, Serpa Neto A, Murugan R, Kellum JA, Bellomo R. Continuous renal replacement therapy: the interaction between fluid balance and net ultrafiltration. *Am J Respir Crit Care Med*. 2021;203(9):1199–1201.
289. Murugan R, Kerti SJ, Chang CH, et al. Association between net ultrafiltration rate and renal recovery among critically ill adults with acute kidney injury receiving continuous renal replacement therapy: an observational cohort study. *Blood Purif*. 2022;51(5):397–409.
290. O'Connor ME, Jones SL, Glassford NJ, Bellomo R, Prowle JR. Defining fluid removal in the intensive care unit: a national and international survey of critical care practice. *J Intensive Care Soc*. 2017;18(4):282–288.
291. Lumlertgul N, Murugan R, Seylanova N, McCready P, Ostermann M. Net ultrafiltration prescription survey in Europe. *BMC Nephrol*. 2020;21(1):522.
292. Murugan R, Ostermann M, Peng Z, et al. Net ultrafiltration prescription and practice among critically ill patients receiving renal replacement therapy: a multinational survey of critical care practitioners. *Crit Care Med*. 2020;48(2):e87–e97.
293. Ledoux-Hutchinson L, Wald R, Malbrain M, et al. Fluid management for critically ill patients with acute kidney injury receiving kidney replacement therapy: an international survey. *Clin J Am Soc Nephrol*. 2023.
294. Teixeira JP, Tolwani AJ. Practice variation in the prescription of volume removal with acute KRT: a call for randomized controlled trials. *Clin J Am Soc Nephrol*. 2023;18(6):702–704.
295. Teixeira JP, Zeidman A, Beaubien-Souligny W, et al. Proceedings of the 2022 UAB CRRT academy: non-invasive hemodynamic monitoring to guide fluid removal with CRRT and proliferation of extracorporeal blood purification devices. *Blood Purif*. 2023:1–23.
296. Shaw AD, Bagshaw SM, Goldstein SL, et al. Major complications, mortality, and resource utilization after open abdominal surgery: 0.9% saline compared to Plasma-Lyte. *Ann Surg*. 2012;255(5):821–829.
297. Raghunathan K, Shaw A, Nathanson B, et al. Association between the choice of IV crystalloid and in-hospital mortality among critically ill adults with sepsis. *Crit Care Med*. 2014;42(7):1585–1591.
298. Sen A, Keener CM, Sileanu FE, et al. Chloride content of fluids used for large-volume resuscitation is associated with reduced survival. *Crit Care Med*. 2017;45(2):e146–e153.
299. Weinberg L, Li M, Churilov L, et al. Associations of fluid amount, type, and balance and acute kidney injury in patients undergoing major surgery. *Anaesth Intensive Care*. 2018;46(1):79–87.
300. Zampieri FG, Ranzani OT, Azevedo LC, Martins ID, Kellum JA, Liborio AB. Lactated ringer is associated with reduced mortality and less acute kidney injury in critically ill patients: a retrospective cohort analysis. *Crit Care Med*. 2016;44(12):2163–2170.
301. Zhou F, Peng ZY, Bishop JV, Cove ME, Singbartl K, Kellum JA. Effects of fluid resuscitation with 0.9% saline versus a balanced electrolyte solution on acute kidney injury in a rat model of sepsis. *Crit Care Med*. 2014;42(4):e270–e278.

302. Wilcox CS. Regulation of renal blood flow by plasma chloride. *J Clin Invest*. 1983;71(3):726–735.
303. Bullivant EM, Wilcox CS, Welch WJ. Intrarenal vasoconstriction during hyperchloremia: role of thromboxane. *Am J Physiol*. 1989;256(1 Pt 2):F152–F157.
304. Chowdhury AH, Cox EF, Francis ST, Lobo DN. A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and plasma-lyte(R) 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg*. 2012;256(1):18–24.
305. Yunos NM, Bellomo R, Hegarty C, Story D, Ho L, Bailey M. Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA*. 2012;308(15):1566–1572.
306. Young P, Bailey M, Beasley R, et al. Effect of a buffered crystalloid solution vs saline on acute kidney injury among patients in the intensive care unit: the SPLIT randomized clinical trial. *JAMA*. 2015;314(16):1701–1710.
307. Self WH, Semler MW, Wanderer JP, et al. Balanced crystalloids versus saline in noncritically ill adults. *N Engl J Med*. 2018;378(9):819–828.
308. Semler MW, Self WH, Wanderer JP, et al. Balanced crystalloids versus saline in critically ill adults. *N Engl J Med*. 2018;378(9):829–839.
309. Zampieri FG, Machado FR, Biondi RS, et al. Effect of intravenous fluid treatment with a balanced solution vs 0.9% saline solution on mortality in critically ill patients: the BaSICS randomized clinical trial. *JAMA*. 2021;326(9):1–12.
310. Connor Jr MJ, Coopersmith CM. Does crystalloid composition or rate of fluid administration make a difference when resuscitating patients in the ICU? *JAMA*. 2021.
311. Finfer S, Micallef S, Hammond N, et al. Balanced multielectrolyte solution versus saline in critically ill adults. *N Engl J Med*. 2022;386(9):815–826.
312. Hammond Naomi E, Zampieri FG, Di TGL, et al. Balanced crystalloids versus saline in critically ill adults — a systematic review with meta-analysis. *NEJM Evidence*. 2022;1(2):EVIDoa2100010.
313. Collins MG, Fahim MA, Pascoe EM, et al. Balanced crystalloid solution versus saline in deceased donor kidney transplantation (BEST-Fluids): a pragmatic, double-blind, randomised, controlled trial. *Lancet*. 2023;402(10396):105–117.
314. Sankar J, Muralidharan J, Lalitha AV, et al. Multiple electrolytes solution versus saline as bolus fluid for resuscitation in pediatric septic shock: a multicenter randomized clinical trial. *Crit Care Med*. 2023;51(11):1449–1460.
315. Finfer S, Bellomo R, Boyce N, et al. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med*. 2004;350(22):2247–2256.
316. Brunkhorst FM, Engel C, Bloos F, et al. Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med*. 2008;358(2):125–139.
317. Myburgh JA, Finfer S, Bellomo R, et al. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med*. 2012;367(20):1901–1911.
318. Annane D, Siami S, Jaber S, et al. Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA*. 2013;310(17):1809–1817.
319. Caironi P, Tognoni G, Masson S, et al. Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med*. 2014;370(15):1412–1421.
320. Bansal M, Farrugia A, Balboni S, Martin G. Relative survival benefit and morbidity with fluids in severe sepsis—a network meta-analysis of alternative therapies. *Curr Drug Saf*. 2013;8(4):236–245.
321. Wiedermann CJ, Joannidis M. Albumin replacement in severe sepsis or septic shock. *N Engl J Med*. 2014;371(1):83.
322. Rochwerf B, Alhazzani W, Sindi A, et al. Fluid resuscitation in sepsis: a systematic review and network meta-analysis. *Ann Intern Med*. 2014;161(5):347–355.
323. Patel A, Laffan MA, Waheed U, Brett SJ. Randomised trials of human albumin for adults with sepsis: systematic review and meta-analysis with trial sequential analysis of all-cause mortality. *BMJ*. 2014;349:g4561.
324. McIntyre LA, Fergusson DA, Cook DJ, et al. Fluid resuscitation with 5% albumin versus normal saline in early septic shock: a pilot randomized, controlled trial. *J Crit Care*. 2012;27(3):317.
325. Futier E, Garot M, Godet T, et al. Effect of hydroxyethyl starch vs saline for volume replacement therapy on death or postoperative complications among high-risk patients undergoing major abdominal surgery: the FLASH randomized clinical trial. *JAMA*. 2020;323(3):225–236. <https://doi.org/10.1001/jama.2019.20833>.
326. Teixeira JP, Van Sant LM, Nielsen ND. Pharmacology and clinical use of plasma expanders. *Anaesth Intensive Care Med*. 2023;24(7):421–427.
327. Smith I, Kumar P, Molloy S, et al. Base excess and lactate as prognostic indicators for patients admitted to intensive care. *Intensive Care Med*. 2001;27(1):74–83.
328. Jung B, Rimmel T, Le Goff C, et al. Severe metabolic or mixed acidemia on intensive care unit admission: incidence, prognosis and administration of buffer therapy. A prospective, multiple-center study. *Crit Care*. 2011;15(5):R238.
329. Kraut JA, Kurtz I. Use of base in the treatment of acute severe organic acidosis by nephrologists and critical care physicians: results of an online survey. *Clin Exp Nephrol*. 2006;10(2):111–117.
330. Jung B, Jaber S. The Janus faces of bicarbonate therapy in the ICU. *Intensive Care Med*. 2020;46(3):516–518.
331. Sakhuja A, Gomez H, Kellum JA. The Janus faces of bicarbonate therapy in the ICU: con. *Intensive Care Med*. 2020;46(3):519–521.
332. Forni LG, Hodgson LE, Selby NM. The Janus faces of bicarbonate therapy in the ICU: not sure. *Intensive Care Med*. 2020;46(3):522–524.
333. Baue AE, Tragus ET, Parkins WM. Effects of sodium chloride and bicarbonate in shock with metabolic acidosis. *Am J Physiol*. 1967;212(1):54–60.
334. Cooper DJ, Herbertson MJ, Werner HA, Walley KR. Bicarbonate does not increase left ventricular contractility during L-lactic acidemia in pigs. *Am Rev Respir Dis*. 1993;148(2):317–322.
335. Kim HJ, Son YK, An WS. Effect of sodium bicarbonate administration on mortality in patients with lactic acidosis: a retrospective analysis. *PLoS One*. 2013;8(6):e65283.
336. Zhang Z, Zhu C, Mo L, Hong Y. Effectiveness of sodium bicarbonate infusion on mortality in septic patients with metabolic acidosis. *Intensive Care Med*. 2018;44(11):1888–1895.
337. Cooper DJ, Walley KR, Wiggs BR, Russell JA. Bicarbonate does not improve hemodynamics in critically ill patients who have lactic acidosis. A prospective, controlled clinical study. *Ann Intern Med*. 1990;112(7):492–498.
338. Mathieu D, Neviere R, Billard V, Fleyfel M, Wattel F. Effects of bicarbonate therapy on hemodynamics and tissue oxygenation in patients with lactic acidosis: a prospective, controlled clinical study. *Crit Care Med*. 1991;19(11):1352–1356.
339. Fang ZX, Li YF, Zhou XQ, et al. Effects of resuscitation with crystalloid fluids on cardiac function in patients with severe sepsis. *BMC Infect Dis*. 2008;8:50.
340. Forsythe SM, Schmidt GA. Sodium bicarbonate for the treatment of lactic acidosis. *Chest*. 2000;117(1):260–267.
341. Boyd JH, Walley KR. Is there a role for sodium bicarbonate in treating lactic acidosis from shock? *Curr Opin Crit Care*. 2008;14(4):379–383.
342. Jaber S, Paugam C, Futier E, et al. Sodium bicarbonate therapy for patients with severe metabolic acidemia in the intensive care unit (BICAR-ICU): a multicentre, open-label, randomised controlled, phase 3 trial. *Lancet*. 2018;392(10141):31–40.
343. Jung B, Huguet H, Molinari N, Jaber S. Sodium bicarbonate for the treatment of severe metabolic acidosis with moderate or severe acute kidney injury in the critically ill: protocol for a randomised clinical trial (BICARICU-2). *BMJ Open*. 2023;13(8):e073487.
344. Bellomo R, Ronco C, Mehta RL, et al. Acute kidney injury in the ICU: from injury to recovery: reports from the 5th Paris International Conference. *Ann Intensive Care*. 2017;7(1):49.
345. Liangos O, Wald R, O'Bell JW, Price L, Pereira BJ, Jaber BL. Epidemiology and outcomes of acute renal failure in hospitalized patients: a national survey. *Clin J Am Soc Nephrol*. 2006;1(1):43–51.
346. Ali T, Khan I, Simpson W, et al. Incidence and outcomes in acute kidney injury: a comprehensive population-based study. *J Am Soc Nephrol*. 2007;18(4):1292–1298.
347. Lo LJ, Go AS, Chertow GM, et al. Dialysis-requiring acute renal failure increases the risk of progressive chronic kidney disease. *Kidney Int*. 2009;76(8):893–899.
348. Wang AY, Bellomo R, Cass A, et al. Health-related quality of life in survivors of acute kidney injury: the prolonged outcomes study of the randomized evaluation of normal versus augmented level replacement therapy study outcomes. *Nephrology*. 2015;20(7):492–498.

349. Pannu N, James M, Hemmelgarn B, Klarenbach S, Alberta Kidney Disease N. Association between AKI, recovery of renal function, and long-term outcomes after hospital discharge. *Clin J Am Soc Nephrol*. 2013;8(2):194–202.
350. Bagshaw SM, Laupland KB, Doig CJ, et al. Prognosis for long-term survival and renal recovery in critically ill patients with severe acute renal failure: a population-based study. *Crit Care*. 2005;9(6):R700–R709.
351. Liano F, Pascual J. Epidemiology of acute renal failure: a prospective, multicenter, community-based study. Madrid Acute Renal Failure Study Group. *Kidney Int*. 1996;50(3):811–818.
352. Mehta RL, Pascual MT, Soroko S, et al. Spectrum of acute renal failure in the intensive care unit: the PICARD experience. *Kidney Int*. 2004;66(4):1613–1621.
353. Ostermann M, Bellomo R, Burdmann EA, et al. Controversies in acute kidney injury: conclusions from a kidney disease: improving global outcomes (KDIGO) conference. *Kidney Int*. 2020;98(2):294–309.
354. Bagshaw SM, Darmon M, Ostermann M, et al. Current state of the art for renal replacement therapy in critically ill patients with acute kidney injury. *Intensive Care Med*. 2017;43(6):841–854.
355. Ostermann M, Dickie H, Barrett NA. Renal replacement therapy in critically ill patients with acute kidney injury—when to start. *Nephrol Dial Transplant*. 2012;27(6):2242–2248.
356. Ostermann M, Joannidis M, Pani A, et al. Patient selection and timing of continuous renal replacement therapy. *Blood Purif*. 2016;42(3):224–237.
357. Tolwani A. Continuous renal-replacement therapy for acute kidney injury. *N Engl J Med*. 2012;367(26):2505–2514.
358. Ostermann M, Bagshaw SM, Lumlertgul N, Wald R. Indications for and timing of initiation of KRT. *Clin J Am Soc Nephrol*. 2023;18(1):113–120.
359. Seabra VF, Balk EM, Liangos O, Sosa MA, Cendoroglo M, Jaber BL. Timing of renal replacement therapy initiation in acute renal failure: a meta-analysis. *Am J Kidney Dis*. 2008;52(2):272–284.
360. Bouman CS, Oudemans-Van Straaten HM, Tijssen JG, Zandstra DF, Kesecioglu J. Effects of early high-volume continuous venovenous hemofiltration on survival and recovery of renal function in intensive care patients with acute renal failure: a prospective, randomized trial. *Crit Care Med*. 2002;30(10):2205–2211.
361. Karvellas CJ, Farhat MR, Sajjad I, et al. A comparison of early versus late initiation of renal replacement therapy in critically ill patients with acute kidney injury: a systematic review and meta-analysis. *Crit Care*. 2011;15(1):R72.
362. Payen D, Mateo J, Cavaillon JM, et al. Impact of continuous venovenous hemofiltration on organ failure during the early phase of severe sepsis: a randomized controlled trial. *Crit Care Med*. 2009;37(3):803–810.
363. Jamale TE, Hase NK, Kulkarni M, et al. Earlier-start versus usual-start dialysis in patients with community-acquired acute kidney injury: a randomized controlled trial. *Am J Kidney Dis*. 2013;62(6):1116–1121.
364. Combes A, Brechot N, Amour J, et al. Early high-volume hemofiltration versus standard care for post-cardiac surgery shock. The HEROICS study. *Am J Respir Crit Care Med*. 2015;192(10):1179–1190.
365. Wald R, Adhikari NK, Smith OM, et al. Comparison of standard and accelerated initiation of renal replacement therapy in acute kidney injury. *Kidney Int*. 2015;88(4):897–904.
366. Zarbock A, Kellum JA, Schmidt C, et al. Effect of early vs delayed initiation of renal replacement therapy on mortality in critically ill patients with acute kidney injury: the ELAIN randomized clinical trial. *JAMA*. 2016;315(20):2190–2199.
367. Gaudry S, Hajage D, Schortgen F, et al. Initiation strategies for renal-replacement therapy in the intensive care unit. *N Engl J Med*. 2016;375(2):122–133.
368. Barbar SD, Clere-Jehl R, Bourredjem A, et al. Timing of renal-replacement therapy in patients with acute kidney injury and sepsis. *N Engl J Med*. 2018;379(15):1431–1442.
369. STARRT-AKI Investigators, Canadian Critical Care Trials Group, Australian New Zealand Intensive Care Society Clinical Trials Group, et al. Timing of initiation of renal-replacement therapy in acute kidney injury. *N Engl J Med*. 2020;383(3):240–251.
370. Gaudry S, Hajage D, Martin-Lefevre L, et al. Comparison of two delayed strategies for renal replacement therapy initiation for severe acute kidney injury (AKIKI 2): a multicentre, open-label, randomised, controlled trial. *Lancet*. 2021;397(10281):1293–1300.
371. Lee K, Jeong JH, Mun CH, et al. Convection-enhanced high-flux hemodialysis. *Artif Organs*. 2007;31(8):653–658.
372. Mistry K. Dialysis disequilibrium syndrome prevention and management. *Int J Nephrol Renovascular Dis*. 2019;12:69–77.
373. Sharma S, Waikar SS. Intradialytic hypotension in acute kidney injury requiring renal replacement therapy. *Semin Dial*. 2017;30(6):553–558.
374. Russo DS, Eugenio CS, Balestrin IG, et al. Comparison of hemodynamic instability among continuous, intermittent and hybrid renal replacement therapy in acute kidney injury: a systematic review of randomized clinical trials. *J Crit Care*. 2022;69:153998.
375. Beaubien-Souligny W, Yang Y, Burns KEA, et al. Intra-dialytic hypotension following the transition from continuous to intermittent renal replacement therapy. *Ann Intensive Care*. 2021;11(1):96.
376. Douvris A, Zeid K, Hiremath S, et al. Mechanisms for hemodynamic instability related to renal replacement therapy: a narrative review. *Intensive Care Med*. 2019;45(10):1333–1346.
377. Chan RJ, Helmecci W, Canney M, Clark EG. Management of intermittent hemodialysis in the critically ill patient. *Clin J Am Soc Nephrol*. 2023;18(2):245–255.
378. Schortgen F, Soubrier N, Delclaux C, et al. Hemodynamic tolerance of intermittent hemodialysis in critically ill patients: usefulness of practice guidelines. *Am J Respir Crit Care Med*. 2000;162(1):197–202.
379. Douvris A, Malhi G, Hiremath S, et al. Interventions to prevent hemodynamic instability during renal replacement therapy in critically ill patients: a systematic review. *Crit Care*. 2018;22(1):41.
380. Schytz PA, Mace ML, Soja AM, et al. Impact of extracorporeal blood flow rate on blood pressure, pulse rate and cardiac output during haemodialysis. *Nephrol Dial Transplant*. 2015;30(12):2075–2079.
381. Trivedi HS, Kukla A, Prowant B, Lim HJ. A study of the extracorporeal rate of blood flow and blood pressure during hemodialysis. *Hemodial Int*. 2007;11(4):424–429.
382. Jardin F, Prost JF, Ozier Y, Margairaz A. Hemodialysis in septic patients: improvements in tolerance of fluid removal with concentrated albumin as the priming fluid. *Crit Care Med*. 1982;10(10):650–652. <https://www.ncbi.nlm.nih.gov/pubmed/7116885>.
383. Hryciw N, Joannidis M, Hiremath S, Callum J, Clark EG. Intravenous albumin for mitigating hypotension and augmenting ultrafiltration during kidney replacement therapy. *Clin J Am Soc Nephrol*. 2021;16(5):820–828.
384. Mustafa RA, Bdair F, Akl EA, et al. Effect of lowering the dialysate temperature in chronic hemodialysis: a systematic review and meta-analysis. *Clin J Am Soc Nephrol*. 2016;11(3):442–457.
385. Selby NM, McIntyre CW. A systematic review of the clinical effects of reducing dialysate fluid temperature. *Nephrol Dial Transplant*. 2006;21(7):1883–1898.
386. Prakash S, Garg AX, Heidenheim AP, House AA. Midodrine appears to be safe and effective for dialysis-induced hypotension: a systematic review. *Nephrol Dial Transplant*. 2004;19(10):2553–2558.
387. Henrich WL, Woodard TD, Blachley JD, Gomez-Sanchez C, Pettinger W, Cronin RE. Role of osmolality in blood pressure stability after dialysis and ultrafiltration. *Kidney Int*. 1980;18(4):480–488.
388. Macedo E, Karl B, Lee E, Mehta RL. A randomized trial of albumin infusion to prevent intradialytic hypotension in hospitalized hypoalbuminemic patients. *Crit Care*. 2021;25(1):18.
389. Mc Causland FR, Claggett B, Sabbiseti VS, Jarolim P, Waikar SS. Hypertonic mannitol for the prevention of intradialytic hypotension: a randomized controlled trial. *Am J Kidney Dis*. 2019;74(4):483–490.
390. Lynch KE, Ghassemi F, Flythe JE, et al. Sodium modelling to reduce intradialytic hypotension during haemodialysis for acute kidney injury in the intensive care unit. *Nephrology*. 2016;21(10):870–877.
391. Davenport A, Will EJ, Davison AM. Continuous vs. intermittent forms of haemofiltration and/or dialysis in the management of acute renal failure in patients with defective cerebral autoregulation at risk of cerebral oedema. *Contrib Nephrol*. 1991;93:225–233.
392. Tuchman S, Khademan ZP, Mistry K. Dialysis disequilibrium syndrome occurring during continuous renal replacement therapy. *Clin Kidney J*. 2013;6(5):526–529.

393. Osgood M, Compton R, Carandang R, Hall W, Kershaw G, Muehlischlegel S. Rapid unexpected brain herniation in association with renal replacement therapy in acute brain injury: caution in the neurocritical care unit. *Neurocritical Care*. 2015;22(2):176–183.
394. Clark WR, Turk JE, Kraus MA, Gao D. Dose determinants in continuous renal replacement therapy. *Artif Organs*. 2003;27(9):815–820.
395. Teixeira JP, Neyra JA, Tolwani A. Continuous KRT: a contemporary review. *Clin J Am Soc Nephrol*. 2023;18(2):256–269.
396. Brunet S, Leblanc M, Geadah D, Parent D, Courteau S, Cardinal J. Diffusive and convective solute clearances during continuous renal replacement therapy at various dialysate and ultrafiltration flow rates. *Am J Kidney Dis*. 1999;34(3):486–492.
397. Uchino S, Fealy N, Baldwin I, Morimatsu H, Bellomo R. Predilution vs. post-dilution during continuous veno-venous hemofiltration: impact on filter life and azotemic control. *Nephron Clin Pract*. 2003;94(4):c94–c98.
398. Friedrich JO, Wald R, Bagshaw SM, Burns KE, Adhikari NK. Hemofiltration compared to hemodialysis for acute kidney injury: systematic review and meta-analysis. *Crit Care*. 2012;16(4):R146.
399. Borthwick EM, Hill CJ, Rabindranath KS, Maxwell AP, McAuley DF, Blackwood B. High-volume haemofiltration for sepsis in adults. *Cochrane Database Syst Rev*. 2017;1(1):CD008075.
400. Clark E, Molnar AO, Joannes-Boyau O, Honore PM, Sikora L, Bagshaw SM. High-volume hemofiltration for septic acute kidney injury: a systematic review and meta-analysis. *Crit Care*. 2014;18(1):R7.
401. Bellomo R, Baldwin I, Fealy N. Prolonged intermittent renal replacement therapy in the intensive care unit. *Crit Care Resusc*. 2002;4(4):281–290. <https://www.ncbi.nlm.nih.gov/pubmed/16573441>.
402. Naka T, Baldwin I, Bellomo R, Fealy N, Wan L. Prolonged daily intermittent renal replacement therapy in ICU patients by ICU nurses and ICU physicians. *Int J Artif Organs*. 2004;27(5):380–387.
403. Edrees F, Li T, Vijayan A. Prolonged intermittent renal replacement therapy. *Adv Chron Kidney Dis*. 2016;23(3):195–202.
404. Levine Z, Vijayan A. Prolonged intermittent kidney replacement therapy. *Clin J Am Soc Nephrol*. 2023;18(3):383–391.
405. Marshall MR, Ma T, Galler D, Rankin AP, Williams AB. Sustained low-efficiency daily dialysis (SLEDD-f) for critically ill patients requiring renal replacement therapy: towards an adequate therapy. *Nephrol Dial Transplant*. 2004;19(4):877–884.
406. Abe M, Okada K, Suzuki M, et al. Comparison of sustained hemodiafiltration with continuous venovenous hemodiafiltration for the treatment of critically ill patients with acute kidney injury. *Artif Organs*. 2010;34(4):331–338.
407. Gashti CN, Salcedo S, Robinson V, Rodby RA. Accelerated venovenous hemofiltration: early technical and clinical experience. *Am J Kidney Dis*. 2008;51(5):804–810.
408. Fieghen HE, Friedrich JO, Burns KE, et al. The hemodynamic tolerability and feasibility of sustained low efficiency dialysis in the management of critically ill patients with acute kidney injury. *BMC Nephrol*. 2010;11:32.
409. Kumar VA, Yeun JY, Depner TA, Don BR. Extended daily dialysis vs. continuous hemodialysis for ICU patients with acute renal failure: a two-year single center report. *Int J Artif Organs*. 2004;27(5):371–379.
410. Kielstein JT, Kretschmer U, Ernst T, et al. Efficacy and cardiovascular tolerability of extended dialysis in critically ill patients: a randomized controlled study. *Am J Kidney Dis*. 2004;43(2):342–349.
411. Wu VC, Wang CH, Wang WJ, et al. Sustained low-efficiency dialysis versus continuous veno-venous hemofiltration for postsurgical acute renal failure. *Am J Surg*. 2010;199(4):466–476.
412. Gabriel DP, Nascimento GV, Caramori JT, Martim LC, Barretti P, Balbi AL. High volume peritoneal dialysis for acute renal failure. *Perit Dial Int*. 2007;27(3):277–282. <https://www.ncbi.nlm.nih.gov/pubmed/17468475>.
413. Ponce D, Berbel MN, Regina de Goes C, Almeida CT, Balbi AL. High-volume peritoneal dialysis in acute kidney injury: indications and limitations. *Clin J Am Soc Nephrol*. 2012;7(6):887–894.
414. Bohorques R, Rivas R, Martinez A. Continuous equilibration peritoneal dialysis (CEPD) in acute renal failure. *Perit Dial Int*. 1990;10(2):183. <https://www.ncbi.nlm.nih.gov/pubmed/2085609>.
415. Steiner RW. Continuous equilibration peritoneal dialysis in acute renal failure. *Perit Dial Int*. 1989;9(1):5–7. <https://www.ncbi.nlm.nih.gov/pubmed/2488181>.
416. Fernando SK, Finkelstein FO, Tidal PD: its role in the current practice of peritoneal dialysis. *Kidney Int Suppl*. 2006;(103):S91–S95.
417. Amerling R, Winchester JF, Ronco C. Continuous flow peritoneal dialysis: update 2012. *Contrib Nephrol*. 2012;178:205–215.
418. Ronco C, Dell'Aquila R, Bonello M, et al. Continuous flow peritoneal dialysis: a new double lumen catheter. *Int J Artif Organs*. 2003;26(11):984–990.
419. Nolph KD. Peritoneal dialysis for acute renal failure. *ASAIO (Am Soc Artif Intern Organs) Trans*. 1988;34(1):54–55. <https://www.ncbi.nlm.nih.gov/pubmed/3377911>.
420. Al Sahlawi M, Ponce D, Charytan DM, Cullis B, Perl J. Peritoneal dialysis in critically ill patients: time for a critical reevaluation? *Clin J Am Soc Nephrol*. 2023;18(4):512–520.
421. Ye Z, Wang Y, Ge L, et al. Comparing renal replacement therapy modalities in critically ill patients with acute kidney injury: a systematic review and network meta-analysis. *Crit Care Explor*. 2021;3(5):e0399.
422. Vinsonneau C, Camus C, Combes A, et al. Continuous venovenous haemodiafiltration versus intermittent haemodialysis for acute renal failure in patients with multiple-organ dysfunction syndrome: a multicentre randomised trial. *Lancet*. 2006;368(9533):379–385.
423. Mehta RL, McDonald B, Gabbai FB, et al. A randomized clinical trial of continuous versus intermittent dialysis for acute renal failure. *Kidney Int*. 2001;60(3):1154–1163.
424. Augustine JJ, Sandy D, Seifert TH, Paganini EP. A randomized controlled trial comparing intermittent with continuous dialysis in patients with ARF. *Am J Kidney Dis*. 2004;44(6):1000–1007.
425. Uehlinger DE, Jakob SM, Ferrari P, et al. Comparison of continuous and intermittent renal replacement therapy for acute renal failure. *Nephrol Dial Transplant*. 2005;20(8):1630–1637.
426. Lins RL, Elseviers MM, Van der Niepen P, et al. Intermittent versus continuous renal replacement therapy for acute kidney injury patients admitted to the intensive care unit: results of a randomized clinical trial. *Nephrol Dial Transplant*. 2009;24(2):512–518.
427. Schefold JC, von Haehling S, Pschowski R, et al. The effect of continuous versus intermittent renal replacement therapy on the outcome of critically ill patients with acute renal failure (CONVINT): a prospective randomized controlled trial. *Crit Care*. 2014;18(1):R11.
428. Rabindranath K, Adams J, Macleod AM, Muirhead N. Intermittent versus continuous renal replacement therapy for acute renal failure in adults. *Cochrane Database Syst Rev*. 2007;(3):CD003773.
429. Bagshaw SM, Berthiaume LR, Delaney A, Bellomo R. Continuous versus intermittent renal replacement therapy for critically ill patients with acute kidney injury: a meta-analysis. *Crit Care Med*. 2008;36(2):610–617.
430. Pannu N, Klarenbach S, Wiebe N, Manns B, Tonelli M, Alberta Kidney Disease N. Renal replacement therapy in patients with acute renal failure: a systematic review. *JAMA*. 2008;299(7):793–805.
431. Kellum JA, Angus DC, Johnson JP, et al. Continuous versus intermittent renal replacement therapy: a meta-analysis. *Intensive Care Med*. 2002;28(1):29–37.
432. Tonelli M, Manns B, Feller-Kopman D. Acute renal failure in the intensive care unit: a systematic review of the impact of dialytic modality on mortality and renal recovery. *Am J Kidney Dis*. 2002;40(5):875–885.
433. Schneider AG, Bellomo R, Bagshaw SM, et al. Choice of renal replacement therapy modality and dialysis dependence after acute kidney injury: a systematic review and meta-analysis. *Intensive Care Med*. 2013;39(6):987–997.
434. Nash DM, Przech S, Wald R, O'Reilly D. Systematic review and meta-analysis of renal replacement therapy modalities for acute kidney injury in the intensive care unit. *J Crit Care*. 2017;41:138–144.
435. Palevsky PM. Indications and timing of renal replacement therapy in acute kidney injury. *Crit Care Med*. 2008;36(4 suppl):S224–S228.
436. Schwenger V, Weigand MA, Hoffmann O, et al. Sustained low efficiency dialysis using a single-pass batch system in acute kidney injury—a randomized interventional trial: the Renal Replacement Therapy Study in Intensive Care Unit PatiEnts. *Crit Care*. 2012;16(4):R140.
437. Zhang L, Yang J, Eastwood GM, Zhu G, Tanaka A, Bellomo R. Extended daily dialysis versus continuous renal replacement

- therapy for acute kidney injury: a meta-analysis. *Am J Kidney Dis*. 2015;66(2):322–330.
438. Marshall MR, Creamer JM, Foster M, et al. Mortality rate comparison after switching from continuous to prolonged intermittent renal replacement for acute kidney injury in three intensive care units from different countries. *Nephrol Dial Transplant*. 2011;26(7):2169–2175.
 439. Dalbhi SA, Alorf R, Alotaibi M, et al. Sustained low efficiency dialysis is non-inferior to continuous renal replacement therapy in critically ill patients with acute kidney injury: a comparative meta-analysis. *Medicine (Baltim)*. 2021;100(51):e28118.
 440. Gabriel DP, Caramori JT, Martim LC, Barretti P, Balbi AL. High volume peritoneal dialysis vs daily hemodialysis: a randomized, controlled trial in patients with acute kidney injury. *Kidney Int Suppl*. 2008;(108):S87–S93.
 441. Ponce D, Berbel MN, Abrao JM, Goes CR, Balbi AL. A randomized clinical trial of high volume peritoneal dialysis versus extended daily hemodialysis for acute kidney injury patients. *Int Urol Nephrol*. 2013;45(3):869–878.
 442. Liu L, Zhang L, Liu GJ, Fu P. Peritoneal dialysis for acute kidney injury. *Cochrane Database Syst Rev*. 2017;12(12):CD011457.
 443. Al-Hwiesh A, Abdul-Rahman I, Finkelstein F, et al. Acute kidney injury in critically ill patients: a prospective randomized study of tidal peritoneal dialysis versus continuous renal replacement therapy. *Ther Apher Dial*. 2018;22(4):371–379.
 444. Mehta RL, Letteri JM. Current status of renal replacement therapy for acute renal failure. A survey of US nephrologists. The National Kidney Foundation Council on Dialysis. *Am J Nephrol*. 1999;19(3):377–382.
 445. Overberger P, Pesacreta M, Palevsky PM, Network VNARFT. Management of renal replacement therapy in acute kidney injury: a survey of practitioner prescribing practices. *Clin J Am Soc Nephrol*. 2007;2(4):623–630.
 446. Palevsky PM, Liu KD, Brophy PD, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. *Am J Kidney Dis*. 2013;61(5):649–672.
 447. Davenport A, Will EJ, Davison AM. Effect of renal replacement therapy on patients with combined acute renal and fulminant hepatic failure. *Kidney Int Suppl*. 1993;41:S245–S251. <https://www.ncbi.nlm.nih.gov/pubmed/8320931>.
 448. Davenport A, Finn R, Goldsmith HJ. Management of patients with renal failure complicated by cerebral oedema. *Blood Purif*. 1989;7(4):203–209.
 449. Rosner MH, Connor Jr MJ. Management of severe hyponatremia with continuous renal replacement therapies. *Clin J Am Soc Nephrol*. 2018;13(5):787–789.
 450. Yessayan LT, Szamosfalvi B, Rosner MH. Management of dysnatremias with continuous renal replacement therapy. *Semin Dial*. 2021;34(6):472–479.
 451. Morgan D, Ho K, Murray C, Davies H, Louw J. A randomized trial of catheters of different lengths to achieve right atrium versus superior vena cava placement for continuous renal replacement therapy. *Am J Kidney Dis*. 2012;60(2):272–279.
 452. Czepizak CA, O'Callaghan JM, Venus B. Evaluation of formulas for optimal positioning of central venous catheters. *Chest*. 1995;107(6):1662–1664.
 453. Parienti JJ, Thirion M, Megarbane B, et al. Femoral vs jugular venous catheterization and risk of nosocomial events in adults requiring acute renal replacement therapy: a randomized controlled trial. *JAMA*. 2008;299(20):2413–2422.
 454. Parienti JJ, Megarbane B, Fischer MO, et al. Catheter dysfunction and dialysis performance according to vascular access among 736 critically ill adults requiring renal replacement therapy: a randomized controlled study. *Crit Care Med*. 2010;38(4):1118–1125.
 455. Kelly YP, Mendu ML. Vascular access for renal replacement therapy in acute kidney injury: are nontunneled catheters the right choice? *Semin Dial*. 2019;32(5):406–410.
 456. Verma S, Palevsky PM. Prescribing continuous kidney replacement therapy in acute kidney injury: a narrative review. *Kidney Med*. 2021;3(5):827–836.
 457. Palevsky PM, Zhang JH, VA/NIH Acute Renal Failure Trial Network, et al. Intensity of renal support in critically ill patients with acute kidney injury. *N Engl J Med*. 2008;359(1):7–20.
 458. Bellomo R, Cass A, RENAL Replacement Therapy Study Investigators, et al. Intensity of continuous renal-replacement therapy in critically ill patients. *N Engl J Med*. 2009;361(17):1627–1638.
 459. Jun M, Heerspink HJ, Ninomiya T, et al. Intensities of renal replacement therapy in acute kidney injury: a systematic review and meta-analysis. *Clin J Am Soc Nephrol*. 2010;5(6):956–963.
 460. Van Wert R, Friedrich JO, Scales DC, Wald R, Adhikari NK, University of Toronto Acute Kidney Injury Research G. High-dose renal replacement therapy for acute kidney injury: systematic review and meta-analysis. *Crit Care Med*. 2010;38(5):1360–1369.
 461. Wang Y, Gallagher M, Li Q, et al. Renal replacement therapy intensity for acute kidney injury and recovery to dialysis independence: a systematic review and individual patient data meta-analysis. *Nephrol Dial Transplant*. 2018;33(6):1017–1024.
 462. Clark EG, Vijayan A. Intensive RRT for AKI: dial down your enthusiasm. *Kidney*. 2022;3(8):1439–1441.
 463. Sharma S, Kelly YP, Palevsky PM, Waikar SS. Intensity of renal replacement therapy and duration of mechanical ventilation: secondary analysis of the acute renal failure trial network study. *Chest*. 2020;158(4):1473–1481.
 464. Mc Causland FR, Asafu-Adjei J, Betensky RA, Palevsky PM, Waikar SS. Comparison of urine output among patients treated with more intensive versus less intensive RRT: results from the acute renal failure trial network study. *Clin J Am Soc Nephrol*. 2016;11(8):1335–1342.
 465. Vijayan A, Delos Santos RB, Li T, Goss CW, Palevsky PM. Effect of frequent dialysis on renal recovery: results from the acute renal failure trial network study. *Kidney Int Rep*. 2018;3(2):456–463.
 466. Klarenbach S, Manns B, Pannu N, et al. Economic evaluation of continuous renal replacement therapy in acute renal failure. *Int J Technol Assess Health Care*. 2009;25(3):331–338.
 467. Uchino S, Toki N, Takeda K, et al. Validity of low-intensity continuous renal replacement therapy. *Crit Care Med*. 2013;41(11):2584–2591.
 468. Calatzis A, Toepfer M, Schramm W, Spannagl M, Schiffel H. Citrate anticoagulation for extracorporeal circuits: effects on whole blood coagulation activation and clot formation. *Nephron*. 2001;89(2):233–236.
 469. Davenport A, Tolwani A. Citrate anticoagulation for continuous renal replacement therapy (CRRT) in patients with acute kidney injury admitted to the intensive care unit. *NDT Plus*. 2009;2(6):439–447.
 470. Legrand M, Tolwani A. Anticoagulation strategies in continuous renal replacement therapy. *Semin Dial*. 2021;34(6):416–422.
 471. Neyra JA, Yessayan L, Thompson Bastin ML, Wille KM, Tolwani AJ. How to prescribe and troubleshoot continuous renal replacement therapy: a case-based review. *Kidney*. 2021;2(2):371–384.
 472. Schneider AG, Journois D, Rimmele T. Complications of regional citrate anticoagulation: accumulation or overload? *Crit Care*. 2017;21(1):281.
 473. Tan JN, Haroon SWP, Mukhopadhyay A, et al. Hyperlactatemia predicts citrate intolerance with regional citrate anticoagulation during continuous renal replacement therapy. *J Intensive Care Med*. 2019;34(5):418–425.
 474. Apsner R, Schwarzenhofer M, Derfler K, Zauner C, Ratheiser K, Kranz A. Impairment of citrate metabolism in acute hepatic failure. *Wien Klin Wochenschr*. 1997;109(4):123–127. <https://www.ncbi.nlm.nih.gov/pubmed/9076929>.
 475. Meier-Kriesche HU, Gitomer J, Finkel K, DuBose T. Increased total to ionized calcium ratio during continuous venovenous hemodialysis with regional citrate anticoagulation. *Crit Care Med*. 2001;29(4):748–752.
 476. Kramer L, Bauer E, Joukhadar C, et al. Citrate pharmacokinetics and metabolism in cirrhotic and noncirrhotic critically ill patients. *Crit Care Med*. 2003;31(10):2450–2455.
 477. Bakker AJ, Boerma EC, Keidel H, Kingma P, van der Voort PH. Detection of citrate overdose in critically ill patients on citrate-anticoagulated venovenous haemofiltration: use of ionised and total/ionised calcium. *Clin Chem Lab Med*. 2006;44(8):962–966.
 478. Zhang W, Bai M, Yu Y, et al. Safety and efficacy of regional citrate anticoagulation for continuous renal replacement therapy in liver failure patients: a systematic review and meta-analysis. *Crit Care*. 2019;23(1):22.
 479. Slowinski T, Morgera S, Joannidis M, et al. Safety and efficacy of regional citrate anticoagulation in continuous venovenous hemodialysis in the presence of liver failure: the Liver Citrate Anticoagulation Threshold (L-CAT) observational study. *Crit Care*. 2015;19:349.

480. Sponholz C, Settmacher U, Bauer M, Kortgen A. Regional citrate anticoagulation for continuous renal replacement therapy in the perioperative care of liver transplant recipients: a single center experience. *Ther Apher Dial*. 2015;19(1):8–15.
481. Bai M, Zhou M, He L, et al. Citrate versus heparin anticoagulation for continuous renal replacement therapy: an updated meta-analysis of RCTs. *Intensive Care Med*. 2015;41(12):2098–2110.
482. Zarbock A, Kullmar M, Kindgen-Milles D, et al. Effect of regional citrate anticoagulation vs systemic heparin anticoagulation during continuous kidney replacement therapy on dialysis filter life span and mortality among critically ill patients with acute kidney injury: a randomized clinical trial. *JAMA*. 2020;324(16):1629–1639.
483. Liu C, Mao Z, Kang H, Hu J, Zhou F. Regional citrate versus heparin anticoagulation for continuous renal replacement therapy in critically ill patients: a meta-analysis with trial sequential analysis of randomized controlled trials. *Crit Care*. 2016;20(1):144.
484. Gerss J, Meersch M, Kindgen-Milles D, et al. The effect of filter lifespan during continuous renal replacement therapy in critically ill patients with acute kidney injury on the rate of new-onset infection: analysis from the RICH randomized controlled trial. *Am J Respir Crit Care Med*. 2022;206(4):511–514.
485. Tan HK, Baldwin I, Bellomo R. Continuous veno-venous hemofiltration without anticoagulation in high-risk patients. *Intensive Care Med*. 2000;26(11):1652–1657.
486. Uchino S, Fealy N, Baldwin I, Morimatsu H, Bellomo R. Continuous venovenous hemofiltration without anticoagulation. *ASAIO J*. 2004;50(1):76–80.
487. Davies H, Leslie G. Maintaining the CRRT circuit: non-anticoagulant alternatives. *Aust Crit Care*. 2006;19(4):133–138.
488. Joannidis M, Oudemans-van Straaten HM. Clinical review: patency of the circuit in continuous renal replacement therapy. *Crit Care*. 2007;11(4):218.
489. Fealy N, Aitken L, du Toit E, Lo S, Baldwin I. Faster blood flow rate does not improve circuit life in continuous renal replacement therapy: a randomized controlled trial. *Crit Care Med*. 2017;45(10):e1018–e1025.
490. Claure-Del Granado R, Mehta RL. Fluid overload in the ICU: evaluation and management. *BMC Nephrol*. 2016;17(1):109.
491. McIntyre CW, Burton JO, Selby NM, et al. Hemodialysis-induced cardiac dysfunction is associated with an acute reduction in global and segmental myocardial blood flow. *Clin J Am Soc Nephrol*. 2008;3(1):19–26.
492. Mahmoud H, Forni LG, McIntyre CW, Selby NM. Myocardial stunning occurs during intermittent haemodialysis for acute kidney injury. *Intensive Care Med*. 2017;43(6):942–944.
493. Slessarev M, Salerno F, Ball IM, McIntyre CW. Continuous renal replacement therapy is associated with acute cardiac stunning in critically ill patients. *Hemodial Int*. 2019;23(3):325–332.
494. Murugan R, Bellomo R, Palevsky PM, Kellum JA. Ultrafiltration in critically ill patients treated with kidney replacement therapy. *Nat Rev Nephrol*. 2021;17(4):262–276.
495. Neyra JA, Mehta RL, Murugan R. Fluid management during continuous renal replacement therapy: a case-based approach. *Nephron*. 2023;147(12):782–787.
496. Matzke GR, Aronoff GR, Atkinson Jr AJ, et al. Drug dosing consideration in patients with acute and chronic kidney disease—a clinical update from Kidney Disease: improving Global Outcomes (KDIGO). *Kidney Int*. 2011;80(11):1122–1137.
497. Bogard KN, Peterson NT, Plumb TJ, Erwin MW, Fuller PD, Olsen KM. Antibiotic dosing during sustained low-efficiency dialysis: special considerations in adult critically ill patients. *Crit Care Med*. 2011;39(3):560–570.
498. Heintz BH, Matzke GR, Dager WE. Antimicrobial dosing concepts and recommendations for critically ill adult patients receiving continuous renal replacement therapy or intermittent hemodialysis. *Pharmacotherapy*. 2009;29(5):562–577.
499. Monard C, Rimmele T, Ronco C. Extracorporeal blood purification therapies for sepsis. *Blood Purif*. 2019;47(suppl 3):1–14.
500. Prince LR, Whyte MK, Sabroe I, Parker LC. The role of TLRs in neutrophil activation. *Curr Opin Pharmacol*. 2011;11(4):397–403.
501. Lee S, Sirich TL, Meyer TW. Improving solute clearances by hemodialysis. *Blood Purif*. 2022;1–12.
502. Botella J, Ghezzi PM, Sanz-Moreno C. Adsorption in hemodialysis. *Kidney Int Suppl*. 2000;76:S60–S65.
503. Ghannoum M, Bouchard J, Nolin TD, Ouellet G, Roberts DM. Hemoperfusion for the treatment of poisoning: technology, determinants of poison clearance, and application in clinical practice. *Semin Dial*. 2014;27(4):350–361.
504. Cheungpasitporn W, Thongprayoon C, Zoghby ZM, Kashani K. MARS: should I use it? *Adv Chron Kidney Dis*. 2021;28(1):47–58.
505. Ronco C, Bellomo R. Hemoperfusion: technical aspects and state of the art. *Crit Care*. 2022;26(1):135.
506. Lee KC, Stadlbauer V, Jalan R. Extracorporeal liver support devices for listed patients. *Liver Transplant*. 2016;22(6):839–848.
507. Seffer MT, Cottam D, Forni LG, Kielstein JT. Heparin 2.0: a new approach to the infection crisis. *Blood Purif*. 2021;50(1):28–34.
508. Monard C, Abraham P, Schneider A, Rimmele T. New targets for extracorporeal blood purification therapies in sepsis. *Blood Purif*. 2022;1–7.
509. Sakai Y. Polymethylmethacrylate membrane with a series of serendipity. *Contrib Nephrol*. 2011;173:137–147.
510. Ding F, Song JH, Jung JY, et al. A biomimetic membrane device that modulates the excessive inflammatory response to sepsis. *PLoS One*. 2011;6(4):e18584.
511. Opal SM, Dellinger RP, Vincent JL, Masur H, Angus DC. The next generation of sepsis clinical trial designs: what is next after the demise of recombinant human activated protein C? *Crit Care Med*. 2014;42(7):1714–1721.
512. Himmelfarb J, Le P, Klenzak J, et al. Impaired monocyte cytokine production in critically ill patients with acute renal failure. *Kidney Int*. 2004;66(6):2354–2360.
513. Gist KM, Faubel S. Infection post-AKI: should we worry? *Nephron*. 2020;144(12):673–676.
514. Ronco C, Chawla L, Husain-Syed F, Kellum JA. Rationale for sequential extracorporeal therapy (SET) in sepsis. *Crit Care*. 2023;27(1):50.
515. Silver SA, Nadim MK, O'Donoghue DJ, et al. Community health care quality standards to prevent acute kidney injury and its consequences. *Am J Med*. 2020;133(5):552–560.
516. Jiesibie ZL, Tung TH, Xu QY, et al. Association of acute kidney injury with frailty in elderly population: a systematic review and meta-analysis. *Ren Fail*. 2019;41(1):1021–1027.
517. Al Rifai A, Sukul N, Wonnacott R, Heung M. Safety of arteriovenous fistulae and grafts for continuous renal replacement therapy: the Michigan experience. *Hemodial Int*. 2018;22(1):50–55.
518. Mayer KP, Joseph-Isang E, Robinson LE, Parry SM, Morris PE, Neyra JA. Safety and feasibility of physical rehabilitation and active mobilization in patients requiring continuous renal replacement therapy: a systematic review. *Crit Care Med*. 2020;48(11):e1112–e1120.
519. Macedo E, Mehta RL. Continuous dialysis therapies: core curriculum 2016. *Am J Kidney Dis*. 2016;68(4):645–657.
520. Moss M, Huang DT, National Heart Lung and Blood Institute PETAL Clinical Trials Network, et al. Early neuromuscular blockade in the acute respiratory distress syndrome. *N Engl J Med*. 2019;380(21):1997–2008.

QUESTIONS

This vignette applies to questions 1 and 2.

A 54-year-old man with diabetes and hypertension is admitted to the hospital with left lower lobe pneumonia. Despite appropriate antibacterial therapy, the following morning he is transferred to the ICU with shock and worsening respiratory distress requiring mechanical ventilation. Vital signs include temperature 38.7°C; BP 102/58 mm Hg on norepinephrine at 0.1 µg/kg/min; and HR 92 beats per min. Serum creatinine is 1.8 mg/dL (baseline 0.9 mg/dL), and lactate is 5.2 mmol/L. Arterial blood gas reveals pH 7.28, pCO₂ 42 mm Hg, and PaO₂ 128 mm Hg on 100% oxygen with positive end-expiratory pressure of 10 cm H₂O. A chest radiograph shows progression of his lung disease with interval development of patchy bilateral opacities, and he is diagnosed with acute respiratory distress syndrome. Lung protective ventilation with low tidal volumes (6 mL/kg ideal body weight) is initiated.

1. What is the most common complication of this ventilation strategy?
 - a. Increased risk of barotrauma leading to pneumothorax
 - b. Hypoxemia leading to increased risk of acute kidney injury and need for kidney replacement therapy
 - c. Respiratory acidosis due to permissive hypercapnia
 - d. Increased intrathoracic pressures resulting in impaired venous return and hypotension requiring additional IV fluids
 - e. Increased risk of ventilator dyssynchrony resulting in increased need for neuromuscular blockade

Answer: c

Rationale: ARDS is frequently associated with impaired ventilation (i.e., increased ventilatory dead space and impaired CO₂ excretion). When lung protective ventilation is used for ARDS, respiratory acidosis due to permissive hypercapnia frequently results. Especially in the setting of AKI, patients may develop severe acidemia from combined metabolic and respiratory acidosis for which supplemental bicarbonate or KRT may be indicated. Lung protective ventilation targets lower plateau airway pressures and tidal volumes and will tend to *decrease* the risk of barotrauma. As shown in the original ARDS Net trial⁸⁷, lung protective ventilation does result in somewhat more hypoxemia, but lung protective ventilation nonetheless results in decreased mortality, underlying the fact that the primary goal of mechanical ventilation for ARDS is not to completely normalize arterial oxygen or CO₂ but rather to minimize the risk of ventilator-induced lung injury while maintaining adequate (but not necessarily normal) gas exchange. Furthermore, in the ARDS Net trial, no signals for increased risk of organ dysfunction were seen in the lung protective ventilation arm, with a trend toward *improved* kidney function, consistent with animal and epidemiologic data, which suggest that injurious mechanical ventilation may increase the risk of AKI.⁸⁸⁻⁹⁰ While the positive intrathoracic pressures generated by all mechanical ventilation may decrease venous return, clinically significant hemodynamic impairment is relatively uncommon. Furthermore, lung protective ventilation does not necessarily decrease venous return more than

other ventilatory strategies, with the effect on venous return depending on the level of PEEP applied and the mean airway pressures achieved. Ventilator dyssynchrony can be a complication of the lower tidal volumes used with lung protective ventilation, but if it occurs, adjustment to the ventilator settings or increased sedation should be tried before implementation of neuromuscular blockade. Though pharmacologic paralysis is necessary in some patients with ARDS, a large multicenter RCT⁵²⁰ demonstrated no mortality benefit to protocolized neuromuscular blockade in ARDS when compared with standard light sedation used in modern practice.

2. Despite appropriate treatment and lung protective ventilation, the patient's hypoxemia worsens and, ultimately, he is initiated on venovenous extracorporeal membrane oxygenation (VV-ECMO) on ICU Day 3. With ECMO, his gas exchange improves, allowing for implementation of "rest" settings on the ventilator. In addition, his shock resolves, allowing him to come off vasopressor therapy; his lactate normalizes; and his AKI improves, with serum creatinine returning to his baseline of 0.9 mg/dL. However, on ECMO Day 5, he develops a leukocytosis, with white blood cell count of 26,000/mm³, and, though he remains off of vasopressors, develops recurrent AKI. Blood cultures are drawn and bronchoscopy with bronchoalveolar lavage (BAL) is performed. On ECMO Day 8 blood and BAL cultures both grow *Pseudomonas aeruginosa*. However, despite appropriate antimicrobial coverage, his AKI worsens, with serum creatinine increasing to 3.1 mg/dL. Despite high-dose IV furosemide, his urine output drops to 346 mL per 24 h, resulting in a net positive fluid balance of >2 L per day. Potassium is 5.6 mEq/L and plasma-free hemoglobin is 34 mg/dL (upper limit of normal 30 mg/dL). Which of the following is the next best step in management?
 - a. Hold off on KRT initiation as the subgroup of ECMO patients in the STARRT-AKI (Standard versus Accelerated Initiation of Renal Replacement Therapy in Acute Kidney Injury) trial had no benefit to early KRT.
 - b. Initiate intermittent hemodialysis (IHD) because it has been shown in trials to be superior to CRRT for management of fluid balance in ICU patients with AKI.
 - c. Obtain renal ultrasound, urinalysis, and fractional excretion of sodium (FENa) because sepsis is unlikely to cause AKI in the absence of hypotension requiring vasopressors.
 - d. Recommend ECMO circuit change as plasma-free hemoglobin above the upper limit of normal implicates excess hemolysis as the cause of AKI.
 - e. Initiate standard-dose continuous KRT (CRRT) for volume management as positive fluid balance is associated with poor outcomes in patients on ECMO.

Answer: e

Rationale: Though experts⁶⁶ recommend against the *preemptive* use of KRT in patients supported with ECMO, fluid overload in these patients is independently associated with an increased risk of mortality⁵⁹⁻⁶⁴ and timely KRT should be

considered in ECMO patients with AKI because they are unlikely to tolerate prolonged oliguria in the setting of cardiopulmonary failure. In this case, given the failure to respond to high-dose loop diuretic resulting in positive fluid balance and some degree of hyperkalemia, initiation of KRT at this time appears reasonable. RCT data to guide the initiation of KRT in specifically ECMO patients are lacking because patients requiring ECMO were not included in STARTRT-AKI or the other recent RCTs analyzing the timing of KRT initiation in the ICU. There are no compelling data to suggest that CRRT results in superior long-term outcomes, such as survival or renal recovery, relative to IHD when used in patients in the ICU with AKI. However, CRRT has been unequivocally shown to be superior in treating or preventing fluid accumulation, as the continuous modality affords 24 hours per day (rather than 4 hours every 1–2 days) to carry out gradual ultrafiltration. As such, where both modalities are available, the preferential use of CRRT over IHD in patients requiring ECMO is reasonable. With the possible exception of FENa, the diagnostic tests recommended in answer c are reasonable. However, sepsis can cause AKI without hypotension, especially in the setting of unequivocal sepsis, such as this scenario with bacteremia with a compelling pathogen, prominent leukocytosis, and new organ dysfunction. While ischemic and septic AKI often coexist in critically ill humans, pathophysiologic studies demonstrate that septic and ischemic AKI are distinct entities as macrovascular renal blood flow may be normal or *high* early in the course of septic AKI. In general, FENa performs poorly in differentiating prerenal azotemia from ATN in the ICU,¹⁰⁶ and, as pigment nephropathy can cause ATN characteristically with a low FENa, it may be especially misleading in this case where some hemolysis is present. However, while the plasma-free hemoglobin is high, it is only marginally elevated in this case. The plasma-free hemoglobin level at which circuit change is necessary is poorly defined by the literature, and the decision to perform an ECMO circuit exchange is complex, but typically much higher elevations in plasma-free hemoglobin (e.g., several times the upper limit of normal) are necessary to justify the potential risk and expense of a circuit change.

3. A 38-year-old woman with cirrhosis is admitted for hepatic encephalopathy (HE) triggered by spontaneous bacterial peritonitis (SBP). She is admitted to the ward and treated with ceftriaxone, IV albumin, and lactulose, with initial improvement in her mental status. On Hospital Day 4 she develops recurrent deterioration in mental status, tachypnea, and AKI with serum creatinine rising within 48 hours to 2.1 mg/dL (above a baseline of 0.6) for which nephrology is consulted and IV albumin is restarted. Vital signs include BP 84/54 with mean arterial pressure (MAP) 64 mm Hg and oxygen saturation of 92% of 4 L/min of oxygen. A chest radiograph shows a new right middle lobe opacification for which her antibiotics are broadened to treat nosocomial pneumonia. Examination reveals jaundice, grade 3 HE, moderately tense abdomen with obvious ascites, and 3 to 4+ generalized pitting edema. Laboratory values include total bilirubin 6.8 mg/dL, serum albumin 3.2 g/dL, INR 2.6, and lactate 1.8 mmol/L. Urine studies include FENa of <0.1%, bland urinalysis, and undetectable urine protein

and urine albumin. Renal ultrasound is normal, and she has had no nephrotoxin exposure. What is the next most appropriate step to treat her AKI?

- a. Stop IV albumin and transfer the patient to the ICU for initiation of norepinephrine targeted to a MAP of 80 to 85 mm Hg for suspected hepatorenal syndrome-AKI (HRS-AKI).
- b. To avoid ICU transfer, she should be placed on continuous pulse oximetry and telemetry monitoring and started on terlipressin on the ward.
- c. Given the FENa <1%, she likely has prerenal azotemia and therefore should be continued on IV albumin at a dose of 20 to 40 gm IV daily.
- d. Given that she has a new pneumonia, she does not meet diagnostic criteria for HRS-AKI and therefore is not a candidate for vasoconstrictor therapy, but she should be transferred to the ICU as she is likely to have cerebral edema.
- e. Given her AKI, therapeutic paracentesis is contraindicated regardless of the severity of her ascites, but since HRS-AKI is the most common cause of AKI in patients with liver disease, she should be started on octreotide and midodrine for her HRS-AKI if she fails to improve with 48 hours of albumin 1 mg/kg IV daily.

Answer: a

Rationale: Though HRS-AKI is overall less common than ATN or prerenal azotemia as a cause of AKI in the setting of cirrhosis,¹⁰³ this patient is likely to have HRS-AKI. She meets nearly all the formal criteria for HRS-AKI^{99,105} including AKI by KDIGO criteria, cirrhosis with ascites, absence of shock or nephrotoxin exposure, and unremarkable urine studies and renal ultrasound. Additional clinical features that strongly suggest HRS-AKI include low MAP and very low FENa (with <0.2% or less being suggestive of HRS-AKI and *not* prerenal azotemia). Though the absence of active infection or sepsis was previously required to diagnose HRS, the newer criteria only preclude a diagnosis of HRS-AKI in the setting of septic shock, as it is now better recognized that infection may trigger HRS-AKI that persists despite appropriate treatment of the infection. Traditional criteria also require failure to respond to 48 hours of IV albumin therapy. However, while a trial of IV albumin is appropriate in most patients with AKI and cirrhosis, assessment of volume status before starting albumin is prudent and albumin should not be given to patients like this one with overt volume overload or significant hypoxemia.^{100-102,122} In addition, albumin does not treat HRS-AKI. Given this presentation is overall highly suggestive of HRS-AKI, delaying vasoconstrictor therapy for 48 hours to evaluate response to albumin may be harmful as response to vasoconstrictors appears best when initiated early.^{99,108,113} She has two contraindications to terlipressin therapy including hypoxemia of any severity and multiorgan dysfunction, with AKI with serum creatinine ≥ 2 mg/dL, grade 3 HE, and coagulopathy with INR ≥ 2.5 (i.e., acute-on-chronic liver failure [ACLF] grade ≥ 3). As such, given that octreotide and midodrine are generally ineffective for HRS, the best option is transfer to the ICU for norepinephrine therapy, which, though not studied in as many total patients, has been shown in multiple studies to have similar efficacy for HRS-AKI as terlipressin.^{99,108,113} Cerebral edema is a potential lethal complication of ALF, being present in a substantial minority

of patients with ALF and grade 3 HE and the majority of ALF patients with grade 4 HE.¹²³ Cerebral edema is characteristically absent in cirrhosis.^{126,127} With new multiorgan dysfunction in the setting of cirrhosis, this patient meets criteria for ACLF, which may occasionally be complicated by cerebral edema, but it is not characteristic of ACLF. Finally, the role of intraabdominal hypertension in HRS-AKI is being increasingly appreciated and, though some caution may be prudent (e.g., limitation of volume removed to ≤ 5 L), careful therapeutic paracentesis is not contraindicated in the setting of kidney dysfunction and may have some potential to *improve* renal function via decreased intraabdominal pressure and increased renal perfusion pressure.^{118-121,139}

4. A 62-year-old man with AKI secondary to septic shock has been initiated on CRRT. He is mechanically ventilated and requires two vasopressors for blood pressure support. His weight is 80 kg and CRRT dose is 25 mL/kg/h. His urine output is 20 mL/h. Within 12 hours, his serum creatinine has decreased from 4.5 mg/dL to 3.8 mg/dL, potassium has decreased from 6.5 mEq/L to 5.2 mEq/L, and bicarbonate has increased from 16 mEq/L

to 22 mEq/L. What is the appropriate next step in the management of this patient?

- a. Decrease the CRRT dose to 20 mL/kg/h.
- b. Increase the CRRT dose to 35 mL/kg/h.
- c. Maintain the current CRRT dose of 25 mL/kg/h.
- d. Discontinue CRRT.

Answer: c

Rationale: The appropriate next step in this patient's management is to maintain the current CRRT dose of 25 mL/kg/h. The recommended minimum dose of CRRT is 20 to 25 mL/kg/h. However, to account for treatment interruptions and the natural decline in filter efficiency over time, guidelines suggest prescribing 25 to 30 mL/kg/h of effluent dose.^{15,395,471} The patient's serum creatinine has decreased since initiation of CRRT, and he has improving serum potassium and bicarbonate levels, indicating that the current dose is effectively removing solutes from the blood. Increasing the CRRT dose may lead to excessive removal of solutes and has not been shown to improve outcomes. Decreasing the CRRT dose may not be sufficient to adequately remove solutes. Discontinuing CRRT at this time may not be appropriate, given the ongoing renal injury, oliguria, and need for continued support.