



## VARYING IMMIGRATION RATE

**1.1 RADS with averaging**

Here we show the difference between rank abundance distributions calculated using two different samplings from the simulations..



(a) From snapshot of final simulation state



(b) Abundances averaged over final 1000 iterations

Figure 1.1: **Average rank abundance distributions** over one hundred simulation runs, for nine different pair values of immigration rate<sub>2</sub> and habitat destruction. Each distribution is calculated using the mean relative abundance of the ranked species. The different colours correspond to different MAI ratios: red = 0.0; green = 0.5; blue = 1.0.

## 1.2 Stationarity testing

To consider stationarity we look at three types of dynamics 1) from our simulation, 2) a random walk and 3) with abundances at each iteration drawn independently from a normal distribution. The means and variances were chosen to be by a comparable size. These are plotted in figure 1.2. We also indicate the results of applying the ADF test for stationarity to the whole timeseries. A value of less than  $-2.86$  means we reject the null hypothesis of non-stationarity at 5% confidence.






Figure 1.2: Three types of dynamics generated using IBM simulation with high IR; a random walk without drift; and independent sampling from a normal distribution.




(a) Sample size = 1000 iterations




(b) Sample size = 5000 iterations

Figure 1.3: The z-statistic used to test the null hypothesis that sample means are drawn from the stationary distribution. Each dot indicates a sample from the dynamics, which is tested.

### **1.3 New results: without vegetarian predators**



(a) Sample size = 1000 iterations



(b) Sample size = 5000 iterations

Figure 1.4: The effect of using different sample sizes on the sample mean and standard deviation. Dynamics generated using IBM simulation model with low and high IR (green and red respectively).




Figure 1.5: **Summary heat maps:** Each heat map shows the value of a certain response metric across a 2-dimensional slice of parameter space. The parameters varied are immigration rate  $IR$  (y-axis) and percentage habitat destruction  $HL$  (x-axis). Each row of plots corresponds to a different MAI ratio as labelled. To construct the heatmaps one hundred repeat simulations were run for each combination of parameter values, with each simulation using a different underlying network. The mean value of the response metrics is taken over the hundred repeats. Therefore each pixel shows an estimate of the expectation value of the metric at those parameter values. The left column shows the expected number of species that are extinct at the end of a simulation; the central column shows the expected biomass (total number of individuals) at the end of a simulation; and the right column shows the expected temporal variability (coefficient of variation of total biomass) of the dynamics during the final thousand iterations of a simulation. The latter is used as a proxy for stability (see text).

Figure 1.6: **Summary heat maps:** Each heat map shows the value of a certain response metric across a 2-dimensional slice of parameter space. The parameters varied are immigration rate  $IR$  (y-axis) and percentage habitat destruction  $HL$  (x-axis). Each row of plots corresponds to a different MAI ratio as labelled. To construct the heatmaps one hundred repeat simulations were run for each combination of parameter values, with each simulation using a different underlying network. The mean value of the response metrics is taken over the hundred repeats. Therefore each pixel shows an estimate of the expectation value of the metric at those parameter values. The left column shows the expected number of species that are extinct at the end of a simulation; the central column shows the expected biomass (total number of individuals) at the end of a simulation; and the right column shows the expected temporal variability (coefficient of variation of total biomass) of the dynamics during the final thousand iterations of a simulation. The latter is used as a proxy for stability (see text).



## BIBLIOGRAPHY

- [1] D. AI, P. DESJARDINS-PROULX, C. CHU, AND G. WANG, *Immigration, local dispersal limitation, and the repeatability of community composition under neutral and niche dynamics*, (2012).
- [2] M. W. CADOTTE, *Dispersal and species diversity: A meta-analysis*, *The American Naturalist*, 167 (2006), pp. 913–924.
- [3] R. CONDIT, R. A. CHISHOLM, AND S. P. HUBBELL, *Thirty years of forest census at barro colorado and the importance of immigration in maintaining diversity*, (2012).
- [4] K. L. DRURY, J. D. SUTER, J. B. RENDALL, A. M. KRAMER, AND J. M. DRAKE, *Immigration can destabilize tri-trophic interactions: implications for conservation of top predators*, *Theoretical Ecology*, (2015), pp. 1–12.
- [5] E. S. EVELEIGH, K. S. MCCANN, P. C. MCCARTHY, S. J. POLLOCK, C. J. LUCAROTTI, B. MORIN, G. A. MCDUGALL, D. B. STRONGMAN, J. T. HUBER, J. UMBANHOWAR, ET AL., *Fluctuations in density of an outbreak species drive diversity cascades in food webs*, *Proceedings of the National Academy of Sciences*, 104 (2007), pp. 16976–16981.
- [6] B. L. FOSTER, T. L. DICKSON, C. A. MURPHY, I. S. KAREL, AND V. H. SMITH, *Propagule pools mediate community assembly and diversity-ecosystem regulation along a grassland productivity gradient*, *Journal of Ecology*, 92 (2004), pp. 435–449.
- [7] R. P. FRECKLETON, A. R. WATKINSON, AND M. REES, *Measuring the importance of competition in plant communities*, *Journal of ecology*, 97 (2009), pp. 379–384.
- [8] N. M. HADDAD, L. A. BRUDVIG, J. CLOBERT, K. F. DAVIES, A. GONZALEZ, R. D. HOLT, T. E. LOVEJOY, J. O. SEXTON, M. P. AUSTIN, C. D. COLLINS, ET AL., *Habitat fragmentation and its lasting impact on earth's ecosystems*, *Science Advances*, 1 (2015), p. e1500052.
- [9] I. HANSKI, G. A. ZURITA, M. I. BELLOCQ, AND J. RYBICKI, *Species–fragmented area relationship*, *Proceedings of the National Academy of Sciences*, 110 (2013), pp. 12715–12720.
- [10] C. KLAUSMEIER, *Habitat destruction and extinction in competitive and mutualistic meta-communities*, *Ecology Letters*, 4 (2001), pp. 57–63.

## BIBLIOGRAPHY

---

- [11] N. M. KORFANTA, W. D. NEWMARK, AND M. J. KAUFFMAN, *Long-term demographic consequences of habitat fragmentation to a tropical understory bird community*, Ecology, 93 (2012), pp. 2548–2559.
- [12] M. LOREAU AND N. MOUQUET, *Immigration and the maintenance of local species diversity*, The American Naturalist, 154 (1999), pp. 427–440.
- [13] N. MOUQUET, P. LEADLEY, J. MÉRIGUET, AND M. LOREAU, *Immigration and local competition in herbaceous plant communities: a three-year seed-sowing experiment*, Oikos, 104 (2004), pp. 77–90.
- [14] T. PUETTKER, A. A. BUENO, C. DOS SANTOS DE BARROS, S. SOMMER, AND R. PARDINI, *Immigration rates in fragmented landscapes-empirical evidence for the importance of habitat amount for species persistence*, PLoS One, 6 (2011), p. e27963.
- [15] W. J. RIPPLE AND R. L. BESCHTA, *Trophic cascades in yellowstone: The first 15 years after wolf reintroduction*, Biological Conservation, 145 (2012), pp. 205–213.
- [16] A. TEYSSÈDRE AND A. ROBERT, *Contrasting effects of habitat reduction, conversion and alteration on neutral and non neutral biological communities*, Oikos, 123 (2014), pp. 857–865.
- [17] J. M. TYLIANAKIS, T. TSCHARNTKE, AND O. T. LEWIS, *Habitat modification alters the structure of tropical host–parasitoid food webs*, Nature, 445 (2007), pp. 202–205.
- [18] B. H. WARREN, D. SIMBERLOFF, R. E. RICKLEFS, R. AGUILÉE, F. L. CONDAMINE, D. GRAVEL, H. MORLON, N. MOUQUET, J. ROSINDELL, J. CASQUET, ET AL., *Islands as model systems in ecology and evolution: prospects fifty years after macarthur-wilson*, Ecology letters, 18 (2015), pp. 200–217.
- [19] C. E. ZARTMAN AND H. E. NASCIMENTO, *Are habitat-tracking metacommunities dispersal limited? inferences from abundance-occupancy patterns of epiphylls in amazonian forest fragments*, Biological Conservation, 127 (2006), pp. 46–54.