

Multifractal analysis of measures arising from random substitutions

ANDREW MITCHELL AND ALEX RUTAR

ABSTRACT. We study regularity properties of frequency measures arising from random substitutions, which are a generalisation of (deterministic) substitutions where the substituted image of each letter is chosen independently from a fixed finite set. In particular, for a natural class of such measures, we derive a closed-form analytic formula for the L^q -spectrum and prove that the multifractal formalism holds. This provides an interesting new class of measures satisfying the multifractal formalism. More generally, we establish results concerning the L^q -spectrum of a broad class of frequency measures. We introduce a new notion called the *inflation word L^q -spectrum* of a random substitution and show that this coincides with the L^q -spectrum of the corresponding frequency measure for all $q \geq 0$. As an application, we obtain closed-form formulas under separation conditions and recover known results for topological and measure theoretic entropy.

CONTENTS

1. Introduction	2
1.1. Entropy and L^q -spectra	4
1.2. Random substitutions	5
1.3. Statement and discussion of main results	5
1.4. Discussion and further work	9
2. Preliminaries	9
2.1. Symbolic notation	10
2.2. Dynamics, entropy and dimension	10
2.3. L^q -spectra and smoothness	12
2.4. Multifractal spectrum and multifractal formalism	14
2.5. Random substitutions and frequency measures	16
2.6. Primitive random substitutions	18
2.7. Compatible random substitutions	18
2.8. Frequency measures	20
2.9. Separation conditions and recognisability	21
3. L^q -spectra of frequency measures	23
3.1. Inflation word L^q -spectra	23
3.2. L^q -spectra for non-negative q	26

2020 *Mathematics Subject Classification.* 37B10, 37C45, 52C23.

Key words and phrases. random substitution, multifractal analysis, L^q -spectrum, multifractal formalism.

3.3. L^q -spectra for negative q : lower bounds	29
3.4. Proof of general bounds for the L^q spectrum	31
3.5. L^q -spectra for negative q under recognisability	32
3.6. Recovering entropy from the L^q -spectrum	33
4. Recognisability and the multifractal formalism	34
4.1. Non-typical local dimensions	34
4.2. Proof of the multifractal formalism	40
5. Examples, counterexamples and applications	41
5.1. Failure of bounds for $q < 0$ without recognisability	41
5.2. Examples with recognisability	43
5.3. Examples without recognisability	46
Acknowledgements	47
References	47

1. INTRODUCTION

A *substitution* is a combinatorial object consisting of a finite alphabet \mathcal{A} along with a set of *transformation rules*. The theory of substitutions, along with statistical properties of the system under repeated iteration, is a large and actively researched field at the interface of combinatorics and symbolic dynamics. A thorough introduction to the statistical properties and dynamics of substitutions can be found in [2, 25]. Associated with a (deterministic) substitution is a *frequency measure*, which encodes the frequency of subwords under repeated iteration. Notably, the corresponding subshift supporting this measure has zero topological entropy, and the frequency measure is the unique ergodic measure supported on this subshift.

Random substitutions are a generalisation of (deterministic) substitutions [13] where we apply a transformation rule to each letter randomly and independently chosen from a finite set of possibilities. Similarly to the deterministic case, subshifts associated with random substitutions support ergodic frequency measures which capture the expected rate of occurrence of subwords under random iteration. But in contrast to the deterministic case, the corresponding subshift typically has positive topological entropy and supports uncountably many ergodic measures. Random substitutions include examples exhibiting deterministic behaviour, while also including examples which are subshifts of finite type. Moreover, there is a large amount of intermediate behaviour: subshifts of random substitutions can simultaneously exhibit long range correlation [3] (an indication of order) and positive topological entropy [14] (an indication of disorder).

In this paper, we study the fine scaling properties of frequency measures associated with random substitutions from the perspective of *multifractal analysis*. This perspective is relevant in a wide variety of contexts, such as the geometry of fractal sets and measures and in dynamical systems, with typical applications to geometric

measure theory and number theory. In our setting, our primary objects of study are the L^q -spectrum, which is a parametrised family of quantities which capture the inhomogeneous scaling properties of a measure, and the *local dimension*, which capture the exponential growth rate of a measure around a point. The L^q -spectrum and local dimensions are related through a heuristic relationship known as the *multifractal formalism* [18]. It is an important and well-studied question to determine settings in which the multifractal formalism holds, and to determine qualitative conditions describing its failure.

Much of the work on multifractal analysis has been done in the setting of local dimensions of self-similar measures (for some examples, see [1, 10, 11, 19, 29]) and Birkhoff sums of potentials in dynamical systems with a finite type property (see, for example, [8, 24] and the reference therein). As a notable recent example, in P. Shmerkin's recent proof of the Furstenberg $\times 2 \times 3$ conjecture [29], he computes the L^q -spectrum of a large class of dynamically self-similar measures and relates such results to the multifractal analysis of slices of sets. This information about L^q -spectra also implies L^p -smoothness properties in the question of absolute continuity of Bernoulli convolutions (see [30] for some background on this classic problem). For more detail on the geometry of measures and multifractal analysis, we refer the reader to the foundational work by L. Olsen [22] and the classic texts of K. Falconer [7] and Ya. Pesin [23].

Returning to our setting, substitution dynamical systems have characteristic features of (dynamical) self-similarity, but in many cases are far from being ergodic measures on shifts of finite type. More generally, frequency measures provide a rich family of shift-invariant ergodic measures which exhibit interesting and unique properties in symbolic dynamics in a natural way. For example, it was proved in [15, Theorem 4.8] that for a certain class of random substitutions, the corresponding subshift supports a frequency measure that is the unique ergodic measure of maximal entropy. However, this measure is not a Gibbs measure with respect to the zero potential, and therefore the system does not satisfy the common *specification property*, which is a well-known strategy for proving intrinsic ergodicity of symbolic dynamical systems (see [5] and the references therein). Moreover, there are examples of random substitutions such that the corresponding subshift supports multiple ergodic measures of maximal entropy [17, Example 6.5]. More generally, many key properties of frequency measures associated with random substitutions are poorly understood.

In this paper, we derive symbolic expressions for the L^q -spectrum of frequency measures associated with random substitutions under certain weak assumptions. Then under an additional assumption (recognisability), we prove a closed-form analytic expression for the L^q -spectrum and a variational formula which together imply the multifractal formalism. We emphasise that this class of examples has novel properties not witnessed before: in general, the unique frequency measure of maximal dimension is not a Gibbs measure with respect to the zero potential and the

corresponding subshift is not sofic. The techniques and results provide important new perspectives on the geometry and dynamics of the respective measures.

1.1. Entropy and L^q -spectra. For a Borel probability measure in a compact metric space, the L^q -spectrum is a well-studied quantity which encodes the scaling properties of the measure, in a weak sub-exponential sense. Specifically, the L^q -spectrum of μ is given by

$$\tau_\mu(q) = \liminf_{r \rightarrow 0} \frac{\log \sup \sum_i \mu(B(x_i, r))^q}{\log r}.$$

where the supremum is taken over $2r$ -separated subsets $\{x_i\}_i$ of the support of μ .

The L^q -spectrum encodes information about the local scaling of the measure μ . We define the *local dimension* of μ at x by

$$\dim_{\text{loc}}(\mu, x) = \lim_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r}$$

when the limit exists. We then define the *multifractal spectrum* of μ by

$$f_\mu(\alpha) = \dim_{\text{H}} \{x \in X : \dim_{\text{loc}}(\mu, x) = \alpha\}.$$

In general, the structure of the set of local dimensions can be very complex—for example, the level sets are often dense uncountable subsets of the support of μ . However, the “multifractal miracle” is the phenomenon that, even though the level sets are very complex, the multifractal spectrum is often a concave analytic function of α .

In fact, the multifractal spectrum and the L^q -spectrum are related through a heuristic relationship called the *multifractal formalism* [18], which speculates that under certain regularity assumptions, the multifractal spectrum is given by the *concave conjugate* of the L^q -spectrum, that is the quantity

$$\tau_\mu^*(\alpha) = \inf_{q \in \mathbb{R}} (q\alpha - \tau_\mu(q)).$$

Generally speaking, $\tau_\mu^*(\alpha) \geq f_\mu(\alpha)$ [19]: in particular, the slopes of the asymptotes of the L^q -spectrum bound the exponential scaling of measures of balls $B(x, r)$ uniformly for all $x \in \text{supp } \mu$.

In our specific setting, where our metric space is the two-sided shift $\mathcal{A}^{\mathbb{Z}}$ and the measure μ is ergodic, the local dimension is precisely the scaling rate of the *information function* of μ . In fact, the Shannon–McMillan–Breiman theorem states that the local dimension of the measure (with an appropriate choice of the metric) is almost surely the entropy of the measure. Thus the L^q -spectrum provides *uniform* control over the scaling rate of the information function. More detail about these notions are given in [Section 2](#).

1.2. Random substitutions. A (deterministic) substitution is a rule which replaces each symbol in a finite or infinite string over an alphabet \mathcal{A} with a finite word over the same alphabet. Random substitutions generalise this notion by substituting a randomly chosen word (according to a fixed finite distribution) independently for each letter. We can also think of a random substitution as a (deterministic) set-valued substitution ϑ , together with a choice of probabilities.

For example, given $p \in (0, 1)$, the *random Fibonacci substitution* ϑ_p is defined by

$$\vartheta_p: \begin{cases} a \mapsto \begin{cases} ab & \text{with probability } p, \\ ba & \text{with probability } 1 - p, \end{cases} \\ b \mapsto a. \end{cases}$$

To a given (primitive) random substitution ϑ_P , one can canonically associate a subshift X_ϑ of the two-sided shift $\mathcal{A}^{\mathbb{Z}}$ along with an ergodic *frequency measure* μ_P , which quantifies the relative occurrence of a given word under repeated application of the random substitution.

As highlighted in the introduction, primitive random substitutions give rise to subshifts and measures with a wide variety of properties. As a result, we will impose additional conditions.

Our main assumption, which we call *compatibility*, asserts that for each $a \in \mathcal{A}$, the number of occurrences of each $b \in \mathcal{A}$ is identical in every possible substituted image of a . For example, the random Fibonacci substitution is compatible since in all the possible images of a , a occurs once and b occurs once. The key feature of compatibility is that the one can define a deterministic *substitution matrix*, such that the Perron–Frobenius eigenvalue is the asymptotic growth rate of lengths of words, and the corresponding right eigenvector encodes the asymptotic frequency with which the individual letters appear. Compatibility is a common assumption: for example, it is assumed in the main results of [3, 14, 21, 27].

Another standard assumption is that of *recognisability*, which heuristically states that each element of the subshift is the *unique* image of another element of the subshift. Recognisability precludes the existence of periodic points [27] and is one of the assumptions required to establish intrinsic ergodicity in [15]. It is also assumed in the main results of [12, 21].

1.3. Statement and discussion of main results. We now give concise statements of the main results in this paper. We refer the reader to [Section 2](#) for full statements of the notation and definitions used in this section.

Fix a random substitution $\vartheta_{\mathbf{P}}$ and let λ and \mathbf{R} denote the Perron–Frobenius eigenvalue and corresponding right eigenvector of the substitution matrix of $\vartheta_{\mathbf{P}}$, respectively. Given $q \in \mathbb{R}$ and $k \in \mathbb{N}$, define

$$\varphi_{\vartheta_{\mathbf{P}},k}(q) = \varphi_k(q) = - \sum_{a \in \mathcal{A}} R_a \log \left(\sum_{s \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = s]^q \right).$$

We define the *inflation word L^q -spectrum* of $\vartheta_{\mathbf{P}}$ by

$$T_{\vartheta,\mathbf{P}}(q) = \liminf_{k \rightarrow \infty} \frac{1}{\lambda^k} \varphi_k(q).$$

We similarly define the upper variant $\bar{T}_{\vartheta,\mathbf{P}}$ by taking a limit superior in place of the limit inferior. Throughout, $\mu_{\mathbf{P}}$ will denote the frequency measure associated with $\vartheta_{\mathbf{P}}$.

Heuristically, the inflation word spectrum approximates the frequency measure $\mu_{\mathbf{P}}$ by the probability distribution on the iterated system. The normalisation factors follow from the observation that the words in the k^{th} iteration have length approximately λ^k when normalised by the left Perron–Frobenius eigenvector.

Our main general result bounding the L^q -spectrum is the following, which states for $q \geq 0$ that $T_{\vartheta,\mathbf{P}}$ and $\tau_{\mu_{\mathbf{P}}}$ coincide, and moreover provides bounds on $\tau_{\mu_{\mathbf{P}}}$ in terms of the functions φ_k for all $q \in \mathbb{R}$.

Theorem A. *Let $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ be a primitive and compatible random substitution with corresponding frequency measure $\mu_{\mathbf{P}}$. Then the limits defining $\tau_{\mu_{\mathbf{P}}}(q)$ and $T_{\vartheta,\mathbf{P}}(q)$ exist and coincide for all $q \geq 0$. Moreover,*

(1) *For all $0 \leq q \leq 1$,*

$$(1.1) \quad \frac{1}{\lambda^k - 1} \varphi_k(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda^k} \varphi_k(q)$$

and $(\lambda^{-k} \varphi_k(q))_{k=1}^{\infty}$ converges monotonically to $\tau_{\mu_{\mathbf{P}}}(q)$ from above.

(2) *For all $q \geq 1$,*

$$(1.2) \quad \frac{1}{\lambda^k} \varphi_k(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda^k - 1} \varphi_k(q)$$

and $(\lambda^{-k} \varphi_k(q))_{k=1}^{\infty}$ converges monotonically to $\tau_{\mu_{\mathbf{P}}}(q)$ from below.

(3) *For all $q < 0$,*

$$(1.3) \quad \frac{1}{\lambda^k - 1} \varphi_k(q) \leq \tau_{\mu_{\mathbf{P}}}(q).$$

The notion of compatibility is defined in [Section 2.7](#), which is key in order to obtain the uniform estimate given in [Lemma 2.9](#). For $q < 0$, it is *not* true in general that $\tau_{\mu_{\mathbf{P}}}(q)$ and $T_{\vartheta,\mathbf{P}}(q)$ coincide (a counterexample is given in [Example 5.2](#)): the problem is essentially “non-uniqueness of cutting points”, as highlighted by the averaging procedure in the construction of the measure (see [Lemma 2.12](#)). In other words, the corresponding upper bound in (1.3) does not hold in general. Despite this, it still follows from (1.3) that $T_{\vartheta,\mathbf{P}}(q)$ provides a lower bound for $\tau_{\mu_{\mathbf{P}}}(q)$.

In **Proposition 3.1**, we prove that if ϑ_P also satisfies the *disjoint set condition*, or the *identical set condition with identical production probabilities* (see **Definition 2.13**), then a closed-form expression can be obtained for the L^q -spectrum of the corresponding frequency measure. By combining this result with **Theorem A**, we obtain the following corollary.

Corollary B. *Let ϑ_P be a primitive and compatible random substitution with corresponding frequency measure μ_P . Then for all $q \geq 0$:*

(1) *If ϑ_P satisfies the disjoint set condition, then*

$$\tau_{\mu_P}(q) = \frac{1}{\lambda - 1} \varphi_1(q).$$

(2) *If ϑ_P satisfies the identical set condition and has identical production probabilities, then*

$$\tau_{\mu_P}(q) = \frac{1}{\lambda} \varphi_1(q).$$

In particular, under the disjoint set condition or identical set condition with identical production probabilities, the L^q -spectrum is analytic on $(0, \infty)$.

In [15], a result analogous to **Theorem A** is obtained for entropy in terms of the *inflation word entropy*, without the compatibility assumption. In fact, to highlight the generality of our results on L^q -spectra, as a consequence of **Theorem A**, we obtain a new proof of this result (as well as a result on topological entropy).

- (a) We obtain the main result of [14] on topological entropy, which states that for subshifts of primitive and compatible random substitutions, the topological entropy can be characterised in terms of the asymptotic growth rate of inflation words.
- (b) For frequency measures corresponding to primitive and compatible random substitutions, we also obtain the characterisation of (measure theoretic) entropy obtained in [15, Theorem 3.3]—under the additional hypothesis that the substitution is compatible.

This is described in the following corollary.

Corollary C. *Let $\vartheta_P = (\vartheta, P)$ be a primitive and compatible random substitution with associated subshift X_ϑ and frequency measure μ_P .*

(1) *The limit*

$$\lim_{k \rightarrow \infty} \frac{1}{\lambda^k} \sum_{a \in \mathcal{A}} R_a \log(\#\vartheta^k(a))$$

exists and is equal to $h_{\text{top}}(X_\vartheta)$.

(2) *The L^q -spectrum of μ_P is differentiable at 1. Moreover, the limit*

$$\lim_{k \rightarrow \infty} \frac{1}{\lambda^k} \sum_{a \in \mathcal{A}} R_a \sum_{v \in \vartheta^k(a)} -\mathbb{P}[\vartheta_P^k(a) = v] \log(\mathbb{P}[\vartheta_P^k(a) = v])$$

exists and is equal to $h_{\mu_P}(X_\vartheta) = \dim_{\text{H}} \mu_P = \tau'_{\mu_P}(1)$.

We now turn our attention to the multifractal spectrum. Firstly, while $\tau_{\mu_P}(q)$ and $T_{\vartheta, P}(q)$ do not coincide in general for $q < 0$, if the random substitution that gives rise to the frequency measure μ_P is additionally assumed to be recognisable (see [Definition 2.14](#)), then the limits defining $\tau_{\mu_P}(q)$ and $T_{\vartheta, P}(q)$ both exist and coincide for all $q \in \mathbb{R}$. Moreover, under recognisability, we prove that the multifractal spectrum is the concave conjugate of the L^q -spectrum: in other words, the multifractal formalism holds for any associated frequency measure μ_P . In particular, we conclude that f_{μ_P} is a concave analytic function.

In fact, in [Proposition 4.5](#) we prove a stronger variational formula for the multifractal spectrum. For each $\alpha \in \mathbb{R}$, we construct measures ν such that $\dim_{\text{H}} \nu \geq \tau^*(\alpha)$ and $\dim_{\text{loc}}(\mu_P, x) = \alpha$ for ν -a.e. $x \in X_{\vartheta}$. In particular, we can take the measures to be frequency measures associated with permissible probabilities for the substitution ϑ .

Theorem D. *Let ϑ_P be a primitive, compatible, and recognisable random substitution with corresponding frequency measure μ_P . Then for all $q \in \mathbb{R}$,*

$$\tau_{\mu_P}(q) = T_{\vartheta, P}(q) = \frac{1}{\lambda - 1} \varphi_1(q).$$

Moreover, $f_{\mu_P}(\alpha) = \tau_{\mu_P}^*(\alpha)$ is an analytic and concave function. In fact, for each $\alpha \in \mathbb{R}$ such that $f_{\mu_P}(\alpha) \geq 0$, there are permissible probabilities \mathbf{Q} such that $f_{\mu_P}(\alpha) = \dim_{\text{H}} \mu_{\mathbf{Q}}$ and $\dim_{\text{loc}}(\mu_P, x) = \alpha$ for $\mu_{\mathbf{Q}}$ -a.e. $x \in X_{\vartheta}$.

To conclude this section, we observe that our results on L^q -spectra also give uniform bounds on the exponential scaling rate of the frequency measures. The following result is a direct application of [Theorem A](#) and [Theorem D](#), combined with [Proposition 4.4](#).

Corollary E. *Let $\vartheta_P = (\vartheta, \mathbf{P})$ be a primitive, compatible, and recognisable random substitution. Then*

$$\begin{aligned} \alpha_{\min} &:= \lim_{q \rightarrow \infty} \frac{\tau_{\mu_P}(q)}{q} = - \sum_{a \in \mathcal{A}} R_a \log \left(\max_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] \right) \\ \alpha_{\max} &:= \lim_{q \rightarrow -\infty} \frac{\tau_{\mu_P}(q)}{q} = - \sum_{a \in \mathcal{A}} R_a \log \left(\min_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] \right). \end{aligned}$$

and for all $x \in X_{\vartheta}$, $\alpha_{\min} \leq \underline{\dim}_{\text{loc}}(\mu_P, x) \leq \overline{\dim}_{\text{loc}}(\mu_P, x) \leq \alpha_{\max}$. Moreover,

$$\{\dim_{\text{loc}}(\mu_P, x) : x \in X_{\vartheta}\} = [\alpha_{\min}, \alpha_{\max}].$$

In particular, when the probabilities \mathbf{P} are chosen so that for each $a \in \mathcal{A}$, $\mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] = 1/\#\vartheta(a)$ for all $s \in \vartheta(a)$, then the L^q -spectrum is the line with slope $h_{\text{top}}(X_{\vartheta})$ passing through $(1, 0)$. Thus the local dimension of μ_P exists at every $x \in X_{\vartheta}$ and is given by the constant value $\alpha_{\min} = \alpha_{\max}$. This can be rephrased in terms of a weak Gibbs-type property, which says that for every $\epsilon > 0$, all n sufficiently large (depending on ϵ), and $u \in \mathcal{L}^n$,

$$(1.4) \quad \exp(-|u|(h_{\text{top}}(X_{\vartheta}) + \epsilon)) \leq \mu_P([u]) \leq \exp(-|u|(h_{\text{top}}(X_{\vartheta}) - \epsilon));$$

see, for example, [29, Lemma 1.4] for the short argument. In general, the error term ϵ cannot be dropped by the addition of a constant factor. Under certain assumptions, one can show that there are infinitely many words with $\mu_P([u]) \approx |u|^{-1} \exp(-|u|(h_{\text{top}}(X_\vartheta)))$, as explained in [15, Lemma 4.12]. These assumptions are satisfied, for example, in Example 5.6.

Of course, similar one-sided results hold for $q \geq 0$ only under the assumption of compatibility, by iterating the formula for φ_k and taking an appropriate maximum at each level. In fact, since $\tau_{\mu_P}(q)$ is differentiable at 1, with derivative giving the entropy, and since $h_{\text{top}}(X_\vartheta) = \tau_{\mu_P}(0)$, it follows that μ_P is a measure of maximal entropy if and only if $\tau'_{\mu_P}(q)$ exists and is constant on the interval $(0, 1)$.

1.4. Discussion and further work. We conclude the introduction with a list of comments and potentially interesting questions.

- (1) What is the L^q -spectrum for a compatible substitution when $q < 0$? We do not know this even for the random substitution given in Example 5.1, which satisfies the identical set condition with identical production probabilities. Obtaining results for $q < 0$ is substantially more challenging, since the sum in $\tau_{\mu_P}(q)$ depends on the measure of cylinders with very small (but non-zero) measure. For example, in the self-similar case, without the presence of strong separation assumptions, little is known (in contrast to the $q \geq 0$ case).
- (2) What happens without compatibility? Do the formulas in Theorem A hold in general? In [15], it suffices to use an almost sure version of Lemma 2.9. However, since the L^q -spectrum is sensitive to scaling at individual points as q tends to $\pm\infty$, such an almost sure result in our case is insufficient.
- (3) Outside the disjoint set condition and the identical set condition, what can be said about differentiability of the L^q -spectrum? For $q \geq 0$, we give the L^q -spectrum as a uniform limit of analytic functions: however, aside from the exceptional point $q = 1$ where we can say more, this is not enough to give information about differentiability.
- (4) Can our results on L^q -spectra and multifractal spectra (which hold for recognisable substitutions) be relaxed to a weaker condition such as the disjoint set condition (see Definition 2.13)?
- (5) Can the error term in (1.4) be determined precisely, up to a constant? The approximate Gibbs-type bounds discussed following Corollary E are closely related to the bounds used in the proof of intrinsic ergodicity given in [15, Theorem 4.8]. It could be worth exploring the relationship between intrinsic ergodicity and Gibbs-type properties given by the L^q -spectrum.

2. PRELIMINARIES

In this section we introduce the key notation and definitions that we will use throughout the paper. After introducing some basic notation, in Section 2.2 we introduce symbolic dynamics on the two-sided shift, as well as our notions of entropy

and dimension. In [Section 2.3](#) we present the key definitions and basic results from multifractal analysis that we work with throughout, including the definitions of the L^q -spectrum and local dimensions of a measure. Then, in the following sections we provide an introduction to random substitutions and their associated dynamical systems. In [Section 2.5](#) we give the definition of a random substitution via its action on words, and define the subshift associated to a random substitution. Then, in [Section 2.6](#) and [Section 2.7](#), we define what it means for a random substitution to be *primitive* and *compatible* and present the key properties of such random substitutions. In [Section 2.8](#), we give the definition of the frequency measure associated to a random substitution and state a key result used in the proof of our main results which relates the measures of cylinder sets via the action of the random substitution. Finally, in [Section 2.9](#), we define what it means for a substitution to satisfy the disjoint or identical set condition, and introduce recognisable random substitutions.

2.1. Symbolic notation. Throughout, we use the following symbolic notation, which is essentially the same as the notation used in [\[2, 20\]](#).

For a set B , we let $\#B$ be the cardinality of B and let $\mathcal{F}(B)$ be the set of non-empty finite subsets of B .

We fix an *alphabet* $\mathcal{A} = \{a_1, \dots, a_d\}$, for some $d \in \mathbb{N}$, which is a finite set of *letters* a_i , and equip it with the discrete topology. Then a *word* u with letters in \mathcal{A} is a finite concatenation of letters, namely $u = a_{i_1} \cdots a_{i_n}$ for some $n \in \mathbb{N}$. We write $|u| = n$ for the length of the word u , and for $m \in \mathbb{N}$, we let \mathcal{A}^m denote the set of all words of length m with letters in \mathcal{A} .

We set $\mathcal{A}^+ = \bigcup_{m \in \mathbb{N}} \mathcal{A}^m$ and let

$$\mathcal{A}^{\mathbb{Z}} = \{(a_{i_n})_{n \in \mathbb{Z}} : a_{i_n} \in \mathcal{A} \text{ for all } n \in \mathbb{Z}\}$$

denote the set of all bi-infinite sequences with elements in \mathcal{A} , and endow $\mathcal{A}^{\mathbb{Z}}$ with the product topology. We also fix a metric on $\mathcal{A}^{\mathbb{Z}}$ as follows. Given points $x = (x_n)_{n \in \mathbb{Z}}$ and $y = (y_n)_{n \in \mathbb{Z}}$, let $N(x, y) = \sup\{n \in \mathbb{Z} : x_j = y_j \text{ for all } |j| \leq n\}$ and let $d(x, y) = e^{-2N(x, y)-1}$. The space X is compact with topology generated by the metric.

We will frequently write sequences $(x_n)_{n \in \mathbb{Z}} \in \mathcal{A}^{\mathbb{Z}}$ as $\cdots x_{-1}x_0x_1\cdots$, with the corresponding notation for finite sequences.

If i and $j \in \mathbb{Z}$ with $i \leq j$, and $x = \cdots x_{-1}x_0x_1\cdots \in \mathcal{A}^{\mathbb{Z}}$, then we let $x_{[i, j]} = x_i x_{i+1} \cdots x_j$. We use the same notation if $v \in \mathcal{A}^+$ and $1 \leq i \leq j \leq |v|$. For u and $v \in \mathcal{A}^+$ (or $v \in \mathcal{A}^{\mathbb{Z}}$), we write $u \triangleleft v$ if u is a subword of v , namely if there exist i and $j \in \mathbb{Z}$ with $i \leq j$ so that $u = v_{[i, j]}$. For u and $v \in \mathcal{A}^+$, we set $|v|_u$ to be the number of (possibly overlapping) occurrences of u as a subword of v . If $u = a_{i_1} \cdots a_{i_n}$ and $v = a_{j_1} \cdots a_{j_m} \in \mathcal{A}^+$, for some n and $m \in \mathbb{N}$, we write uv for the *concatenation* of u and v . The *abelianisation* of a word $u \in \mathcal{A}^+$ is the vector $\Phi(u) \in \mathbb{N}_0^{\#\mathcal{A}}$, defined by $\Phi(u)_a = |u|_a$ for all $a \in \mathcal{A}$.

2.2. Dynamics, entropy and dimension. We equip the space $\mathcal{A}^{\mathbb{Z}}$ with invertible dynamics from the *left-shift map* $S: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$. Throughout, we will work with a

subshift $X \subset \mathcal{A}^{\mathbb{Z}}$, which is compact and *shift-invariant*, that is $S^{-1}(X) = X$. Then μ will denote an ergodic and S -invariant Borel probability measure with support contained in X .

The metric structure on $\mathcal{A}^{\mathbb{Z}}$ enables us to define the *Hausdorff dimension* of Borel subsets of X . Using this, we define the *Hausdorff dimension of μ* to be the quantity

$$\dim_{\text{H}} \mu = \inf\{\dim_{\text{H}} E : \mu(E) > 0\}$$

where the infimum is taken over Borel sets E . Here, we use Hausdorff dimension as inherited from the underlying metric; though it would also be appropriate to use Bowen's generalisation of topological entropy to non-compact sets [4]. We also define the *lower local dimension of μ at x* by

$$\underline{\dim}_{\text{loc}}(\mu, x) = \liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r}$$

We define the *upper local dimension* $\overline{\dim}_{\text{loc}}(\mu, x)$ analogously using the limit superior in place of the limit inferior, and when the limits coincide, we refer to the shared quantity as the *local dimension* and denote it by $\dim_{\text{loc}}(\mu, x)$.

Local dimensions and Hausdorff dimension are closely related: it follows from, for instance, [7, Proposition 10.1] that

$$(2.1) \quad \dim_{\text{H}} \mu = \sup\{s : \underline{\dim}_{\text{loc}}(\mu, x) \geq s \text{ for } \mu\text{-a.e. } x\}.$$

Now fix a partition ξ so that with $\xi_k = \bigvee_{i=-k}^k S^{-i}(\xi)$, $\{\xi_k\}_{k=1}^{\infty}$ generates the Borel σ -algebra on X . We recall that the *entropy* of μ with respect to S is given by

$$h_{\mu}(X) = \lim_{k \rightarrow \infty} \frac{1}{2k+1} \sum_{A \in \xi_k} -\mu(A) \log(\mu(A)).$$

where, by the classical Kolmogorov–Sinaï theorem, the quantity does not depend on the choice of partition.

Now given $x \in X$, let $\xi_k(x)$ denote the unique element in the partition ξ_k containing x . Then the Shannon–McMillan–Breiman theorem states that the entropy of μ is almost surely the *information rate* of μ , that is for μ -a.e. $x \in X$,

$$\lim_{k \rightarrow \infty} \frac{-\log \mu(\xi_k(x))}{2k+1} = h_{\mu}(X).$$

We refer the reader to [6] for greater detail.

Now suppose $\xi = \{E_a\}_{a \in \mathcal{A}}$ is the partition of X where $E_a = \{(x_n)_{n \in \mathbb{Z}} \in X : x_0 = a\}$. For the remainder of this paper, ξ will always denote this partition. Then given $x = (x_n)_{n \in \mathbb{Z}} \in X$,

$$\xi_k(x) = \{y \in X : x_j = y_j \text{ for all } |j| \leq k\} = B(x, e^{-(2k+1)})$$

and therefore

$$\dim_{\text{loc}}(\mu, x) = \lim_{k \rightarrow \infty} \frac{-\log \mu(\xi_k(x))}{2k+1}$$

where both limits exist if either limit exists. Since the limit on the right is μ almost surely $h_\mu(X)$, it follows from (2.1) that $\dim_{\text{H}} \mu = h_\mu(X)$.

Finally, the *topological entropy* of X is given by

$$h_{\text{top}}(X) = \lim_{k \rightarrow \infty} \frac{-\log \#\{E \in \xi_k : E \cap X \neq \emptyset\}}{2k+1}.$$

Of course, $h_{\text{top}}(X) = \dim_{\text{B}} X$, the box counting dimension of X .

2.3. L^q -spectra and smoothness. Given $q \in \mathbb{R}$, we define

$$\bar{S}_{\mu,r}(q) = \sup_{\{x_i\}_i \in \mathcal{P}(r)} \sum_i \mu(B(x_i, r))^q$$

where $\mathcal{P}(r)$ is the set of discrete $2r$ -separated subsets of X , that is $\mathcal{P}(r) = \{\{x_i\}_i : x_i \in X, d(x_i, x_j) \geq 2r \text{ for } i \neq j\}$. We then define the L^q -spectrum of μ to be the function

$$\tau_\mu(q) = \liminf_{q \rightarrow 0} \frac{\log \bar{S}_{\mu,r}(q)}{\log r}.$$

For convenience, we also denote the upper variant $\bar{\tau}_\mu(q)$ by taking a limit superior in place of the limit inferior. It is a standard consequence of Hölder's inequality that $\tau_\mu(q)$ is a concave increasing function of q (note that this need not hold for $\bar{\tau}_\mu(q)$).

Of course, the preceding definitions hold more generally in an arbitrary metric space, but in our particular setting we can rephrase the L^q -spectrum in terms of more familiar sums over cylinders. Recall that ξ denotes the partition of X_ϑ into cylinders at 0 corresponding to the letters in \mathcal{A} . Then set

$$S_{\mu,n}(q) = \sum_{E \in \xi_k} \mu(E)^q.$$

Since distinct elements in the partition ξ_k are $e^{-(2k+1)}$ -separated,

$$S_{\mu,n}(q) = \bar{S}_{\mu, e^{-(2n+1)}}(q).$$

It follows immediately that

$$\tau_\mu(q) = \liminf_{n \rightarrow \infty} \frac{-\log S_{\mu,n}(q)}{2n+1}$$

with the analogous result for $\bar{\tau}_\mu(q)$. In particular, we observe that $\tau_\mu(0) = h_{\text{top}}(X)$ assuming μ is fully supported on X .

Finally, by shift invariance, we can characterise the subshift X in terms of a *language* on X . Given $n \in \mathbb{N}$, we set

$$\mathcal{L}^n(X) = \{w \in \mathcal{A}^n : w \triangleleft x \text{ for some } x \in X\}.$$

Given $w \in \mathcal{L}^n(X)$, we let $[w] = \{(x_n)_{n \in \mathbb{Z}} \in X : x_i = w_i \text{ for all } 1 \leq i \leq n\}$. Of course, by shift invariance, there is a measure-preserving bijection between $\mathcal{L}^{2n+1}(X)$ and X_n ,

so it follows again that

$$\tau_\mu(q) = \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sum_{u \in \mathcal{L}^n(X)} \mu([u])^q.$$

We will primarily use this characterisation throughout the paper.

We first list some basic properties of the L^q -spectrum of the measure μ . Here, (a) is a direct consequence of Hölder's inequality, (b) is standard (see, for example, [29, Lemma 1.4]) and (c) was proven in [9, Theorem 1.4].

Lemma 2.1. *Let μ be a shift-invariant measure on X .*

- (a) *The L^q -spectrum $\tau_\mu(q)$ is continuous, increasing and concave on \mathbb{R} .*
- (b) *Let $\alpha_{\min} = \lim_{q \rightarrow \infty} \tau_\mu(q)/q$ and $\alpha_{\max} = \lim_{q \rightarrow -\infty} \tau_\mu(q)/q$. Then for every $s < \alpha_{\min} \leq \alpha_{\max} < t$, all n sufficiently large and $u \in \mathcal{L}^n$, $e^{-tn} \leq \mu([u]) \leq e^{-sn}$. In particular, the local dimensions satisfy*

$$\alpha_{\min} \leq \inf_{x \in X} \underline{\dim}_{\text{loc}}(\mu, x) \leq \sup_{x \in X} \overline{\dim}_{\text{loc}}(\mu, x) \leq \alpha_{\max}.$$

- (c) *The left and right derivatives of τ_μ at $q = 1$ bound the Hausdorff dimension of μ , that is $\tau_\mu^+(1) \leq \dim_{\text{H}} \mu \leq \tau_\mu^-(1)$.*

In fact, (a) gives intuition for why the L^q -spectrum encodes smoothness: rather than obtain almost sure information on local dimensions, the L^q -spectrum contains *uniform* information about local dimensions.

Finally, we prove a simple result concerning the L^q -spectrum which will be useful later in the paper.

Lemma 2.2. *Let $\zeta > 1$ be arbitrary. Then*

$$(2.2) \quad \tau_\mu(q) = \frac{1}{\zeta} \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}^{\lfloor \zeta n \rfloor}(X)} \mu([u])^q \right)$$

and

$$(2.3) \quad \bar{\tau}_\mu(q) = \frac{1}{\zeta} \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}^{\lfloor \zeta n \rfloor}(X)} \mu([u])^q \right).$$

Proof. Of course, it always holds that

$$\begin{aligned} \tau_\mu(q) &\leq \frac{1}{\zeta} \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}^{\lfloor \zeta n \rfloor}(X)} \mu([u])^q \right) \\ \bar{\tau}_\mu(q) &\geq \frac{1}{\zeta} \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}^{\lfloor \zeta n \rfloor}(X)} \mu([u])^q \right). \end{aligned}$$

First, let $q < 0$ and let $n \in \mathbb{N}$ be arbitrary. Let k_n be minimal so that $\lfloor \zeta k_n \rfloor \geq n$. Observe that there is some $M \in \mathbb{N}$ (independent of n) so that $\lfloor \zeta k_n \rfloor \leq n + M$: it follows that $\lim_{n \rightarrow \infty} n/k_n = \zeta$. Then if $v \in \mathcal{L}^{\lfloor \zeta k_n \rfloor}(X)$ is arbitrary, $[v] \subset [u]$ for some $u \in \mathcal{L}^n(X)$ and $\mu([v])^q \geq \mu([u])^q$. Thus

$$S_{\lfloor \zeta k_n \rfloor, \mu}(q) \geq S_{n, \mu}(q).$$

which gives (2.2) for $q < 0$ since $\lim n/k_n = \zeta$.

Similarly, for $q \geq 0$, since there are at most $(\#\mathcal{A})^M$ words $v \in \mathcal{L}^{\lfloor \zeta k_n \rfloor}(X)$ with $[v] \subset [u]$, pigeonholing, for each $u \in \mathcal{L}^n(X)$ there is some $v(u) \in \mathcal{L}^{\lfloor \zeta k_n \rfloor}(X)$ such that $\mu([v(u)])^q \geq (\#\mathcal{A})^{-qM} \mu([u])^q$. Thus

$$S_{\lfloor \zeta k_n \rfloor, \mu}(q) \geq (\#\mathcal{A})^{-qM} S_{n, \mu}(q).$$

This gives (2.2) for $q \geq 0$.

The arguments for (2.3) follow analogously by choosing k_n maximal so that $\lfloor \zeta k_n \rfloor \leq n$. \square

2.4. Multifractal spectrum and multifractal formalism. The L^q -spectrum of a measure is related to the (*fine*) *multifractal spectrum*. Let μ be a shift-invariant measure on a subshift X . We recall that the local dimension of μ at $x \in X$ is given by

$$\dim_{\text{loc}}(\mu, x) = \lim_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mu([x_{[-n, n]}])$$

when the limit exists. Given $\alpha \in \mathbb{R}$, set

$$F_\mu(\alpha) = \{x \in X : \dim_{\text{loc}}(\mu, x) = \alpha\}.$$

We then define the *multifractal spectrum* of μ by

$$f_\mu(\alpha) = \dim_{\text{H}} F_\mu(\alpha)$$

using the convention that $\dim_{\text{H}} \emptyset = -\infty$.

The multifractal spectrum is related to the L^q -spectrum by the following result. Let $g: \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a concave function. For $x \in \mathbb{R}$, we let $g^+(x)$ (resp. $g^-(x)$) denote the right (resp. left) derivative of g at x . Such limits necessarily exist by concavity. We denote the *subdifferential* of g at x by $\partial g(x) = [g^+(x), g^-(x)]$. We then recall that the *concave conjugate* of g is given by

$$g^*(\alpha) = \inf_{q \in \mathbb{R}} \{q\alpha - g(q)\}.$$

Note that g^* is always concave since it is the infimum of a family of affine functions. For more detail concerning the theory of concave functions, we refer the reader to [26].

Now, we say that μ satisfies the *multifractal formalism* when $f_\mu = \tau_\mu^*$. In general, the multifractal formalism need not hold, but it is well-known that the concave conjugate of the L^q -spectrum is an upper bound for the multifractal spectrum. For the convenience of the reader, we provide a short self-contained proof, which follows the main ideas of [19, Theorem 4.1].

Proposition 2.3. *Let μ be a shift-invariant measure on a subshift X . Then $f_\mu(\alpha) \leq \tau^*(\alpha)$ for all $\alpha \in \mathbb{R}$.*

Proof. Recall that ξ_n denotes the partition of X into cylinders corresponding to words of length $2n + 1$, each of which has diameter precisely e^{-2n-1} . For $\alpha \in \mathbb{R}$, $n \in \mathbb{N}$ and $\epsilon > 0$, let

$$\mathcal{M}_{n,\epsilon}(\alpha) = \left\{ I \in \xi_n : e^{-(2n+1)(\alpha+\epsilon)} \leq \mu(I) \leq e^{-(2n+1)(\alpha-\epsilon)} \right\}.$$

In other words, $\mathcal{M}_{n,\epsilon}(\alpha)$ is an ϵ -approximation of $F_\mu(\alpha)$ at level n . Our strategy is to control the size of the sets $\mathcal{M}_{n,\epsilon}(\alpha)$ in terms of the L^q -spectrum of μ , and then use these sets to build a good cover of $F_\mu(\alpha)$. Let $q \in \partial\tau^*(\alpha)$: we prove this in the case that $q \geq 0$; the case $q < 0$ is analogous.

First, observe that

$$(2.4) \quad S_{2n+1,\mu}(q) = \sum_{I \in \xi_n} \mu(I)^q \geq \sum_{u \in \mathcal{M}_{n,\epsilon}(\alpha)} \mu(I)^q \geq e^{-(2n+1)(\alpha+\epsilon)q} \# \mathcal{M}_{n,\epsilon}(\alpha).$$

Since $\tau_\mu(q) = \liminf_{n \rightarrow \infty} (\log S_{2n+1,\mu}(q)) / (-2n-1)$ by [Lemma 2.2](#), there is some $N_\epsilon \in \mathbb{N}$ so that for all $n \geq N_\epsilon$, $S_{2n+1,\mu}(q) \leq e^{-(2n+1)(\tau_\mu(q)-\epsilon)}$. Combining this with (2.4),

$$(2.5) \quad \# \mathcal{M}_{n,\epsilon}(\alpha) \leq e^{-(2n+1)(\tau(q)-\epsilon)} \cdot e^{(2n+1)(\alpha+\epsilon)q} = e^{(2n+1)(\tau^*(\alpha)+(q+1)\epsilon)}$$

for all $n \geq N_\epsilon$ where we have used the fact that $q \in \partial\tau^*(\alpha)$.

Now for each $x \in F_\mu(\alpha)$, we can find some $n_x \in \mathbb{N}$ so that for all $n \geq n_x$, $\mu(\xi_n(x)) \geq e^{-(2n+1)(\alpha+\epsilon)}$. In particular,

$$\mathcal{G}_\epsilon := \bigcup_{n=N_\epsilon}^{\infty} \mathcal{M}_{n,\epsilon}(\alpha)$$

is a Vitali cover for $F_\mu(\alpha)$.

Now suppose $\{I_j\}_{j=1}^{\infty}$ is any disjoint subcollection of \mathcal{G}_ϵ : then with $s = \tau^*(\alpha) + 2\epsilon(1+q)$,

$$\begin{aligned} \sum_{j=1}^{\infty} (\text{diam } I_j)^s &\leq \sum_{n=N_\epsilon}^{\infty} \sum_{I \in \mathcal{M}_{n,\epsilon}(\alpha)} (\text{diam } I)^s \leq \sum_{n=N_\epsilon}^{\infty} e^{-(2n+1)s} \# \mathcal{M}_{n,\epsilon}(\alpha) \\ &\leq \sum_{n=N_\epsilon}^{\infty} e^{-(2n+1)s} e^{(2n+1)(\tau^*(\alpha)+(q+1)\epsilon)} \\ &= \sum_{n=N_\epsilon}^{\infty} (e^{-(1+q)\epsilon})^{2n+1} < \infty \end{aligned}$$

by (2.5). Thus by the Vitali covering theorem for Hausdorff measure, there is a cover $\{E_i\}_{i=1}^{\infty}$ for $F_\mu(\alpha)$ such that

$$\mathcal{H}^s(F_\mu(\alpha)) \leq \sum_{i=1}^{\infty} (\text{diam } E_i)^s < \infty$$

and thus $\dim_{\mathbb{H}} F_{\mu}(\alpha) \leq \tau^*(\alpha) + 2\epsilon(1+q)$. But $\epsilon > 0$ was arbitrary, so the desired result follows. \square

2.5. Random substitutions and frequency measures. We now introduce our primary objects of interest: *random substitutions*, and their associated *frequency measures*. In a similar manner to [14, 15], we define a random substitution by the data required to determine its action on letters. We then extend this to a random map on words.

Definition 2.4. Let $\mathcal{A} = \{a_1, \dots, a_d\}$ be a finite alphabet. A *random substitution* $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ is a *set-valued substitution* $\vartheta: \mathcal{A} \rightarrow \mathcal{F}(\mathcal{A}^+)$ together with a set of non-degenerate probability vectors

$$\mathbf{P} = \left\{ \mathbf{p}_i = (p_{i,1}, \dots, p_{i,r_i}) : r_i = \#\vartheta(a_i); \mathbf{p}_i \in (0, 1]^{r_i}; \sum_{j=1}^{r_i} p_{i,j} = 1 \text{ for all } i = 1, \dots, d \right\},$$

such that

$$\vartheta_{\mathbf{P}}: a_i \mapsto \begin{cases} s^{(i,1)} & \text{with probability } p_{i,1}, \\ \vdots & \vdots \\ s^{(i,r_i)} & \text{with probability } p_{i,r_i}, \end{cases}$$

for every $1 \leq i \leq d$, where $\vartheta(a_i) = \{s^{(i,j)}\}_{1 \leq j \leq r_i}$.

We call each $s^{(i,j)}$ a *realisation* of $\vartheta_{\mathbf{P}}(a_i)$. If $r_i = 1$ for all $i \in \{1, \dots, d\}$, then we call $\vartheta_{\mathbf{P}}$ *deterministic*.

Example 2.5 (Random Fibonacci). Let $\mathcal{A} = \{a, b\}$, and let $p \in (0, 1)$. The *random Fibonacci substitution* $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ is the random substitution given by

$$\vartheta_{\mathbf{P}}: \begin{cases} a \mapsto \begin{cases} ab & \text{with probability } p, \\ ba & \text{with probability } 1-p, \end{cases} \\ b \mapsto a \end{cases}$$

with defining data $r_a = 2, r_b = 1, s^{(a,1)} = ab, s^{(a,2)} = ba, s^{(b,1)} = a, \mathbf{P} = \{\mathbf{p}_a = (p, 1-p), \mathbf{p}_b = (1)\}$ and corresponding set-valued substitution $\vartheta: a \mapsto \{ab, ba\}, b \mapsto \{a\}$.

In the following we describe how a random substitution $\vartheta_{\mathbf{P}}$ determines a (countable state) Markov matrix Q , indexed by $\mathcal{A}^+ \times \mathcal{A}^+$. We interpret the entry $Q_{u,v}$ as the probability of mapping a word u to a word v under the random substitution. Formally, $Q_{a_i, s^{(i,j)}} = p_{i,j}$ for all $j \in \{1, \dots, r_i\}$ and $Q_{a_i, v} = 0$ if $v \notin \vartheta(a_i)$. We extend the action of $\vartheta_{\mathbf{P}}$ to finite words by mapping each letter *independently* to one of its realisations. More precisely, given $n \in \mathbb{N}$, $u = a_{i_1} \dots a_{i_n} \in \mathcal{A}^n$ and $v \in \mathcal{A}^+$ with $|v| \geq n$, we let

$$\mathcal{D}_n(v) = \{(v^{(1)}, \dots, v^{(n)}) \in (\mathcal{A}^+)^n : v^{(1)} \dots v^{(n)} = v\}$$

denote the set of all decompositions of v into n individual words and set

$$Q_{u,v} = \sum_{(v^{(1)}, \dots, v^{(n)}) \in \mathcal{D}_n(v)} \prod_{j=1}^n Q_{a_{i_j}, v^{(j)}}.$$

In words, $\vartheta_P(u) = v$ with probability $Q_{u,v}$.

For $u \in \mathcal{A}^+$, let $(\vartheta_P^n(u))_{n \in \mathbb{N}}$ be a stationary Markov chain on some probability space $(\Omega_u, \mathcal{F}_u, \mathbb{P}_u)$, with Markov matrix given by Q ; that is,

$$\mathbb{P}_u[\vartheta_P^{n+1}(u) = w \mid \vartheta_P^n(u) = v] = \mathbb{P}_v[\vartheta_P(v) = w] = Q_{v,w}$$

for all v and $w \in \mathcal{A}^+$, and $n \in \mathbb{N}$. In particular,

$$\mathbb{P}_u[\vartheta_P^n(u) = v] = (Q^n)_{u,v}$$

for all u and $v \in \mathcal{A}^+$, and $n \in \mathbb{N}$. We often write \mathbb{P} for \mathbb{P}_u if the initial word is understood. In this case, we also write \mathbb{E} for the expectation with respect to \mathbb{P} . As before, we call v a *realisation* of $\vartheta_P^n(u)$ if $(Q^n)_{u,v} > 0$ and set

$$\vartheta^n(u) = \{v \in \mathcal{A}^+ : (Q^n)_{u,v} > 0\}$$

to be the set of all realisations of $\vartheta_P^n(u)$. Conversely, we may regard $\vartheta_P^n(u)$ as the set $\vartheta^n(u)$ endowed with the additional structure of a probability vector. If $u = a \in \mathcal{A}$ is a letter, we call a word $v \in \vartheta^k(a)$ a *level- k inflation word*, or *exact inflation word*.

To a given random substitution $\vartheta_P = (\vartheta, P)$ one can associate a subshift. First, we say that a word $u \in \mathcal{A}^+$ is *(ϑ -)legal* if there exists an $a_i \in \mathcal{A}$ and $k \in \mathbb{N}$ such that u appears as a subword of some word in $\vartheta^k(a_i)$. We define the *language* of ϑ by $\mathcal{L}_\vartheta = \{u \in \mathcal{A}^+ : u \text{ is } \vartheta\text{-legal}\}$ and, for $w \in \mathcal{A}^+ \cup \mathcal{A}^\mathbb{Z}$, we let $\mathcal{L}(w) = \{u \in \mathcal{A}^+ : u \triangleleft w\}$ denote the language of w .

Definition 2.6. The *random substitution subshift* of a random substitution $\vartheta_P = (\vartheta, P)$ is the system (X_ϑ, S) , where $X_\vartheta = \{w \in \mathcal{A}^\mathbb{Z} : \mathcal{L}(w) \subseteq \mathcal{L}_\vartheta\}$ and S denotes the (left) shift map, defined by $S(w)_i = w_{i+1}$ for each $w \in X_\vartheta$.

Under very mild assumptions, the space X_ϑ is non-empty [17]. This holds, for example, if the generating random substitution is primitive (we give a definition in Section 2.6). We endow X_ϑ with the subspace topology inherited from $\mathcal{A}^\mathbb{Z}$, and since X_ϑ is defined in terms of a language, it is a compact S -invariant subspace of $\mathcal{A}^\mathbb{Z}$. Hence, X_ϑ is a subshift. For $n \in \mathbb{N}$, we write $\mathcal{L}_\vartheta^n = \mathcal{L}_\vartheta \cap \mathcal{A}^n$ and $\mathcal{L}^n(w) = \mathcal{L}(w) \cap \mathcal{A}^n$ to denote the subsets of \mathcal{L}_ϑ and $\mathcal{L}(w)$, respectively, consisting of words of length n . The set-valued function ϑ naturally extends to X_ϑ , where for $w = \dots w_{-1}w_0w_1\dots \in X_\vartheta$ we let $\vartheta(w)$ denotes the (infinite) set of sequences of the form $v = \dots v_{-2}v_{-1}.v_0v_1\dots$, with $v_j \in \vartheta(w_j)$ for all $j \in \mathbb{Z}$. It is easily verified that $\vartheta(X_\vartheta) \subset X_\vartheta$.

The notation X_ϑ reflects the fact that the random substitution subshift does not depend on the choice of (non-degenerate) probabilities P . In fact, this is the case for many structural properties of ϑ_P . In these cases, one sometimes refers to ϑ instead of ϑ_P as a random substitution, see for instance [14, 16, 27, 28]. On the other hand, for

some applications, one needs additional structure on the probability space. In fact, there is an underlying branching process, similar to a Galton–Watson process, that allows one to construct more refined random variables, see [17] for further details. The measure theoretic properties we consider are typically dependent on the choice of probabilities; however, some of the auxiliary results we use only depend on the set-valued substitution ϑ . To avoid confusion, for results where there is no dependence on the choice of probabilities we will give the statement in terms of the set-valued substitution ϑ and omit the dependence on \mathbf{P} in the notation.

2.6. Primitive random substitutions. A standard assumption in the study of substitutions (both deterministic and random) is that of *primitivity*. Given a random substitution $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ over an alphabet $\mathcal{A} = \{a_1, \dots, a_d\}$ with cardinality $d \in \mathbb{N}$, we define the *substitution matrix* $M = M_{\vartheta_{\mathbf{P}}} \in \mathbb{R}^{d \times d}$ of $\vartheta_{\mathbf{P}}$ by

$$M_{i,j} = \mathbb{E}[|\vartheta_{\mathbf{P}}(a_j)|_{a_i}] = \sum_{k=1}^{r_j} p_{j,k} |s^{(j,k)}|_{a_i}.$$

Since M has only non-negative entries, it has a real eigenvalue λ of maximal modulus. Observe that $\lambda \geq 1$, with $\lambda = 1$ precisely if M is column-stochastic, so that the random substitution is non-expanding. To avoid this degenerate situation, we will assume that $\lambda > 1$. If the matrix M is *primitive* (that is if there exists a $k \in \mathbb{N}$ such that all the entries of M^k are positive), the Perron–Frobenius theorem gives that λ is a simple eigenvalue and that the corresponding (right) eigenvector $\mathbf{R} = (R_1, \dots, R_d)$ can be chosen to have strictly positive entries. We will normalise this eigenvector so that $\|\mathbf{R}\|_1 = 1$. We will refer to λ as the *Perron–Frobenius eigenvalue* of the random substitution, $\vartheta_{\mathbf{P}}$, with corresponding *Perron–Frobenius eigenvector* \mathbf{R} .

Definition 2.7. We say that $\vartheta_{\mathbf{P}}$ is *primitive* if $M = M_{\vartheta_{\mathbf{P}}}$ is primitive and its Perron–Frobenius eigenvalue satisfies $\lambda > 1$.

We emphasise that for a random substitution $\vartheta_{\mathbf{P}}$, being primitive is independent of the (non-degenerate) choice of probabilities \mathbf{P} . In this sense, primitivity is a property of ϑ rather than $\vartheta_{\mathbf{P}}$.

Since $M_{\vartheta_{\mathbf{P}}}^k = M_{\vartheta_{\mathbf{P}}^k}$, the Perron–Frobenius eigenvalue of $\vartheta_{\mathbf{P}}^k$ is λ^k .

2.7. Compatible random substitutions. Another standard assumption in the study of random substitutions is that of *compatibility*, which gives that exact inflation words have a well-defined abelianisation. In particular, the matrix of a compatible random substitution is independent of the choice of probabilities, so the letter frequencies are uniform and do not depend on the realisation. As discussed in the introduction, the existence of uniform letter frequencies is fundamental in the proofs of our main results.

Definition 2.8. We say that a random substitution $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ is *compatible* if for all $a \in \mathcal{A}$, and $u, v \in \vartheta(a)$, we have $\Phi(u) = \Phi(v)$.

Observe that compatibility is independent of the choice of probabilities, and that a random substitution $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ is compatible if and only if for all $u \in \mathcal{A}^+$, we have that $|s|_a = |t|_a$ for all s and $t \in \vartheta(u)$, and $a \in \mathcal{A}$. We write $|\vartheta(u)|_a$ to denote this common value, and let $|\vartheta(u)|$ denote the common length of words in $\vartheta(u)$. For convenience, we also set $|\vartheta| = \max_{a \in \mathcal{A}} |\vartheta(a)|$. For a random substitution that is both primitive and compatible, the (uniform) letter frequencies are encoded by the right Perron–Frobenius eigenvector of the substitution matrix, which by compatibility is independent of the choice of probabilities. In particular, we have the following (see [25] for a proof in the deterministic case, which also holds in the random case by compatibility).

Lemma 2.9 (Letter frequency bounds). *If $\vartheta_{\mathbf{P}}$ is a primitive and compatible random substitution, then for all $\varepsilon > 0$ there is an integer N such that every word v of length at least N satisfies*

$$|v|(R_a - \varepsilon) < |v|_a < |v|(R_a + \varepsilon)$$

for all $a \in \mathcal{A}$.

The random Fibonacci substitution defined in Example 2.5 is compatible, since $\Phi(ab) = \Phi(ba) = (1, 1)$. It is also primitive, since the square of its substitution matrix is positive. For any choice of probabilities, the right Perron–Frobenius eigenvector is given by (τ^{-1}, τ^{-2}) , where τ denotes the golden ratio. In terms of letter frequencies, this means that in all sufficiently long legal words, approximately τ^{-1} proportion of the letters are a and τ^{-2} proportion are b .

The following consequence of Lemma 2.9 is useful in the proof of Theorem A.

Lemma 2.10. *Let $\vartheta_{\mathbf{P}} = (\vartheta, \mathbf{P})$ be a primitive and compatible random substitution and let $q \geq 1$. For all $\varepsilon > 0$, there is an $M \in \mathbb{N}$ such that for every $m \geq M$ and $v \in \mathcal{L}_{\vartheta}^m$,*

$$\begin{aligned} \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{m(R_a + \varepsilon)} &\leq \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q \\ &\leq \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{m(R_a - \varepsilon)}. \end{aligned}$$

For $q \leq 1$, the same result holds with reversed inequalities.

Proof. Since $\vartheta_{\mathbf{P}}$ is compatible, the cutting points of inflation tiles are well-defined, so breaking the sum into inflation tiles we obtain

$$\begin{aligned} \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q &= \sum_{w^1 \in \vartheta(v_1)} \mathbb{P}[\vartheta_{\mathbf{P}}(v_1) = w^1]^q \sum_{w^2 \in \vartheta(v_2)} \cdots \sum_{w^m \in \vartheta(v_m)} \mathbb{P}[\vartheta_{\mathbf{P}}(v_m) = w^m]^q \\ &= \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{|v|_a}. \end{aligned}$$

The result then follows by applying [Lemma 2.9](#) to bound $|v|_a$, noting that for all $a \in \mathcal{A}$ we have $\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \leq 1$ if $q \geq 1$ and $\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \geq 1$ if $q \leq 1$. \square

2.8. Frequency measures. The main object that we associate with a given primitive random substitution ϑ_P is the *frequency measure* μ_P . This measure quantifies the relative occurrence of a given word in a random substitution. We now define this measure precisely.

First, we define the *expected frequency* of a word $v \in \mathcal{L}_\vartheta$ by

$$\text{freq}(v) = \lim_{k \rightarrow \infty} \frac{\mathbb{E}[|\vartheta_P^k(a)|_v]}{\mathbb{E}[|\vartheta_P^k(a)|]},$$

where, by primitivity, this limit is independent of the choice of $a \in \mathcal{A}$. In fact, we have the stronger property that the word frequencies exist \mathbb{P} -almost surely in the limit of large inflation words and are given by $\text{freq}(v)$ for all $v \in \mathcal{L}_\vartheta$ (see [\[17\]](#) for further details). Recalling that $\xi(X_\vartheta)$ is the algebra of cylinder sets on X_ϑ that specify the origin, we define $\mu_P: \xi(X_\vartheta) \cup \{\emptyset\} \rightarrow [0, 1]$ by $\mu_P(\emptyset) = 0$, $\mu_P(X_\vartheta) = 1$, and $\mu_P([v]_m) = \text{freq}(v)$ for $v \in \mathcal{L}_\vartheta$ and $m \in \{1 - |v|, 2 - |v|, \dots, 0\}$. This set function extends to a unique measure.

Proposition 2.11 ([\[17, Proposition 5.3 and Theorem 5.9\]](#)). *The set function μ_P is a content with mass one which extends uniquely to a shift-invariant ergodic Borel probability measure on X_ϑ .*

We call the measure μ_P defined in [Proposition 2.11](#) the *frequency measure* corresponding to the random substitution ϑ_P . Observe that frequency measures are dependent on the probabilities of the substitution. As such, for the subshift of a primitive random substitution that is non-deterministic, there exist uncountably many frequency measures supported on this subshift [\[17\]](#). In contrast, the subshift of a primitive deterministic substitution has precisely one frequency measure, which is the unique ergodic measure [\[25\]](#).

Frequency measures corresponding to primitive and compatible random substitutions satisfy the following renormalisation lemma, which relates the measure of a cylinder set of a legal word to measures of cylinder sets of shorter words via the production probabilities of the random substitution. This result first appeared in [\[17\]](#) and is central to the proof of the main result in [\[15\]](#).

Lemma 2.12 (Renormalisation). *Let ϑ_P be a primitive and compatible random substitution with corresponding frequency measure μ_P . Let $n \in \mathbb{N}$ and let k be an integer such that every $v \in \mathcal{L}_\vartheta^k$ has $|\vartheta(v)| \geq n + |\vartheta(v_1)|$. Then for every $u \in \mathcal{L}_\vartheta^n$,*

$$\mu_P([u]) = \frac{1}{\lambda} \sum_{v \in \mathcal{L}_\vartheta^k} \mu_P([v]) \sum_{j=1}^{|\vartheta(v_1)|} \mathbb{P}[\vartheta_P(v)_{[j, j+m-1]} = u].$$

[Lemma 2.12](#) is key to the proof of [Theorem A](#), as it relates the sums $\sum_{u \in \mathcal{L}_\vartheta^n} \mu_P([u])$ to sums over smaller words via the production probabilities. This in turn allows us

to obtain relations between τ_{μ_P} and φ_k . Under additional assumptions, simplified reformulations of [Lemma 2.12](#) can be obtained (see, for example, [Lemma 2.18](#), which is used in [Theorem D](#)).

2.9. Separation conditions and recognisability. In this section, we introduce additional common assumptions which either (1) impose a certain separation on inflation words, or (2) impose a certain uniformity of the inflation and the probabilities. Under these conditions, we can obtain closed-form formulas for the L^q -spectrum.

Definition 2.13. A random substitution $\vartheta_P = (\vartheta, P)$ satisfies the *disjoint set condition* if

$$u \text{ and } v \in \vartheta(a) \text{ with } u \neq v \implies \vartheta^k(u) \cap \vartheta^k(v) = \emptyset$$

for all $a \in \mathcal{A}$ and $k \in \mathbb{N}$. It satisfies the *identical set condition* if

$$u \text{ and } v \in \vartheta(a) \implies \vartheta^k(u) = \vartheta^k(v)$$

for all $a \in \mathcal{A}$ and $k \in \mathbb{N}$. Moreover, we say that ϑ_P has *identical production probabilities* if for all $a \in \mathcal{A}$, $k \in \mathbb{N}$ and $v \in \vartheta^k(a)$,

$$\mathbb{P}[\vartheta_P^{k-1}(u_1) = v] = \mathbb{P}[\vartheta_P^{k-1}(u_2) = v]$$

for all u_1 and $u_2 \in \vartheta(a)$.

A consequence of the disjoint set condition is that for every $a \in \mathcal{A}$, $k \in \mathbb{N}$ and $w \in \vartheta^k(a)$, there is a unique $v \in \vartheta^{k-1}(a)$ such that $w \in \vartheta(v)$. In other words, every exact inflation word can be uniquely de-substituted. The following definition extends this idea of unique de-substitution from inflation words to all elements in the subshift.

Definition 2.14. Let $\vartheta_P = (\vartheta, P)$ be a primitive and compatible random substitution. We call ϑ_P *recognisable* if for every $x \in X_\vartheta$ there exists a unique $y = \cdots y_{-1}y_0y_1 \cdots \in X_\vartheta$ and a unique integer $k \in \{0, \dots, |\vartheta(y_0)| - 1\}$ with $S^{-k}(x) \in \vartheta(y)$.

The following follows routinely from the definition of recognisability (a proof is given in [\[15, Lemma 4.5\]](#)).

Lemma 2.15. *If ϑ_P is a primitive, compatible and recognisable random substitution, then ϑ_P satisfies the disjoint set condition.*

In contrast to the disjoint set condition, recognisability is stable under taking powers (see [\[15, Lemma 4.6\]](#)).

Lemma 2.16. *Let ϑ_P be a primitive and compatible random substitution and $m \in \mathbb{N}$. If ϑ_P is recognisable, then so is ϑ_P^m .*

An alternative characterisation of recognisability is the following *local* version. Intuitively, local recognisability means that applying a *finite window* to a sequence is enough to determine the position and the type of the inflation word in the middle of that window. The following result is given in [\[15, Lemma 4.4\]](#) (see also [\[12, Proposition 5.7\]](#)).

Lemma 2.17. *Let $\vartheta_P = (\vartheta, P)$ be a primitive and compatible random substitution. If ϑ_P is recognisable, then there exists a smallest natural number $\kappa(\vartheta)$, called the recognisability radius of ϑ_P , with the following property: if $x \in \vartheta([a])$ for some $a \in \mathcal{A}$ and $x_{[-\kappa(\vartheta), \kappa(\vartheta)]} = y_{[-\kappa(\vartheta), \kappa(\vartheta)]}$ for some $y \in X_\vartheta$, then $y \in \vartheta([a])$.*

As a consequence of this local characterisation of recognisability, for every legal word u with length greater than twice the radius of recognisability there exists an inflation word w , appearing as a subword of u , which has a unique decomposition into exact inflation words. We call the largest such w the *recognisable core* of u .

Local recognisability allows us to obtain a stronger version of [Lemma 2.12](#) for recognisable random substitutions. This result is key to obtaining the coincidence of the L^q -spectrum and its inflation word analogue under recognisability for $q < 0$, and thus the conclusion of [Theorem D](#).

Lemma 2.18. *Let $\vartheta_P = (\vartheta, P)$ be a primitive and compatible random substitution, with corresponding frequency measure μ_P and $u \in \mathcal{L}_\vartheta$. If $v \in \mathcal{L}_\vartheta$ and $w \in \vartheta(v)$ contains u as a subword, then*

$$\mu_P([u]) \geq \frac{1}{\lambda} \mu_P([v]) \mathbb{P}[\vartheta_P(v) = w].$$

If, additionally, ϑ_P is recognisable, $|u| > 2\kappa(\vartheta)$ and w' is the recognisable core of u with $v' \in \mathcal{L}_\vartheta$ the unique legal word such that $w' \in \vartheta(v')$, then

$$\mu_P([u]) \leq \frac{\kappa(\vartheta)}{\lambda} \mu_P([v']) \mathbb{P}[\vartheta_P(v') = w'].$$

Proof. If u is a subword of $w \in \vartheta(v)$, then $\mu_P([u]) \geq \mu_P([w])$. Thus by [Lemma 2.12](#) applied to $\mu_P([w])$,

$$\mu_P([u]) \geq \frac{1}{\lambda} \mu_P([v]) \mathbb{P}[\vartheta_P(v) = w].$$

Now, assume that ϑ_P is recognisable, $|u| > 2\kappa(\vartheta)$ and $w' \in \vartheta(v')$ is the recognisable core of u . Let k be an integer such that every $t \in \mathcal{L}_\vartheta^k$ has $|\vartheta(t)| \geq k + |\vartheta(v_1)|$. Since there are at most $\kappa(\vartheta)$ letters of u preceding the recognisable core, if $t \in \mathcal{L}_\vartheta^k$ is a word for which $u \in \vartheta(t)_{[j, j+|u|-1]}$ for some $j \in \{1, \dots, |\vartheta(t_1)|\}$, then $t_i \cdots t_{i+|v|-1} = v'$ for some $i \in \{1, \dots, \kappa(\vartheta)\}$. Moreover, since there is a unique way to decompose w' into exact inflation words, for each $t \in \mathcal{L}_\vartheta^k$ there can be at most one $j \in \{1, \dots, \vartheta(t_1)\}$ such that

$u \in \vartheta(t)_{[j, j+|u|-1]}$. Hence, it follows by [Lemma 2.12](#) that

$$\begin{aligned} \mu_P([u]) &= \frac{1}{\lambda} \sum_{t \in \mathcal{L}^k} \mu_P([t]) \sum_{j=1}^{|\vartheta(t_1)|} \mathbb{P}[\vartheta_P(t)_{[j, j+|u|-1]} = u] \\ &\leq \frac{1}{\lambda} \sum_{i=1}^{\kappa(\vartheta)} \sum_{\substack{t \in \mathcal{L}_{\vartheta}^k \\ t_i \cdots t_{i+|v|-1} = v'}} \mu_P([t]) \mathbb{P}[\vartheta_P(v') = w'] \\ &= \frac{\kappa(\vartheta)}{\lambda} \mu_P([v']) \mathbb{P}[\vartheta_P(v') = w'], \end{aligned}$$

which completes the proof. \square

3. L^q -SPECTRA OF FREQUENCY MEASURES

In this section, we prove our main results on L^q -spectra of frequency measures. Here, we relate the L^q -spectrum to a certain “symbolic” L^q -spectrum, which we call the *inflation word L^q -spectrum*. Heuristically, the inflation word L^q -spectrum is the natural guess for the L^q -spectrum if you do not account for non-uniqueness in the positions in which legal words can appear in inflation words. This notion is introduced in [Section 3.1](#), where we also state and prove some of its key properties. In particular, in [Proposition 3.1](#), we prove a simple closed-form formula for the inflation word L^q -spectrum under the disjoint set condition or the identical set condition with identical production probabilities. In [Proposition 3.2](#), we establish basic monotonicity results.

Then, in [Section 3.2](#) and [Section 3.3](#), we establish the general bounds for the L^q -spectrum in terms of the inflation word L^q -spectrum, giving [Theorem A](#) (the proof is given in [Section 3.4](#)). We also prove that this bound is sharp in [Section 3.5](#), under the recognisability assumption. This proves the first part of [Theorem D](#). However this bound need not hold in general: we discuss a counterexample in [Example 5.2](#). Finally, in [Section 3.6](#), we prove differentiability of the L^q -spectrum at $q = 1$ and show how to recover known results for measure theoretic and topological entropy from our results concerning L^q -spectra.

3.1. Inflation word L^q -spectra. Given a primitive random substitution $\vartheta_P = (\vartheta, P)$, we can define an analogue of the L^q -spectrum in terms of its production probabilities, in a similar manner to the inflation word analogue of entropy introduced in [\[15\]](#). In many cases, this notion coincides with the L^q -spectrum of the frequency measure associated to ϑ_P . For each $k \in \mathbb{N}$ and $q \in \mathbb{R}$, define

$$\varphi_k(q) = - \sum_{a \in \mathcal{A}} R_a \log \left(\sum_{s \in \vartheta^k(a)} \mathbb{P}[\vartheta_P^k(a) = s]^q \right),$$

where $\mathbf{R} = (R_a)_{a \in \mathcal{A}}$ is the right Perron–Frobenius eigenvector of the substitution matrix of $\vartheta_{\mathbf{P}}$. We define the *inflation word L^q -spectrum* of $\vartheta_{\mathbf{P}}$ by

$$T_{\vartheta, \mathbf{P}}(q) = \liminf_{k \rightarrow \infty} \frac{\varphi_k(q)}{\lambda^k}.$$

We similarly define the upper variant $\overline{T}_{\vartheta, \mathbf{P}}$ by taking a limit supremum in place of the limit infimum.

We first state some key properties of $T_{\vartheta, \mathbf{P}}(q)$ which follow easily from the definition. Firstly, if the random substitution $\vartheta_{\mathbf{P}}$ is compatible and satisfies either the disjoint set condition or the identical set condition with identical production probabilities, then the limit defining $T_{\vartheta, \mathbf{P}}(q)$ exists for all $q \in \mathbb{R}$ and is given by a closed form expression. For $q \geq 0$, these properties transfer to the L^q -spectrum by [Theorem A](#).

Proposition 3.1. *Let $\vartheta_{\mathbf{P}}$ be a primitive and compatible random substitution and $q \in \mathbb{R}$. If $\vartheta_{\mathbf{P}}$ satisfies the disjoint set condition, then the limit defining $T_{\vartheta, \mathbf{P}}(q)$ exists and*

$$T_{\vartheta, \mathbf{P}}(q) = \frac{1}{\lambda - 1} \varphi_1(q).$$

If $\vartheta_{\mathbf{P}}$ satisfies the identical set condition and has identical production probabilities, then the limit defining $T_{\vartheta, \mathbf{P}}(q)$ exists and

$$T_{\vartheta, \mathbf{P}}(q) = \frac{1}{\lambda} \varphi_1(q).$$

Proof. Fix $q \in \mathbb{R}$. By the Markov property of $\vartheta_{\mathbf{P}}$, for all $a \in \mathcal{A}$, $k \in \mathbb{N}$ and $v \in \vartheta^k(a)$,

$$(3.1) \quad \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = v] = \sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = v].$$

If $\vartheta_{\mathbf{P}}$ satisfies the disjoint set condition, then for every $v \in \vartheta^k(a)$ there is a unique $s(v) \in \vartheta(a)$ such that $v \in \vartheta^{k-1}(s(v))$. Thus, for all $s \in \vartheta(a)$ such that $s \neq s(v)$, we have $\mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = v] = 0$, and so it follows by (3.1) that

$$\begin{aligned} \sum_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = v]^q &= \sum_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s(v)]^q \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s(v)) = v]^q \\ &= \sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \sum_{u \in \vartheta^{k-1}(s)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = u]^q \\ &= \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right) \cdot \prod_{b \in \mathcal{A}} \left(\sum_{u \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = u]^q \right)^{|\vartheta(a)|_b}, \end{aligned}$$

where in the final equality we use compatibility to split the second sum into inflation tiles. Thus

$$\begin{aligned}\varphi_k(q) &= - \sum_{a \in \mathcal{A}} R_a \sum_{b \in \mathcal{A}} |\vartheta(a)|_b \log \left(\sum_{u \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = u]^q \right) \\ &\quad - \sum_{a \in \mathcal{A}} R_a \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right) \\ &= \lambda \varphi_{k-1}(q) + \varphi_1(q),\end{aligned}$$

noting that $\sum_{a \in \mathcal{A}} R_a |\vartheta(a)|_b = \lambda R_b$. It follows inductively that

$$\frac{1}{\lambda^k} \varphi_k(q) = \sum_{j=1}^k \frac{1}{\lambda^j} \varphi_1(q) \xrightarrow{k \rightarrow \infty} \frac{1}{\lambda - 1} \varphi_1(q),$$

so the limit defining $T_{\vartheta, \mathbf{P}}(q)$ exists and is equal to $(\lambda - 1)^{-1} \varphi_1(q)$.

On the other hand, if $\vartheta_{\mathbf{P}}$ satisfies the identical set condition and has identical production probabilities, then $\mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s^1) = u] = \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s^2) = u]$ for all $s^1, s^2 \in \vartheta(a)$. Hence, it follows by (3.1) that

$$\sum_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = v]^q = \sum_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = v]^q$$

for any choice of $s \in \vartheta(a)$. By compatibility and the independence of the action,

$$\sum_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = v]^q = \prod_{b \in \mathcal{A}} \left(\sum_{u \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = u]^q \right)^{|\vartheta(a)|_b},$$

and thus

$$\varphi_k(q) = \sum_{b \in \mathcal{A}} \sum_{a \in \mathcal{A}} R_a |\vartheta(a)|_b \log \left(\sum_{v \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = v]^q \right) = \lambda \varphi_{k-1}(q),$$

noting that $\sum_{a \in \mathcal{A}} R_a |\vartheta(a)|_b = R_b$. It follows by induction that $\varphi_k(q)/\lambda^k = \varphi_1(q)/\lambda$ for all $k \in \mathbb{N}$, so we conclude that $T_{\vartheta, \mathbf{P}}(q)$ exists and equals $\lambda^{-1} \varphi_1(q)$. \square

Proposition 3.2. *Let $\vartheta_{\mathbf{P}}$ be a primitive and compatible random substitution. For all $q > 1$ and $q < 0$, the sequence $(\lambda^{-k} \varphi_k(q))_k$ is non-decreasing; and for all $0 < q < 1$, the sequence is non-increasing.*

Proof. This is largely a consequence of Jensen's inequality. Note that on the interval $(0, 1]$, the function $x \mapsto x^q$ is convex if $q > 1$ or $q < 0$, and concave if $0 < q < 1$. We

first prove this for the case when $q > 1$ or $q < 0$. Observe that for all $a \in \mathcal{A}$, $k \in \mathbb{N}$ with $k \geq 2$ and $v \in \vartheta^k(a)$, it follows by the Markov property of $\vartheta_{\mathbf{P}}$ that

$$\begin{aligned} \sum_{v \in \vartheta^k} \mathbb{P}[\vartheta_{\mathbf{P}}^k(a) = v]^q &= \sum_{v \in \vartheta^k(a)} \left(\sum_{s \in \vartheta(a): v \in \vartheta^{k-1}(s)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = v] \right)^q \\ &\leq \sum_{v \in \vartheta^k(a)} \left(\frac{\sum_{s \in \vartheta(a): v \in \vartheta^{k-1}(s)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s] \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = v]^q}{\sum_{s \in \vartheta(a): v \in \vartheta^{k-1}(s)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]} \right) \\ &\leq \prod_{b \in \mathcal{A}} \left(\sum_{w \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = w]^q \right)^{|\vartheta(a)|_b}. \end{aligned}$$

In the second line, we apply Jensen's inequality, and in the third line, we use compatibility to decompose each probability $\mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(s) = w]$ into inflation tiles. It follows that

$$\frac{1}{\lambda^k} \varphi_k(q) \geq -\frac{1}{\lambda^k} \sum_{b \in \mathcal{A}} R_b \sum_{a \in \mathcal{A}} R_a |\vartheta(a)|_b \log \left(\sum_{w \in \vartheta^{k-1}(b)} \mathbb{P}[\vartheta_{\mathbf{P}}^{k-1}(b) = w]^q \right) = \frac{1}{\lambda^{k-1}} \varphi_{k-1}(q),$$

noting that $\sum_{a \in \mathcal{A}} R_a |\vartheta(a)|_b = \lambda$.

The $0 < q < 1$ case follows similarly, with Jensen's inequality giving the opposite inequality since $x \mapsto x^q$ is concave. \square

An analogous monotonicity result does not hold in general for the $(\lambda^k - 1)^{-1} \varphi_k(q)$ bounds, even when $q \geq 0$. A counterexample is given by the random period doubling substitution (Example 5.7) with non-uniform probabilities.

3.2. L^q -spectra for non-negative q . The majority of the work in proving Theorem A lies in proving the bounds in (1.1), (1.2) and (1.3). Observe that it suffices to prove the bound for the case $k = 1$, since we then obtain the bound for other $k \in \mathbb{N}$ by considering higher powers of the random substitution. We first prove the upper bound for the case $q > 1$.

Throughout this section, we assume that the random substitution is primitive and compatible.

Proposition 3.3. *For all $q > 1$,*

$$\bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda - 1} \varphi_1(q).$$

Proof. Fix $q > 1$. Let $\varepsilon > 0$ and, for each $n \in \mathbb{N}$, let $m(n)$ be the integer defined by

$$m(n) = \left\lceil \frac{n}{\lambda - \varepsilon} \right\rceil.$$

Then the integers n and $m(n)$ satisfy the conditions of [Lemma 2.12](#), so it follows that

$$\sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_{\mathbf{P}}([u])^q = \sum_{u \in \mathcal{L}_{\vartheta}^n} \left(\frac{1}{\lambda} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v]) \sum_{j=1}^{|\vartheta(v_1)|} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j,j+n-1]} = u] \right)^q.$$

Since $q > 1$, the function $x \mapsto x^q$ is superadditive on the interval $[0, 1]$, so

$$\begin{aligned} \sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_{\mathbf{P}}([u])^q &\geq \sum_{u \in \mathcal{L}_{\vartheta}^n} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v])^q \left(\frac{1}{\lambda} \sum_{j=1}^{|\vartheta(v_1)|} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j,j+n-1]} = u] \right)^q \\ &\geq \frac{1}{\lambda^q} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v])^q \sum_{j=1}^{|\vartheta(v_1)|} \sum_{u \in \mathcal{L}_{\vartheta}^n} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j,j+n-1]} = u]^q. \end{aligned}$$

We now bound the probability on the right of this expression by the production probability of an inflation word. Observe that if $w(u) \in \vartheta(v)$ contains u as a subword in position j , then $\mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j,j+n-1]} = u] \geq \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w(u)]$. Hence,

$$\sum_{u \in \mathcal{L}_{\vartheta}^n} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j,j+n-1]} = u]^q \geq \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q$$

for all $j \in \{1, \dots, |\vartheta(v_1)|\}$.

Since $\vartheta_{\mathbf{P}}$ is compatible, by [Lemma 2.10](#) there exists an $N \in \mathbb{N}$ such that for all $n \geq N$ and all $v \in \mathcal{L}_{\vartheta}^{m(n)}$

$$\sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q \geq \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{m(n)(R_a + \varepsilon)}.$$

Hence,

$$\sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_{\mathbf{P}}([u])^q \geq \frac{1}{\lambda^q} \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{m(n)(R_a + \varepsilon)} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v])^q.$$

Taking logarithms, rearranging and dividing by n gives

$$\begin{aligned} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_{\mathbf{P}}([u])^q \right) &\leq -\frac{1}{n} \log \left(\sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v])^q \right) + \frac{1}{n} \log \lambda^q \\ &\quad - \frac{m(n)}{n} \sum_{a \in \mathcal{A}} (R_a + \varepsilon) \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right). \end{aligned}$$

Noting that $m(n)/n \rightarrow (\lambda - \varepsilon)^{-1}$ as $n \rightarrow \infty$, it follows by [Lemma 2.2](#) that

$$\bar{\tau}_{\mu_P}(q) \leq \frac{1}{\lambda - \varepsilon} \bar{\tau}_{\mu_P}(q) + \frac{1}{\lambda - \varepsilon} \sum_{a \in \mathcal{A}} \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \right) + c\varepsilon$$

where $c := (\#\mathcal{A}) \max_{a \in \mathcal{A}} \log(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q)$. But $\varepsilon > 0$ was arbitrary; letting $\varepsilon \rightarrow 0$ and rearranging we obtain

$$\bar{\tau}_{\mu_P}(q) \leq \frac{1}{\lambda - 1} \varphi_1(q),$$

which completes the proof. \square

We now prove the corresponding lower bound.

Proposition 3.4. *For all $q > 1$,*

$$\tau_{\mu_P}(q) \geq \frac{1}{\lambda} \varphi_1(q).$$

Proof. Let $\varepsilon > 0$ and, for each $n \in \mathbb{N}$, let $m(n)$ be the integer defined by

$$m(n) = \left\lceil \frac{n}{\lambda - \varepsilon} \right\rceil.$$

Since $q > 1$, the function $x \mapsto x^q$ is convex on the interval $[0, 1]$. Hence, it follows by [Lemma 2.12](#) and two applications of Jensen's inequality that

$$\begin{aligned} \sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_P([u])^q &= \sum_{u \in \mathcal{L}_{\vartheta}^n} \left(\frac{1}{\lambda} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v]) \sum_{j=1}^{|\vartheta(v_1)|} \mathbb{P}[\vartheta_P(v)_{[j, j+n-1]} = u] \right)^q \\ &\leq \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v]) \sum_{u \in \mathcal{L}_{\vartheta}^n} \left(\frac{1}{\lambda} \sum_{j=1}^{|\vartheta(v_1)|} \mathbb{P}[\vartheta_P(v)_{[j, j+n-1]} = u] \right)^q \\ &\leq \frac{|\vartheta|^{q-1}}{\lambda^q} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v]) \sum_{j=1}^{|\vartheta(v_1)|} \sum_{u \in \mathcal{L}_{\vartheta}^n} \mathbb{P}[\vartheta_P(v)_{[j, j+n-1]} = u]^q. \end{aligned}$$

We bound above the probability on the right of this expression by the production probability of a sufficiently large inflation word contained in u . By compatibility, there is an integer $k(n)$ such that $j + n \leq |\vartheta(v_{[1, m(n)-k(n)]})|$ for all $n \in \mathbb{N}$ and $v \in \mathcal{L}_{\vartheta}^{m(n)}$, where $\lim k(n)/n = 0$. In particular, for every $v \in \mathcal{L}_{\vartheta}^n$, a realisation of $\vartheta(v_{[2, m(n)-k(n)]})$ is contained in u as an inflation word, so

$$\sum_{u \in \mathcal{L}_{\vartheta}^n} \mathbb{P}[\vartheta_P(v)_{[j, j+n-1]} = u]^q \leq \sum_{w \in \vartheta(v_2 \cdots v_{m(n)-k(n)})} \mathbb{P}[\vartheta(v_2 \cdots v_{m(n)-k(n)}) = w]^q.$$

We now bound this quantity uniformly for all $v \in \mathcal{L}_\vartheta^{m(n)}$. By [Lemma 2.10](#) and the above, there is an $N \in \mathbb{N}$ such that for all $n \geq N$

$$\sum_{u \in \mathcal{L}_\vartheta^n} \mu_{\mathbf{P}}([u])^q \leq \frac{|\vartheta|^{q-1}}{\lambda^q} \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{(m(n)-k(n)-1)(R_a-\varepsilon)}.$$

Taking logarithms, rearranging and dividing by n gives

$$\begin{aligned} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}_\vartheta^n} \mu_{\mathbf{P}}([u])^q \right) &\geq \frac{m(n) - k(n) - 1}{n} \sum_{a \in \mathcal{A}} (R_a - \varepsilon) \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right) \\ &\quad - \frac{\log(|\vartheta|^{q-1}/\lambda^q)}{n} \\ &\xrightarrow{n \rightarrow \infty} \frac{1}{\lambda - \varepsilon} \sum_{a \in \mathcal{A}} (R_a - \varepsilon) \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right), \end{aligned}$$

But $\varepsilon > 0$ was arbitrarily, so

$$\tau_{\mu_{\mathbf{P}}}(q) \geq \frac{1}{\lambda} \varphi_1(q),$$

which completes the proof. \square

We now state the bounds for the $q \in (0, 1)$ case. We do not give a proof here since the arguments mirror the proofs of [Proposition 3.3](#) and [Proposition 3.4](#), except with reversed inequalities since $x \mapsto x^q$ is concave rather than convex and subadditive as opposed to superadditive.

Proposition 3.5. *If $q \in (0, 1)$, then*

$$\frac{1}{\lambda - 1} \varphi_1(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda} \varphi_1(q).$$

3.3. L^q -spectra for negative q : lower bounds. For $q < 0$, there exist primitive and compatible random substitutions for which $\tau_{\mu_{\mathbf{P}}}(q)$ and $T_{\vartheta, \mathbf{P}}(q)$ do not coincide (see, for instance, [Example 5.2](#)). However, we still obtain that $\tau_{\mu_{\mathbf{P}}}(q) \geq T_{\vartheta, \mathbf{P}}(q)$ for all $q < 0$. To prove this, it suffices to show the sequence of bounds in [\(1.3\)](#) holds. Again, we only need to prove the bound for $k = 1$ since the remaining bounds follow by considering powers of the random substitution.

Proposition 3.6. *If $\vartheta_{\mathbf{P}}$ is a primitive and compatible random substitution, then for all $q < 0$,*

$$\tau_{\mu_{\mathbf{P}}}(q) \geq \frac{1}{\lambda - 1} \varphi_1(q).$$

Proof. Let $\varepsilon > 0$ be sufficiently small and for n sufficiently large, let $m(n)$ be the integer defined by

$$m(n) = \left\lceil \frac{n}{\lambda - \varepsilon} \right\rceil.$$

To avoid division by zero, we rewrite [Lemma 2.12](#) in a form where we do not sum over elements equal to zero. Here, we write $u \blacktriangleleft \vartheta(v)$ to mean there is a realisation w of $\vartheta(v)$ for which u appears as a subword of w . For each $v \in \mathcal{L}_\vartheta^{m(n)}$ and $u \in \mathcal{L}_\vartheta^n$, let $\mathcal{J}(v, u) = \{j \in \{1, \dots, |\vartheta(v)|\} : u \in \vartheta(v)_{[j, j+n-1]}\}$. Observe that if $j \notin \mathcal{J}(u, v)$, then $\mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u] = 0$, and if u does not appear as a subword of any realisations of $\vartheta(v)$, then $\mathcal{J}(u, v) = \emptyset$. Therefore, we can rewrite [Lemma 2.12](#) as

$$\mu_{\mathbf{P}}([u]) = \frac{1}{\lambda} \sum_{\substack{v \in \mathcal{L}_\vartheta^{m(n)} \\ u \blacktriangleleft \vartheta(v)}} \mu_{\mathbf{P}}([v]) \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u].$$

Hence, by the subadditivity of the function $x \mapsto x^q$ on the domain $(0, 1]$,

$$\begin{aligned} \sum_{u \in \mathcal{L}_\vartheta^n} \mu_{\mathbf{P}}([u])^q &= \sum_{u \in \mathcal{L}_\vartheta^n} \left(\frac{1}{\lambda} \sum_{\substack{v \in \mathcal{L}_\vartheta^{m(n)} \\ u \blacktriangleleft \vartheta(v)}} \mu_{\mathbf{P}}([v]) \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u] \right)^q \\ &\leq \frac{1}{\lambda^q} \sum_{u \in \mathcal{L}_\vartheta^n} \sum_{\substack{v \in \mathcal{L}_\vartheta^{m(n)} \\ u \blacktriangleleft \vartheta(v)}} \mu_{\mathbf{P}}([v])^q \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u]^q \\ &= \frac{1}{\lambda^q} \sum_{v \in \mathcal{L}_\vartheta^{m(n)}} \mu_{\mathbf{P}}([v])^q \sum_{\substack{u \in \mathcal{L}_\vartheta^n \\ u \blacktriangleleft \vartheta(v)}} \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u]^q. \end{aligned}$$

For each $j \in \mathcal{J}(v, u)$, let $w_j(u) \in \vartheta(v)$ be a word such that $w_j(u)_{[j, j+n-1]} = u$. Note that there are at most $K := 2^{|\vartheta|}(\#\mathcal{A})^{|\vartheta|}$ different $u \in \mathcal{L}_\vartheta^n$ such that $w_j(u)_{[j, j+n-1]} = u$. Hence,

$$\begin{aligned} \sum_{\substack{u \in \mathcal{L}_\vartheta^n \\ u \blacktriangleleft \vartheta(v)}} \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j, j+n-1]} = u]^q &\leq \sum_{\substack{u \in \mathcal{L}_\vartheta^n \\ u \blacktriangleleft \vartheta(v)}} \sum_{j \in \mathcal{J}(v, u)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w_j(u)]^q \\ &\leq K \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q \end{aligned}$$

and it follows that

$$\sum_{u \in \mathcal{L}_\vartheta^n} \mu_{\mathbf{P}}([u])^q \leq \lambda^{-q} K \sum_{v \in \mathcal{L}_\vartheta^{m(n)}} \mu_{\mathbf{P}}([v])^q \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = w]^q.$$

Thus, by [Lemma 2.10](#), for all $\varepsilon > 0$ there is an integer N such that for all $n \geq N$

$$\sum_{u \in \mathcal{L}_\vartheta^n} \mu_{\mathbf{P}}([u])^q \leq \lambda^{-q} K \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right)^{m(n)(R_a + \varepsilon)} \left(\sum_{v \in \mathcal{L}_\vartheta^{m(n)}} \mu_{\mathbf{P}}([v])^q \right).$$

Taking logarithms, rearranging and dividing by n gives

$$\begin{aligned} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}_{\vartheta}^n} \mu_{\mathbf{P}}([u])^q \right) &\geq -\frac{1}{n} \log \left(\sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_{\mathbf{P}}([v])^q \right) + \frac{1}{n} \log(\lambda^{-q} K) \\ &\quad - \frac{m(n)}{n} \sum_{a \in \mathcal{A}} (R_a + \varepsilon) \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right). \end{aligned}$$

Noting that $m(n)/n \rightarrow (\lambda - \varepsilon)^{-1}$ as $n \rightarrow \infty$, it follows by [Lemma 2.2](#) that

$$\tau_{\mu_{\mathbf{P}}}(q) \geq \frac{1}{\lambda - \varepsilon} \tau_{\mu_{\mathbf{P}}}(q) + \frac{1}{\lambda - \varepsilon} \sum_{a \in \mathcal{A}} \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q \right) + c\varepsilon$$

where $c := (\#\mathcal{A}) \max_{a \in \mathcal{A}} \log(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q)$. Letting $\varepsilon \rightarrow 0$ and rearranging, we obtain

$$\bar{\tau}_{\mu_{\mathbf{P}}}(q) \geq \frac{1}{\lambda - 1} \varphi_1(q),$$

which completes the proof. \square

3.4. Proof of general bounds for the L^q spectrum. Using the bounds proven in the prior two sections, we can now complete the proof of [Theorem A](#).

Proof of Theorem A. Since, for each $k \in \mathbb{N}$, the random substitution $\vartheta_{\mathbf{P}}^k$ gives rise to the same frequency measure as $\vartheta_{\mathbf{P}}$, applying [Proposition 3.3](#), [Proposition 3.4](#) and [Proposition 3.5](#) to $\vartheta_{\mathbf{P}}^k$,

$$\frac{1}{\lambda^k} \varphi_k(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda^k - 1} \varphi_k(q)$$

for all $q > 1$ and

$$\frac{1}{\lambda^k - 1} \varphi_k(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda^k} \varphi_k(q)$$

for $0 < q < 1$. Letting $k \rightarrow \infty$ gives

$$\tau_{\mu_{\mathbf{P}}}(q) = \bar{\tau}_{\mu_{\mathbf{P}}}(q) = T_{\vartheta, \mathbf{P}}(q) = \bar{T}_{\vartheta, \mathbf{P}}(q)$$

for all $q \in (0, 1) \cup (1, \infty)$, so the limits defining $\tau_{\mu_{\mathbf{P}}}(q)$ and $T_{\vartheta, \mathbf{P}}(q)$ both exist and coincide. The same holds for $q = 0$ and $q = 1$ by continuity. The monotonicity of the bounds $\lambda^{-k} \varphi_k(q)$ follows by [Proposition 3.2](#). Finally for $q < 0$, for each $k \in \mathbb{N}$, applying [Proposition 3.6](#) to $\vartheta_{\mathbf{P}}^k$ gives that

$$\tau_{\mu_{\mathbf{P}}}(q) \geq \frac{1}{\lambda^k - 1} \varphi_k(q).$$

Passing to the limit completes the proof. \square

3.5. L^q -spectra for negative q under recognisability. While the upper bound does not hold in general for $q < 0$, for recognisable random substitutions we can obtain this using [Lemma 2.18](#), which we recall is a refinement of [Lemma 2.12](#) using recognisability.

Proposition 3.7. *If ϑ_P is a primitive, compatible and recognisable random substitution, then for all $q < 0$,*

$$\bar{\tau}_{\mu_P}(q) \leq \frac{1}{\lambda - 1} \varphi_1(q).$$

Proof. Let $\varepsilon > 0$ be sufficiently small and, for each $n \in \mathbb{N}$ sufficiently large, let $m(n)$ be the integer defined by

$$m(n) = \left\lfloor \frac{n}{\lambda - \varepsilon} \right\rfloor.$$

For each $u \in \mathcal{L}_{\vartheta}^{n+2\kappa(\vartheta)}$, let $w(u)$ denote the recognisable core of u . Further, let $v(u)$ denote the unique legal word such that $w(u) \in \vartheta(v(u))$. Then, by [Lemma 2.18](#), we have

$$(3.2) \quad \mu_P([u]) \leq \frac{\kappa(\vartheta)}{\lambda} \mu_P([v(u)]) \mathbb{P}[\vartheta(v(u)) = w(u)].$$

Observe that for all $u \in \mathcal{L}_{\vartheta}^{n+2\kappa(\vartheta)}$, the recognisable core $w(u)$ has length at least n so, by compatibility, there is an integer N such that if $n \geq N$, then $|v(u)| \geq m(n)$ for all $u \in \mathcal{L}_{\vartheta}^{n+2\kappa(\vartheta)}$. In particular, for every u there exists a $v \in \mathcal{L}_{\vartheta}^{m(n)}$ such that $\mu_P([v(u)]) \leq \mu_P([v])$ and a $w \in \vartheta(v)$ such that $\mathbb{P}[\vartheta_P(v(u)) = w(u)] \leq \mathbb{P}[\vartheta_P(v) = w]$. Hence, it follows by (3.2) and [Lemma 2.10](#) that

$$\begin{aligned} \sum_{u \in \mathcal{L}_{\vartheta}^{n+2\kappa(\vartheta)}} \mu_P([u])^q &\geq \frac{1}{\lambda^q} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v])^q \sum_{w \in \vartheta(v)} \mathbb{P}[\vartheta_P(v) = w]^q \\ &\geq \frac{1}{\lambda^q} \prod_{a \in \mathcal{A}} \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \right)^{m(R_a - \varepsilon)} \sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v])^q, \end{aligned}$$

noting that since $q < 0$, the function $x \mapsto x^q$ is decreasing on $(0, 1]$. Taking logarithms, rearranging and dividing by n gives

$$\begin{aligned} -\frac{1}{n} \log \left(\sum_{u \in \mathcal{L}_{\vartheta}^{n+2\kappa(\vartheta)}} \mu_P([u])^q \right) &\leq -\frac{1}{n} \log \left(\sum_{v \in \mathcal{L}_{\vartheta}^{m(n)}} \mu_P([v])^q \right) + \frac{1}{n} \log \lambda^q \\ &\quad - \frac{m(n)}{n} \sum_{a \in \mathcal{A}} (R_a - \varepsilon) \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \right). \end{aligned}$$

Noting that $m(n)/n \rightarrow (\lambda - \varepsilon)^{-1}$ as $n \rightarrow \infty$, it follows by [Lemma 2.2](#) that

$$\bar{\tau}_{\mu_P}(q) \leq \frac{1}{\lambda - \varepsilon} \bar{\tau}_{\mu_P}(q) + \frac{1}{\lambda - \varepsilon} \sum_{a \in \mathcal{A}} \log \left(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q \right) + c\varepsilon$$

where $c := (\#\mathcal{A}) \max_{a \in \mathcal{A}} \log(\sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_P(a) = s]^q)$. Letting $\varepsilon \rightarrow 0$ and rearranging, we obtain

$$\bar{\tau}_{\mu_P}(q) \leq \frac{1}{\lambda - 1} \varphi_1(q),$$

which completes the proof. \square

3.6. Recovering entropy from the L^q -spectrum. Since the L^q -spectrum encodes both topological and measure theoretic entropy, [Theorem A](#) provides an alternative means of proving the coincidence of these quantities with the inflation word analogues introduced in [\[14, 15\]](#).

For notational simplicity, set

$$\rho_k = - \sum_{a \in \mathcal{A}} R_a \sum_{s \in \vartheta^k(a)} \mathbb{P}[\vartheta_P^k(a) = s] \log(\mathbb{P}[\vartheta_P^k(a) = s]).$$

Proof of Corollary C. We first establish the result for topological entropy. By [Theorem A](#), the limit defining $T_{\vartheta, P}(0)$ exists; in particular,

$$\lim_{m \rightarrow \infty} \frac{1}{\lambda^k} \sum_{a \in \mathcal{A}} R_a \log(\#\vartheta^m(a))$$

exists. Since $h_{\text{top}}(X_{\vartheta}) = -\tau_{\mu_P}(0) = -T_{\vartheta, P}(0)$, we conclude that

$$h_{\text{top}}(X_{\vartheta}) = - \lim_{m \rightarrow \infty} \frac{1}{\lambda^k} \sum_{a \in \mathcal{A}} R_a \log(\#\vartheta^m(a))$$

as claimed.

Now we consider measure theoretic entropy. We first make the following elementary observation: if f and g are concave functions with $f(1) = g(1)$ and $f(x) \leq g(x)$ for all $x \geq 1$, then $f^+(1) \leq g^+(1)$. Indeed, for all $\epsilon > 0$,

$$\frac{f(1 + \epsilon) - f(1)}{\epsilon} \leq \frac{g(1 + \epsilon) - g(1)}{\epsilon},$$

and taking the limit as ϵ goes to 0 (which always exists by concavity) yields the desired inequality.

Recall that τ_{μ_P} and $\lambda^{-k} \varphi_k$ are concave functions with $\tau_{\mu_P}(1) = \varphi_k(1) = 0$ for all $k \in \mathbb{N}$. Moreover, φ_k is differentiable for all $k \in \mathbb{N}$ with $\varphi'_k(1) = \rho_k$ and by [Proposition 3.2](#) and [Theorem A](#), $(\lambda^{-k} \varphi_k)_{j=1}^{\infty}$ converges monotonically to τ_{μ_P} from

below. In particular, ρ_k/λ^k is a monotonically increasing sequence bounded above by $\tau_{\mu_P}^+(1)$, so that the limit indeed exists. Thus

$$\tau_{\mu_P}^+(1) = \lim_{k \rightarrow \infty} \frac{\rho_k}{\lambda^k}$$

since $\varphi_k(q)/(\lambda^k - 1) \geq \tau_{\mu_P}(q)$ for all $q \in (0, \infty)$, using the preceding observation.

The result for $\tau_{\mu_P}^-(1)$ follows by an identical argument, instead using monotonicity and the corresponding bounds for $q \in (0, 1)$. Thus $\tau_{\mu_P}'(1) = \lim_{k \rightarrow \infty} \rho_k/\lambda^k$, so the desired result follows by [Lemma 2.1\(c\)](#). \square

4. RECOGNISABILITY AND THE MULTIFRACTAL FORMALISM

In this section we establish the multifractal formalism as stated in [Theorem D](#). To do this, we prove a variational principle by considering typical local dimensions of one frequency measure μ_P relative to another frequency measure μ_Q . Our strategy is to prove the almost-sure existence of *relative letter frequencies* in [Lemma 4.3](#): this result, combined with recognisability, gives [Proposition 4.5](#). The multifractal formalism then follows from this dimensional result combined with the formula for the L^q -spectrum proven in [Proposition 3.7](#)—the proof is given in [Section 4.2](#).

4.1. Non-typical local dimensions. To prove the multifractal formalism for a given frequency measure μ_P , we show that for every $\alpha \in [\alpha_{\min}, \alpha_{\max}]$, there exists another frequency measure μ_Q such that $\dim_{\mathbb{H}} \mu_Q \geq \tau_{\mu_P}^*(\alpha)$ and $\dim_{\text{loc}}(\mu_P, x) = \alpha$ for μ_Q -almost every $x \in X_{\vartheta}$. Given a primitive set-valued substitution ϑ , permissible probabilities P and Q , $m \in \mathbb{N}$ and $a \in \mathcal{A}$, define the quantity $H_{P,Q}^{m,a}(\vartheta)$ by

$$H_{P,Q}^{m,a}(\vartheta) = \sum_{v \in \vartheta^m(a)} -\mathbb{P}[\vartheta_Q^m(a) = v] \log \mathbb{P}[\vartheta_P^m(a) = v].$$

Further, let $\mathbf{H}_{P,Q}^m(\vartheta)$ denote the vector $(H_{P,Q}^{m,a}(\vartheta))_{a \in \mathcal{A}}$. We first prove some properties of the quantity $\mathbf{H}_{P,Q}^m(\vartheta)$ which we will use in the proof of [Proposition 4.5](#).

Lemma 4.1. *If ϑ is a primitive and compatible set-valued substitution and P and Q are permissible probabilities, then for all $m \in \mathbb{N}$, $a \in \mathcal{A}$ and $s \in \vartheta(a)$,*

$$\sum_{v \in \vartheta^m(s)} \mathbb{P}[\vartheta_Q^m(s) = v] \log \mathbb{P}[\vartheta_P^m(s) = v] = \sum_{b \in \mathcal{A}} |\vartheta(a)|_b H_{P,Q}^{m,b}(\vartheta).$$

Proof. Since ϑ is compatible, we can decompose each $v \in \vartheta^m(s)$ into inflation words $v = v^1 \dots v^{|\vartheta(a)|}$. By the Markov property of ϑ_P (respectively ϑ_Q), we have that

$$\mathbb{P}[\vartheta_P^m(s) = v] = \mathbb{P}[\vartheta_P^m(s_1) = v^1] \dots \mathbb{P}[\vartheta_P^m(s_{|\vartheta(a)|}) = v^{|\vartheta(a)|}].$$

Therefore

$$\begin{aligned} \sum_{v \in \vartheta^m(s)} \mathbb{P}[\vartheta_Q^m(s) = v] \log \mathbb{P}[\vartheta_P^m(s) = v] &= \sum_{b \in \mathcal{A}} |\vartheta(a)|_b \sum_{w \in \vartheta^m(b)} \mathbb{P}[\vartheta_Q^m(b) = w] \log \mathbb{P}[\vartheta_P^m(b) = w] \\ &= \sum_{b \in \mathcal{A}} |\vartheta(a)|_b H_{P,Q}^{m,b}(\vartheta), \end{aligned}$$

which completes the proof. \square

Lemma 4.2. *If ϑ is a primitive and compatible set-valued substitution satisfying the disjoint set condition, with right Perron–Frobenius eigenvector \mathbf{R} , and \mathbf{P} and \mathbf{Q} are permissible probabilities, then the random substitutions $\vartheta_P = (\vartheta, \mathbf{P})$ and $\vartheta_Q = (\vartheta, \mathbf{Q})$ satisfy*

$$\frac{1}{\lambda^m} \mathbf{H}_{P,Q}^m(\vartheta) \cdot \mathbf{R} \rightarrow \frac{1}{\lambda - 1} \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R}$$

as $m \rightarrow \infty$.

Proof. Since ϑ satisfies the disjoint set condition, for all $m \in \mathbb{N}$ and $a \in \mathcal{A}$,

$$\begin{aligned} \mathbf{H}_{P,Q}^{m+1}(\vartheta) \cdot \mathbf{R} &= \sum_{a \in \mathcal{A}} R_a \sum_{v \in \vartheta^{m+1}(a)} \mathbb{P}[\vartheta_Q^{m+1}(a) = v] \log \mathbb{P}[\vartheta_P^{m+1}(a) = v] \\ &= \sum_{a \in \mathcal{A}} R_a \sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_Q(a) = s] \log \mathbb{P}[\vartheta_P(a) = s] \\ &\quad + \sum_{a \in \mathcal{A}} R_a \sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_Q(a) = s] \sum_{v \in \vartheta^m(s)} \mathbb{P}[\vartheta_Q^m(s) = v] \log \mathbb{P}[\vartheta_P^m(s) = v] \\ &= \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R} + \sum_{b \in \mathcal{A}} H_{P,Q}^{m,b}(\vartheta) \sum_{a \in \mathcal{A}} |\vartheta(a)|_b R_a \\ &= \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R} + \lambda \sum_{b \in \mathcal{A}} R_b H_{P,Q}^{m,b}(\vartheta) \\ &= \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R} + \lambda \mathbf{H}_{P,Q}^m(\vartheta) \cdot \mathbf{R}. \end{aligned}$$

In the second equality we use the Markov property of ϑ_P and ϑ_Q , laws of logarithms, and that $\sum_{v \in \vartheta^m(s)} \mathbb{P}[\vartheta_Q^m(s) = v] = 1$ for all $s \in \vartheta(a)$; in the third we apply [Lemma 4.1](#); in the fourth we use that $M_\vartheta \mathbf{R} = \lambda \mathbf{R}$. Applying the above inductively,

$$\frac{1}{\lambda^m} \mathbf{H}_{P,Q}^m(\vartheta) \cdot \mathbf{R} = \sum_{j=1}^m \frac{1}{\lambda^j} \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R} \xrightarrow{m \rightarrow \infty} \frac{1}{\lambda - 1} \mathbf{H}_{P,Q}^1(\vartheta) \cdot \mathbf{R},$$

which completes the proof. \square

Any bi-infinite sequence x in the subshift of a recognisable random substitution can be written as a bi-infinite concatenation of exact inflation words (w^{n,a_n}) , where w^{n,a_n} is an inflation word generated from the letter a_n . Given a recognisable set-valued

substitution ϑ , $a \in \mathcal{A}$ and $w \in \vartheta(a)$, we define the inflation word frequency of (a, w) in $x \in X_\vartheta$ by

$$f_x(a, w) = \lim_{n \rightarrow \infty} f_x^n(a, w)$$

$$f_x^n(a, w) = \frac{1}{2n+1} \# \{m : a_m = a, w^{m, a_m} = w, w^{m, a_m} \text{ in } x_{[-n, n]}\},$$

provided the limit exists. For a given frequency measure μ_P , the inflation word frequency of a μ_P -typical word is determined by the production probabilities. More specifically, we have the following.

Lemma 4.3. *Let $\vartheta_P = (\vartheta, P)$ be a primitive, compatible and recognisable random substitution with corresponding frequency measure μ_P . For μ_P -almost every $x \in X_\vartheta$, the inflation word frequency exists and is given by*

$$f_x(a, w) = \frac{1}{\lambda} R_a \mathbb{P}[\vartheta_P(a) = w],$$

for all $a \in \mathcal{A}$ and $w \in \vartheta(a)$.

Proof. Let $A_{a,w}$ be the set of points $x \in X_\vartheta$ such that the above does *not* hold. We show that $A_{a,w}$ is a null set. Taking the complement and then the intersection over all a, w gives a full-measure set with the required property. Given $\varepsilon > 0$, let $E(n, \varepsilon)$ be the set of $x \in X_\vartheta$ such that

$$|f_x^n(a, w) - \frac{1}{\lambda} R_a \mathbb{P}[\vartheta_P(a) = w]| > \varepsilon.$$

By the Borel–Cantelli lemma, it suffices to show that

$$\sum_{n \in \mathbb{N}} \mu_P(E(n, \varepsilon)) < \infty$$

for all $\varepsilon > 0$ in order to conclude that $A_{a,w}$ is a nullset. To this end, we show that $\mu_P(E(n, \varepsilon))$ decays exponentially with n . Given u with $|u| = 2n+1 > 2\kappa(\vartheta)$, let u^R denote the recognisable core of u , which has length at least $|u| - 2\kappa(\vartheta)$. [Lemma 2.18](#) gives that

$$\mu_P([u]) \leq \frac{\kappa(\vartheta)}{\lambda} \mu_P([v]) \mathbb{P}[\vartheta_P(v) = u^R] = \frac{\kappa(\vartheta)}{\lambda} \mu_P([v]) \prod_{i=1}^{|v|} \mathbb{P}[\vartheta_P(v_i) = w^{i, v_i}]$$

where each w^{i, v_i} is the inflated image of v_i in u^R . By compatibility, we can choose an integer N such that every v of length at least N satisfies $|v|(R_a - \varepsilon/3) \leq |v|_a \leq |v|(R_a + \varepsilon/3)$ for all $a \in \mathcal{A}$. For each v and $a \in \mathcal{A}$, let $A_a(v)$ denote the set of $u' \in \vartheta(v)$ such that the frequency of indices $i \in \{j : a_j = a\}$ with $w^{i, a} = w$ deviates from $\mathbb{P}[\vartheta(a) = w]$ by more than $\varepsilon/3$. Since ϑ_P acts independently on letters, it follows by Cramér’s theorem that the sum $\sum_{u' \in A_a(v)} \mathbb{P}[\vartheta_P(v) = u']$ decays exponentially with $|v|_a$

(and hence with $|v|$). In particular, there is a constant $C > 0$, independent of the choice of v , such that

$$(4.1) \quad \sum_{u' \in A(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = u'] \leq e^{-Cn}.$$

Note that if u is a sufficiently long legal word and has $[u] \cap E(n, \varepsilon) = \emptyset$, then we require that $u^R \in A(v)$. Indeed, if $u' \notin A(v)$ and $|v| \geq N$, then the relative inflation word frequency of w is bounded above by

$$\begin{aligned} \frac{|\{j : a_j = a\}|}{|v|} \frac{|v|}{|u|} \left(\mathbb{P}[\vartheta_{\mathbf{P}}(a) = w] + \frac{\varepsilon}{3} \right) &\leq \frac{1}{\lambda} \left(R_a + \frac{\varepsilon}{3} \right) \left(\mathbb{P}[\vartheta_{\mathbf{P}}(a) = w] + \frac{\varepsilon}{3} \right) \\ &\leq \frac{1}{\lambda} R_a \mathbb{P}[\vartheta_{\mathbf{P}}(a) = w] + \varepsilon \end{aligned}$$

and, similarly, bounded below by $R_a \mathbb{P}[\vartheta_{\mathbf{P}}(a) = w] / \lambda - \varepsilon$; hence, $[u^R] \cap E(n, \varepsilon) = \emptyset$. Let \mathcal{V}_n denote set of all words which appear as the (unique) preimage of the recognisable core of a word of length n . It then follows by [Lemma 2.18](#) that

$$\mu_{\mathbf{P}}(E(n, \varepsilon)) \leq \sum_{\substack{u \in \mathcal{L}_{\vartheta}^n \\ [u] \cap E(n, \varepsilon) \neq \emptyset}} \mu_{\mathbf{P}}([u]) \leq \frac{\kappa(\vartheta)}{\lambda} \sum_{v \in \mathcal{V}_n} \mu_{\mathbf{P}}([v]) \sum_{u' \in A(v)} \mathbb{P}[\vartheta_{\mathbf{P}}(v) = u'] \leq e^{-Cn},$$

where in the final inequality we have used (4.1) and that

$$\sum_{v \in \mathcal{V}_n} \mu_{\mathbf{P}}([v]) \leq \sum_{j=1}^n \sum_{v \in \mathcal{L}_{\vartheta}^j} \mu_{\mathbf{P}}([v]) \leq n,$$

absorbing this contribution and the $\kappa(\vartheta)/\lambda$ factor into the constant C . It follows that

$$\sum_{n=1}^{\infty} \mu_{\mathbf{P}}(E(n, \varepsilon)) \leq \sum_{n=1}^{\infty} e^{-Cn} < \infty,$$

and the result then follows by the Borel–Cantelli lemma. \square

Finally, we require the following bounds on the exponential scaling rate of measures of cylinders, which is essentially a consequence of [Theorem A](#). In particular, these give bounds on the possible local dimensions of the measure.

Proposition 4.4. *If $\vartheta_{\mathbf{P}}$ is a primitive and compatible random substitution, then there are values $0 < s_1 < s_2 < \infty$ and $c_1, c_2 > 0$ such that for all $n \in \mathbb{N}$ and $v \in \mathcal{L}^n(X_{\vartheta})$,*

$$s_1 \cdot n + c_1 \leq \log \mu_{\mathbf{P}}([v]) \leq s_2 \cdot n + c_2$$

Proof. By [Theorem A](#), for all $k \in \mathbb{N}$ and $q > 1$,

$$\tau_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda^k - 1} \varphi_k(q);$$

and for $q < 0$,

$$\frac{1}{\lambda^k - 1} \varphi_k(q) \leq \tau_{\mu_P}(q),$$

Moreover, for each k , with

$$\begin{aligned} \beta_{k,\min} &:= \lim_{q \rightarrow \infty} \frac{\varphi_k(q)}{q(\lambda^k - 1)} = -\frac{1}{\lambda^k - 1} \sum_{a \in \mathcal{A}} R_a \log \left(\min_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_P^k(a) = v] \right) \\ \beta_{k,\max} &:= \lim_{q \rightarrow -\infty} \frac{\varphi_k(q)}{q(\lambda^k - 1)} = -\frac{1}{\lambda^k - 1} \sum_{a \in \mathcal{A}} R_a \log \left(\max_{v \in \vartheta^k(a)} \mathbb{P}[\vartheta_P^k(a) = v] \right), \end{aligned}$$

it follows that $[\beta_{k,\min}, \beta_{k,\max}] \subset (0, \infty)$ is a decreasing nested sequence of intervals, so with $\beta_{\min} = \lim_{k \rightarrow \infty} \beta_{k,\min}$ and $\beta_{\max} = \lim_{k \rightarrow \infty} \beta_{k,\max}$,

$$0 < \beta_{\min} \leq \lim_{q \rightarrow \infty} \tau_{\mu_P}(q) \leq \lim_{q \rightarrow -\infty} \tau_{\mu_P}(q) \leq \beta_{\max} < \infty.$$

Applying [Lemma 2.1\(b\)](#) gives the result. \square

Finally, we obtain our main conclusion concerning relative local dimensions.

Proposition 4.5. *Let ϑ be a primitive, compatible and recognisable set-valued substitution, let P and Q be permissible probabilities, and let μ_P and μ_Q denote the frequency measure corresponding to ϑ endowed with the probabilities P and Q , respectively. Then, for μ_Q -almost all $x \in X_\vartheta$,*

$$(4.2) \quad \dim_{\text{loc}}(\mu_P, x) = \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a \sum_{v \in \vartheta(a)} -\mathbb{P}[\vartheta_Q^m(a) = v] \log \mathbb{P}[\vartheta_P^m(a) = v].$$

Proof. Fix $m \in \mathbb{N}$. It follows by [Lemma 2.16](#) that since ϑ_P is recognisable, so is ϑ_P^m . For each $x \in X_\vartheta$ and $n \in \mathbb{N}$ with $n > \kappa(\vartheta^m)$, let $u_-^n(x)$ denote the recognisable core of $x_{[-n,n]}$ and let $u_+^n(x)$ denote an inflation word of minimal length that contains $x_{[-n,n]}$. By compatibility, $|u_-^n(x)|/(2n+1) \rightarrow \lambda^{-m}$ and $|u_+^n(x)|/(2n+1) \rightarrow \lambda^{-m}$ as $n \rightarrow \infty$. Further, let $v_-^n(x)$ be the legal word such that $u_-^n(x) \in \vartheta^m(v_-^n(x))$ and $v_+^n(x)$ be the legal word such that $u_+^n(x) \in \vartheta^m(v_+^n(x))$. Then, it follows by [Lemma 2.18](#) and the definition of local dimension that

$$\begin{aligned} &\liminf_{n \rightarrow \infty} \left(-\frac{1}{2n+1} \log \mu_P([u_-^n(x)]) - \frac{1}{2n+1} \log \mathbb{P}[\vartheta_P(v_-^n(x)) = u_-^n(x)] \right) \\ &\leq \underline{\dim}_{\text{loc}}(\mu_P, x) \leq \overline{\dim}_{\text{loc}}(\mu_P, x) \\ &\leq \limsup_{n \rightarrow \infty} \left(-\frac{1}{2n+1} \log \mu_P([u_+^n(x)]) - \frac{1}{2n+1} \log \mathbb{P}[\vartheta_P(v_+^n(x)) = u_+^n(x)] \right). \end{aligned}$$

By [Proposition 4.4](#), there exists a constant $C \geq 0$ such that for all $x \in X_\vartheta$,

$$0 \leq \liminf_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mu_P([u_-^n(x)]) \leq \limsup_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mu_P([u_+^n(x)]) \leq C.$$

Hence, it follows from the above that

$$\begin{aligned}
 (4.3) \quad & \liminf_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mathbb{P}[\vartheta_{\mathbf{P}}(v_-^n(x)) = u_-^n(x)] \\
 & \leq \underline{\dim}_{\text{loc}}(\mu_{\mathbf{P}}, x) \leq \overline{\dim}_{\text{loc}}(\mu_{\mathbf{P}}, x) \\
 & \leq \limsup_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mathbb{P}[\vartheta_{\mathbf{P}}(v_+^n(x)) = u_+^n(x)] + \frac{C}{\lambda^m}.
 \end{aligned}$$

We now show that for μ_Q -almost all $x \in X_{\vartheta}$,

$$\begin{aligned}
 \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}[\vartheta_{\mathbf{P}}(v_-^n(x)) = u_-^n(x)] &= \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \mathbb{P}[\vartheta_{\mathbf{P}}(v_+^n(x)) = u_+^n(x)] \\
 &= \frac{1}{\lambda^m} \mathbf{H}_{\mathbf{P}, \mathbf{Q}}^m(\vartheta) \cdot \mathbf{R}.
 \end{aligned}$$

By compatibility, we can decompose the production probabilities into inflation tiles as

$$\mathbb{P}[\vartheta_{\mathbf{P}}^m(v_-^n(x)) = u_-^n(x)] = \prod_{a \in \mathcal{A}} \prod_{w \in \vartheta^m(a)} \mathbb{P}[\vartheta_{\mathbf{P}}^m(a) = w]^{N_{a,w}(x,n)},$$

where, for each $a \in \mathcal{A}$ and $w \in \vartheta^m(a)$, $N_{a,w}(x,n)$ denotes the number of a 's in $v_-^n(x)$ which map to w . It follows by [Lemma 4.3](#), applied to $\vartheta_{\mathbf{Q}}^m$, that for μ_Q -almost all $x \in X_{\vartheta}$, we have

$$\frac{1}{2n+1} N_{a,w}(x,n) \rightarrow \frac{1}{\lambda^m} R_a \mathbb{P}[\vartheta_{\mathbf{Q}}^m(a) = w]$$

for all $a \in \mathcal{A}$ and $w \in \vartheta^m(a)$. Hence, it follows that

$$\begin{aligned}
 & \lim_{n \rightarrow \infty} -\frac{1}{2n+1} \log \mathbb{P}[\vartheta_{\mathbf{P}}^m(v_-^n(x)) = u_-^n(x)] \\
 &= \frac{1}{\lambda^m} \sum_{a \in \mathcal{A}} R_a \sum_{v \in \vartheta^m(a)} \mathbb{P}[\vartheta_{\mathbf{Q}}^m(a) = v] \log \mathbb{P}[\vartheta_{\mathbf{P}}^m(a) = v] \\
 &= \frac{1}{\lambda^m} \mathbf{H}_{\mathbf{P}, \mathbf{Q}}^m(\vartheta) \cdot \mathbf{R},
 \end{aligned}$$

with the same convergence holding for $u_+^n(x)$ by identical arguments. Thus, it follows from [\(4.3\)](#) that

$$\frac{1}{\lambda^m} \mathbf{H}_{\mathbf{P}, \mathbf{Q}}^m(\vartheta) \cdot \mathbf{R} \leq \underline{\dim}_{\text{loc}}(\mu_{\mathbf{P}}, x) \leq \overline{\dim}_{\text{loc}}(\mu_{\mathbf{P}}, x) \leq \frac{1}{\lambda^m} \mathbf{H}_{\mathbf{P}, \mathbf{Q}}^m(\vartheta) \cdot \mathbf{R} + \frac{C}{\lambda^m}.$$

Since the above holds for all $m \in \mathbb{N}$, by letting $m \rightarrow \infty$ it follows by [Lemma 4.2](#) that $\dim_{\text{loc}}(\mu_{\mathbf{P}}, x)$ exists and

$$\dim_{\text{loc}}(\mu_{\mathbf{P}}, x) = \frac{1}{\lambda - 1} \mathbf{H}_{\mathbf{P}, \mathbf{Q}}^1(\vartheta) \cdot \mathbf{R},$$

which completes the proof. \square

4.2. Proof of the multifractal formalism. In this section, we apply the results obtained in the previous section, along with results on the L^q -spectrum under recognisability, to prove [Theorem D](#).

Proof of Theorem D. We first obtain the results for the L^q -spectrum. Since every recognisable random substitution satisfies the disjoint set condition, [Proposition 3.1](#) gives that $T_{\vartheta, \mathbf{P}}(q) = (\lambda - 1)^{-1} \varphi_1(q)$ for all $q \in \mathbb{R}$. If $q < 0$, then by [Theorem A](#) and [Proposition 3.7](#),

$$\frac{1}{\lambda - 1} \varphi_1(q) = T_{\vartheta, \mathbf{P}}(q) \leq \tau_{\mu_{\mathbf{P}}}(q) \leq \bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq \frac{1}{\lambda - 1} \varphi_1(q),$$

so we conclude that $\tau_{\mu_{\mathbf{P}}}(q)$ exists and equals $(\lambda - 1)^{-1} \varphi_1(q)$. For $q \geq 0$, the result follows already from [Corollary B](#).

We now obtain the results on the multifractal spectrum. In light of [Proposition 2.3](#), it remains to show that $f_{\mu_{\mathbf{P}}}(\alpha) \geq \tau_{\mu_{\mathbf{P}}}^*(\alpha)$ for each $\alpha \in \mathbb{R}$. As proven above, for all $q \in \mathbb{R}$,

$$\tau_{\mu_{\mathbf{P}}}(q) = \frac{1}{\lambda - 1} \varphi_1(q) = \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a T_a(q)$$

where for each $a \in \mathcal{A}$

$$T_a(q) = -\log \sum_{s \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q.$$

First, fix $\alpha \in (\alpha_{\min}, \alpha_{\max})$ and let $q \in \mathbb{R}$ be chosen so that $\tau'_{\mu_{\mathbf{P}}}(q) = \alpha$. Observe that $q\alpha - \tau_{\mu_{\mathbf{P}}}(q) = \tau_{\mu_{\mathbf{P}}}^*(\alpha)$. Then define \mathbf{Q} by the rule

$$\mathbb{P}[\vartheta_{\mathbf{Q}}(a) = s] = \mathbb{P}[\vartheta_{\mathbf{P}}(a) = s]^q e^{T_a(q)}$$

for all $a \in \mathcal{A}$ and $s \in \vartheta(a)$. Then by [Corollary C](#),

$$\begin{aligned} \dim_{\mathbb{H}} \mu_{\mathbf{Q}} &= \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a \left(- \sum_{v \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{Q}}(a) = v] \log \mathbb{P}[\vartheta_{\mathbf{Q}}(a) = v] \right) \\ &= q \cdot \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a \left(- \sum_{v \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{Q}}(a) = v] \log \mathbb{P}[\vartheta_{\mathbf{P}}(a) = v] \right) \\ &\quad - \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a T_a(q) \sum_{v \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{Q}}(a) = v] \\ &= q\alpha - \tau_{\mu_{\mathbf{P}}}(q) = \tau_{\mu_{\mathbf{P}}}^*(\alpha) \end{aligned}$$

since

$$\begin{aligned} \tau'_{\mu_{\mathbf{P}}}(q) &= \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a \frac{- \sum_{v \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{P}}(a) = v]^q \log \mathbb{P}[\vartheta_{\mathbf{P}}(a) = v]}{e^{-T_a(q)}} \\ &= \frac{1}{\lambda - 1} \sum_{a \in \mathcal{A}} R_a \left(- \sum_{v \in \vartheta(a)} \mathbb{P}[\vartheta_{\mathbf{Q}}(a) = v] \log \mathbb{P}[\vartheta_{\mathbf{P}}(a) = v] \right). \end{aligned}$$

In fact, this shows that $\dim_{\text{loc}}(\mu_P, x) = \alpha$ for μ_Q -almost all $x \in X_\vartheta$ by [Proposition 4.5](#). Thus $f_{\mu_P}(\alpha) \geq \dim_{\text{H}} \mu_Q = \tau_{\mu_P}^*(\alpha)$, as required.

The result for $\alpha = \alpha_{\min}$ (resp. $\alpha = \alpha_{\max}$) follows similarly by taking a degenerate probability vector Q assigning equal value to the realisations of $\vartheta(a)$ with maximal (resp. minimal) probabilities given by P , and zero otherwise. The corresponding non-degenerate sub-substitution is also compatible and recognisable, so the same arguments yield the corresponding bounds. \square

5. EXAMPLES, COUNTEREXAMPLES AND APPLICATIONS

5.1. Failure of bounds for $q < 0$ without recognisability. In the following two examples, we show the results in [Theorem A](#) do not extend in general to give an upper bound for the L^q -spectrum in terms of the inflation word L^q -spectrum, for $q < 0$. In [Example 5.1](#), we construct a class of frequency measures on the full-shift on two letters for which the L^q -spectrum and inflation word analogue differ in the $q < 0$ case. The random substitutions that give rise to these frequency measures are not compatible, but in [Example 5.2](#) we present a compatible analogue.

In contrast, in [Example 5.3](#), we give an example showing that the results for $q < 0$ can hold for all $q \in \mathbb{R}$ under the identical set condition with identical production probabilities.

Example 5.1. Let $p_1 < p_2 \in (0, 1)$ such that $p_1 + 3p_2 = 1$ and let ϑ_P be the random substitution defined by

$$\vartheta_P: a, b \mapsto \begin{cases} ab & \text{with probability } p_1 \\ ba & \text{with probability } p_2 \\ aa & \text{with probability } p_2 \\ bb & \text{with probability } p_2 \end{cases}$$

We will show for all sufficiently small $q < 0$ that $\tau_{\mu_P}(q) > T_{\vartheta, P}(q)$. Observe that, for each $k \in \mathbb{N}$, the word $v^k = (ab)^{2^k} \in \vartheta^{k+1}(a) \cap \vartheta^{k+1}(b)$ occurs with probability

$$\mathbb{P}[\vartheta_P^{k+1}(a) = v^k] = \mathbb{P}[\vartheta_P^{k+1}(b) = v^k] = p_1^{2^k}.$$

Clearly, this is the minimal possible probability with which a level- k inflation word can occur, so it follows that

$$\lim_{q \rightarrow -\infty} \frac{T_{\vartheta, P}(q)}{q} = -\frac{1}{2} \log p_1.$$

Now, let $u \in \mathcal{L}_\vartheta^{2^{k+1}}$ be arbitrary. We show that $\mu_P([u]) \geq p_1^{2^{k-1}} p_2^{2^{k-1}}/2$. Since $\vartheta(a) = \vartheta(b)$ with identical production probabilities, it follows by [Lemma 2.12](#) that for any choice of $w \in \mathcal{L}_\vartheta^{2^{k+1}}$

$$\mu_P([u]) = \frac{1}{2} \left(\mathbb{P}[\vartheta_P(w)_{[1, 2^{k+1}]} = u] + [\vartheta_P(w)_{[2, 2^{k+1}+1]} = u] \right).$$

If $\mathbb{P}[\vartheta_{\mathbf{P}}(w)_{[1,2^{k+1}]} = u] \geq p_1^{2^{k-1}} p_2^{2^{k-1}}$, then we are done, otherwise at least half of the letters in v must be sent to ab . But then for u to appear from the second letter, at least half of the letters in v must be sent to ba or bb , so $\mathbb{P}[\vartheta_{\mathbf{P}}(w)_{[2,2^{k+1}+1]} = u] \geq p_1^{2^{k-1}} p_2^{2^{k-1}}$. Hence, $\mu_{\mathbf{P}}([u]) \geq p_1^{2^{k-1}} p_2^{2^{k-1}}/2$ so, in particular,

$$\min_{u \in \mathcal{L}_{\vartheta}^{2^{k+1}}} \mu_{\mathbf{P}}([u]) \geq \frac{1}{2} p_1^{2^{k-1}} p_2^{2^{k-1}}.$$

It follows that

$$\lim_{q \rightarrow -\infty} \frac{\tau_{\mu_{\mathbf{P}}}(q)}{q} \leq -\frac{1}{4}(\log p_1 + \log p_2) < -\frac{1}{2} \log p_1 = \lim_{q \rightarrow -\infty} \frac{T_{\vartheta, \mathbf{P}}(q)}{q}.$$

By a slight modification of this example, we can construct a compatible random substitution for which the two notions do not coincide.

Example 5.2. Let $p_1 < p_2 \in (0, 1)$ such that $p_1 + 3p_2 = 1$ and let $\vartheta_{\mathbf{P}}$ be the random substitution defined by

$$\vartheta_{\mathbf{P}}: a, b \mapsto \begin{cases} ab \, ba & \text{with probability } p_1 \\ ba \, ab & \text{with probability } p_2 \\ ab \, ab & \text{with probability } p_2 \\ ba \, ba & \text{with probability } p_2 \end{cases}$$

By similar arguments to the previous example, we obtain

$$\lim_{q \rightarrow -\infty} \frac{\tau_{\mu_{\mathbf{P}}}(q)}{q} \leq -\frac{1}{8}(\log p_1 + \log p_2) < -\frac{1}{4} \log p_1 = \lim_{q \rightarrow -\infty} \frac{T_{\vartheta, \mathbf{P}}(q)}{q}.$$

The random substitution in [Example 5.2](#) satisfies the identical set condition with identical production probabilities. These conditions are also satisfied by the following example. However, here the L^q -spectrum and inflation word analogue coincide for all $q \in \mathbb{R}$ by a direct argument.

Example 5.3. We will show that for the random substitution

$$\vartheta_{\mathbf{P}}: a, b \mapsto \begin{cases} ab & \text{with probability } p \\ ba & \text{with probability } 1 - p \end{cases}$$

the limit defining $\tau_{\mu_{\mathbf{P}}}(q)$ exists for all $q \in \mathbb{R}$, and

$$\tau_{\mu_{\mathbf{P}}}(q) = T_{\vartheta, \mathbf{P}}(q) = \frac{1}{\lambda} \varphi_1(q) = -\frac{1}{2} \log(p^q + (1-p)^q).$$

[Corollary B](#) gives the result for all $q > 0$ and that $\tau_{\mu_{\mathbf{P}}}(q) \geq T_{\vartheta, \mathbf{P}}(q) = 2^{-1} \varphi_1(q)$ for all $q < 0$, so it only remains to verify for all $q < 0$ that

$$\bar{\tau}_{\mu_{\mathbf{P}}}(q) \leq T_{\vartheta, \mathbf{P}}(q).$$

Since $\vartheta(v^1) = \vartheta(v^2)$ for all $v^1, v^2 \in \mathcal{L}_\vartheta$, it follows from [Lemma 2.12](#) that for all $u \in \mathcal{L}_\vartheta^{2m}$ and any $v \in \mathcal{L}_\vartheta^{m+1}$,

$$\mu_{\mathbf{P}}([u]) = \frac{1}{2} (\mathbb{P}[\vartheta(v)_{[1,1+2m-1]} = u] + \mathbb{P}[\vartheta(v)_{[2,2+2m-1]} = u]).$$

Let $\mathcal{V}_{2m} = \{(ab)^m, (ba)^m\}$. If $u \in \mathcal{L}_\vartheta^{2m} \setminus \mathcal{V}_{2m}$, then u must contain bb as a subword. This uniquely determines the cutting points in any inflation word decomposition, so there exists a unique v and $j(u) \in \{1, 2\}$ such that $u \in \vartheta(v)_{[j(u), 2m+j(u)-1]}$. It follows that

$$\begin{aligned} \sum_{u \in \mathcal{L}_\vartheta^{2m}} \mu_{\mathbf{P}}([u])^q &\geq \sum_{u \in \mathcal{L}_\vartheta^{2m} \setminus \mathcal{V}_{2m}} \left(\frac{1}{2} \mathbb{P}[\vartheta_{\mathbf{P}}(v)_{[j(u), j(u)+2m-1]} = u] \right)^q \\ &\geq \frac{1}{2^q} \sum_{u \in \mathcal{L}_\vartheta^{2m} \setminus \mathcal{V}_{2m}} \mathbb{P}[\vartheta_{\mathbf{P}}(v_2 \cdots v_m) = u_{[3-j(u), 2-j(u)+2m]}]^q. \end{aligned}$$

Now, observe that for every $w \in \vartheta(v_2 \cdots v_m)$ there is a u such that $w = u_{[3-j(u), 2-j(u)+2m]}$. Hence,

$$\sum_{u \in \mathcal{L}_\vartheta^{2m}} \mu_{\mathbf{P}}([u])^q \geq \frac{1}{2^q} \sum_{w \in \vartheta(v_2 \cdots v_m)} \mathbb{P}[\vartheta_{\mathbf{P}}(v_2 \cdots v_m) = w]^q$$

and the conclusion follows by similar arguments to those used in the proofs of the main theorems.

5.2. Examples with recognisability. We first provide examples of random substitutions for which the multifractal formalism holds.

Example 5.4. Let $p > 0$ and let ϑ_p be the random substitution defined by

$$\vartheta_p: \begin{cases} a \mapsto \begin{cases} abb & \text{with probability } p \\ bab & \text{with probability } 1-p \end{cases} \\ b \mapsto aa \end{cases}$$

Certainly ϑ_p is compatible, with corresponding primitive substitution matrix

$$M = \begin{pmatrix} 1 & 2 \\ 2 & 0 \end{pmatrix},$$

Perron–Frobenius eigenvalue $(1 + \sqrt{17})/2$, and (normalised) right Perron–Frobenius eigenvector

$$\left(\frac{-3 + \sqrt{17}}{2}, \frac{5 - \sqrt{17}}{2} \right).$$

One can verify that ϑ is recognisable since every occurrence of aa intersects an image of b and the adjacent letters then determine the cutting points. Thus by [Theorem D](#), for all $q \in \mathbb{R}$

$$\tau_{\mu_p}(q) = T_{\vartheta, \mathbf{P}}(q) = \frac{1}{\lambda - 1} \varphi_1(q) = -\frac{7 - \sqrt{17}}{8} \log(p^q + (1-p)^q)$$




FIGURE 1. L^q -spectra and multifractal spectra corresponding to a recognisable substitution for $p \in \{1/5, 2/5\}$.

and measure μ_p satisfies the multifractal formalism. The asymptotes have slopes $-(7 - \sqrt{17}) \log(p)/8$ and $-(7 - \sqrt{17}) \log(1 - p)/8$. A plot of the L^q -spectra and multifractal spectra for two choices of p is given in Figure 1.

For $p = 1/2$, the L^q -spectrum of the measure μ_p is a straight line and the multifractal spectrum is equal to $h_{\text{top}}(X_\vartheta)$ at $h_{\text{top}}(X_\vartheta)$, and $-\infty$ otherwise.

In the following example, we highlight that the multifractal spectrum need not have value 0 at the endpoints.

Example 5.5. Let ϑ_p be the random substitution defined by

$$\vartheta_p: \begin{cases} a \mapsto \begin{cases} abb & \text{with probability } p \\ bab & \text{with probability } p \\ bba & \text{with probability } 1 - 2p \end{cases} \\ b \mapsto aaa \end{cases}$$

Similarly to Example 5.4, ϑ_p is primitive, compatible and recognisable. Hence, Theorem D gives that

$$\tau_{\mu_p}(q) = -\frac{3}{10} \log(2p^q + (1 - 2p)^q).$$

The asymptotes have slopes $-3 \log(p)/10$ and $-3 \log(1 - 2p)/10$. For $p = 1/5$ and $p = 2/5$, the L^q -spectrum and multifractal spectrum of μ_p are plotted in Figure 2. Here, we highlight that the endpoints of the multifractal spectrum need not be equal to zero.




FIGURE 2. L^q -spectra and multifractal spectra corresponding to a recognisable substitution for $p \in \{1/5, 2/5\}$.

Example 5.6. Consider the random substitution on the three-letter alphabet $\mathcal{A} = \{a, b, c\}$ defined by

$$\vartheta_{\mathbf{P}}: \begin{cases} a \mapsto \begin{cases} bbc & \text{with probability } p_1 \\ cbb & \text{with probability } 1 - p_1 \end{cases} \\ b \mapsto \begin{cases} cca & \text{with probability } p_2 \\ acc & \text{with probability } 1 - p_2 \end{cases} \\ c \mapsto \begin{cases} aab & \text{with probability } p_3 \\ baa & \text{with probability } 1 - p_3 \end{cases} \end{cases}$$

for p_1, p_2 , and p_3 in $(0, 1)$. It is immediate that this substitution is compatible, and by considering the occurrences of 2, 3, or 4 letter repetitions, we observe that this substitution is also recognisable. Moreover, the hypotheses of [15, Theorem 4.8] are satisfied since ϑ is constant length and $\#\vartheta(a) = \#\vartheta(b) = \#\vartheta(c)$. In particular, the corresponding subshift X_{ϑ} is intrinsically ergodic with unique measure of maximal entropy given by taking $p_1 = p_2 = p_3 = 1/2$.

It follows from [15, Lemma 4.12] that the measure of maximal entropy is not a Gibbs measure with respect to the zero potential, so the system does not satisfy the usual specification property. For this choice of uniform probabilities, the L^q -spectrum is a straight line passing through the point $(1, 0)$ with slope $h_{\text{top}}(X_{\vartheta}) = \log(2)/2$. More generally, the L^q -spectrum is given for all $q \in \mathbb{R}$ by the formula

$$\tau_{\mu_{\mathbf{P}}}(q) = -\frac{1}{6} \left(\log((1 - p_1)^q + p_1^q) + \log((1 - p_2)^q + p_2^q) + \log((1 - p_3)^q + p_3^q) \right)$$

and the multifractal formalism is satisfied.

For an example on an alphabet of size two, one may consider the random substitution

$$\vartheta_{\mathbf{P}}: \begin{cases} a \mapsto \begin{cases} ababbb & \text{with probability } p_1 \\ abbabb & \text{with probability } 1 - p_1 \end{cases} \\ b \mapsto \begin{cases} baabaa & \text{with probability } p_2 \\ babaaa & \text{with probability } 1 - p_2 \end{cases} \end{cases}$$

for p_1 and p_2 in $(0, 1)$. The analysis of this example proceeds identically as above.

5.3. Examples without recognisability. Finally, we consider the two most commonly studied examples of random substitutions: random period doubling and random Fibonacci.

Example 5.7. Given $p \in (0, 1)$, let ϑ_p be the random period doubling substitution defined by

$$\vartheta_p: \begin{cases} a \mapsto \begin{cases} ab & \text{with probability } p \\ ba & \text{with probability } 1 - p \end{cases} \\ b \mapsto aa \end{cases}$$

and let μ_p denote the corresponding frequency measure. The substitution ϑ_p satisfies the disjoint set condition, so for all $q \in [0, \infty)$,

$$\tau_{\mu_p}(q) = -\frac{2}{3} \log(p^q + (1 - p)^q).$$

The asymptote as $q \rightarrow \infty$ has slope $-2 \log(\max\{p, 1 - p\})/3$, which gives a sharp lower bound on the local dimensions of μ_p .

If $p = 1/2$, then the measure μ_p has linear L^q -spectrum for $q \geq 0$ given by

$$\tau_{\mu_{1/2}}(q) = \frac{2}{3}(q - 1) \log 2.$$

Since the substitution satisfies the disjoint set condition but is not recognisable, our results do not give the L^q -spectrum for $q < 0$.

Example 5.8. The random Fibonacci substitution ϑ_p defined by

$$\vartheta_p: \begin{cases} a \mapsto \begin{cases} ab & \text{with probability } p \\ ba & \text{with probability } 1 - p \end{cases} \\ b \mapsto a \end{cases}$$

does not satisfy either the identical set condition nor the disjoint set condition. Hence, we cannot apply [Corollary B](#) to obtain a closed-form formula for $\tau_{\mu_p}(q)$. However, we can still apply [Theorem A](#) to obtain a sequence of lower and upper bounds. The case $k = 1$ gives the following bounds for all $0 < q < 1$:

$$-\frac{1}{\phi^2} \log(p^q + (1 - p)^q) = \frac{1}{\phi} \varphi_1(q) \leq \tau_{\mu_p}(q) \leq \frac{1}{\phi - 1} \varphi_1(q) = -\log(p^q + (1 - p)^q),$$




FIGURE 3. Upper and lower bounds on the L^q -spectrum of the frequency measure corresponding to the random Fibonacci substitution with $p = 1/2$, for $k = 3, 5, 7$.

where ϕ denotes the golden ratio. Reversing the inequalities yields the corresponding bounds for $q > 1$. Of course, by considering larger k we can obtain better bounds.

For $p = 1/2$, the bounds given by [Theorem A](#) for $k = 3, 5, 7$ are shown in [Figure 3](#).

ACKNOWLEDGEMENTS

The authors are grateful to Philipp Gohlke for his detailed comments on a draft version of this manuscript, which helped to remove some technical assumptions from [Theorem D](#). They also thank Dan Rust and Tony Samuel for valuable input. AM thanks SFB1283 and the Universität Bielefeld for supporting a research visit during the summer of 2022, where some of the work on this project was undertaken. AM was supported by EPSRC DTP and the University of Birmingham. AR was supported by EPSRC Grant EP/V520123/1 and the Natural Sciences and Engineering Research Council of Canada. The authors thank the organisers of the Junior Ergodic Theory Meeting hosted at the ICMS in Edinburgh in March 2022, where this project began.

REFERENCES

1. Matthias Arbeiter and Norbert Patzschke, *Random Self-Similar Multifractals*, Math. Nachr. **181** (1996), no. 1, 5–42. [3](#)
2. Michael Baake and Uwe Grimm, *Aperiodic order*, Encyclopedia of Mathematics and Its Applications, no. 149, Cambridge University Press, Cambridge ; New York, 2013. [2](#), [10](#)
3. Michael Baake, Timo Spindeler, and Nicolae Strungaru, *Diffraction of compatible random substitutions in one dimension*, Indag. Math. **29** (2018), no. 4, 1031–1071. [2](#), [5](#)
4. Rufus Bowen, *Topological entropy for noncompact sets*, Trans. Amer. Math. Soc. **184** (1973), 125–136. [11](#)

5. Vaughn Climenhaga and Daniel J. Thompson, *Beyond Bowen's Specification Property, Thermodynamic Formalism* (Mark Pollicott and Sandro Vaienti, eds.), vol. 2290, Springer International Publishing, Cham, 2021, pp. 3–82. [3](#)
6. Manfred Einsiedler, Elon Lindenstrauss, and Thomas Ward, *Entropy in Ergodic Theory and Topological Dynamics*, preprint. [11](#)
7. Kenneth J. Falconer, *Techniques in fractal geometry*, Wiley, Chichester ; New York, 1997. [3](#), [11](#)
8. Ai-Hua Fan, De-Jun Feng, and Jun Wu, *Recurrence, dimension, and entropy*, J. Lond. Math. Soc. **64** (2001), no. 1, 229–244. [3](#)
9. Ai-Hua Fan, Ka-Sing Lau, and Hui Rao, *Relationships between Different Dimensions of a Measure*, Monatsh. Math. **135** (2002), no. 3, 191–201. [13](#)
10. De-Jun Feng, *The limited Rademacher functions and Bernoulli convolutions associated with Pisot numbers*, Adv. Math. **195** (2005), no. 1, 24–101. [3](#)
11. De-Jun Feng and Ka-Sing Lau, *Multifractal formalism for self-similar measures with weak separation condition*, J. Math. Pures Appl. **92** (2009), no. 4, 407–428. [3](#)
12. Robbert Fokkink, Dan Rust, and Ville Salo, *Automorphism groups of random substitution subshifts*, (preprint). [5](#), [21](#)
13. Claude Godrèche and Jean-Marc Luck, *Quasiperiodicity and randomness in tilings of the plane*, J. Stat. Phys. **55** (1989), no. 1-2, 1–28. [2](#)
14. Philipp Gohlke, *Inflation word entropy for semi-compatible random substitutions*, Monatsh. Math. **192** (2020), no. 1, 93–110. [2](#), [5](#), [7](#), [16](#), [17](#), [33](#)
15. Philipp Gohlke, Andrew Mitchell, Dan Rust, and Tony Samuel, *Measure Theoretic Entropy of Random Substitution Subshifts*, Ann. Henri Poincaré (to appear). [3](#), [5](#), [7](#), [9](#), [16](#), [20](#), [21](#), [23](#), [33](#), [45](#)
16. Philipp Gohlke, Dan Rust, and Timo Spindeler, *Shifts of finite type and random substitutions*, Discrete Contin. Dyn. Syst. **39** (2019), no. 9, 5085–5103. [17](#)
17. Philipp Gohlke and Timo Spindeler, *Ergodic frequency measures for random substitutions*, Studia Math. **255** (2020), no. 3, 265–301. [3](#), [17](#), [18](#), [20](#)
18. Thomas C. Halsey, Mogens H. Jensen, Leo P. Kadanoff, Itamar Procaccia, and Boris I. Shraiman, *Fractal measures and their singularities: The characterization of strange sets*, Phys. Rev. A **33** (1986), no. 2, 1141–1151. [3](#), [4](#)
19. Ka-Sing Lau and Sze-Man Ngai, *Multifractal Measures and a Weak Separation Condition*, Adv. Math. **141** (1999), no. 1, 45–96. [3](#), [4](#), [14](#)
20. Douglas Lind and Brian Marcus, *An Introduction to Symbolic Dynamics and Coding*, first ed., Cambridge University Press, November 1995. [10](#)
21. Eden Miro, Dan Rust, Lorenzo Sadun, and Gwendolyn S. Tadeo, *Topological Mixing of Random Substitutions*, (preprint). [5](#)
22. Lars Olsen, *A Multifractal Formalism*, Adv. Math. **116** (1995), no. 1, 82–196. [3](#)
23. Yakov B. Pesin, *Dimension theory in dynamical systems: Contemporary views and applications*, Chicago Lectures in Mathematics Series, University of Chicago Press, Chicago, 1997. [3](#)
24. Mark Pollicott and Howard Weiss, *Multifractal Analysis of Lyapunov Exponent for Continued Fraction and Manneville-Pomeau Transformations and Applications to Diophantine Approximation*, Comm. Math. Phys. **207** (1999), no. 1, 145–171. [3](#)
25. Martine Queffélec, *Substitution Dynamical Systems-Spectral Analysis*, Lecture Notes in Mathematics, no. 1294, Springer Berlin Heidelberg, Berlin, Heidelberg, 1987. [2](#), [19](#), [20](#)
26. R. Tyrrell Rockafellar, *Convex analysis*, Princeton Mathematical Series, no. 28, Princeton University Press, Princeton, N.J., 1970. [14](#)
27. Dan Rust, *Periodic points in random substitution subshifts*, Monatsh. Math. **193** (2020), no. 3, 683–704. [5](#), [17](#)
28. Dan Rust and Timo Spindeler, *Dynamical systems arising from random substitutions*, Indag. Math. **29** (2018), no. 4, 1131–1155. [17](#)

- 29. Pablo Shmerkin, *On Furstenberg's intersection conjecture, self-similar measures, and the L^q norms of convolutions*, Ann. Math. **189** (2019), no. 2, 319. [3](#), [9](#), [13](#)
- 30. Péter P. Varjú, *Recent progress on Bernoulli convolutions*, Proceedings of the 7th European Congress of Mathematics (Berlin), January 2018. [3](#)

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, B15 2TT

Email address: `acm925@student.bham.ac.uk`

MATHEMATICAL INSTITUTE, UNIVERSITY OF ST ANDREWS, ST ANDREWS, KY16 9SS

Email address: `alex@rutar.org`