

A review of abnormal prolactin response in ME/CFS

forestglip*

September 19, 2026

Abstract

Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a disabling health condition, characterized by post-exertional malaise and accompanied by symptoms such as fatigue, pain, brain fog, and orthostatic intolerance. Affecting about 1 in 200 people, ME/CFS has no validated treatments and very few insights into its pathophysiology. Case-control studies of ME/CFS, or the related condition, postviral fatigue syndrome, have consistently demonstrated that patients exhibit an abnormally large increase in prolactin following administration of buspirone, a drug which acts at serotonin and dopamine receptors. Less replicated evidence of abnormal prolactin response comes from studies that tested d-fenfluramine, exercise, and insulin-induced hypoglycemia. The implications of these findings are not yet clear, though potential explanations include alterations in neurotransmitter systems or sex hormone pathways. This review discusses what is known about prolactin response in ME/CFS and suggests future research directions.

Introduction

ME/CFS

Around 0.5-1% of the population suffers from myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) [1], a debilitating condition characterized by symptoms such as post-exertional malaise (PEM), fatigue, pain, brain fog, and orthostatic intolerance. PEM is defined as the exacerbation or new onset of symptoms following minor physical, cognitive, emotional, or social activities, and which can persist for days or weeks, or even lead to longer term worsening. In severe cases of ME/CFS, individuals may be bedbound and require a dark and silent living environment to minimize symptoms and prevent worsening. There is no known cure for this condition [2].

Because ME/CFS is a heterogenous syndrome defined by symptoms, specific criteria for ME/CFS vary, with the name "ME/CFS" itself an umbrella term that encompasses the two other most common names for this syndrome: "chronic fatigue syndrome" and "myalgic encephalomyelitis" [3]. In contrast to older ME/CFS criteria, such as those proposed by Holmes et al. [4], Sharpe et al. [5], or Fukuda et al. [6], more recent case criteria, such as the 2003 Canadian consensus criteria [7] or 2015 Institute of Medicine criteria [3] require the presence of PEM for diagnosis.

As yet, no biomarkers have been validated for ME/CFS and little is known about its pathophysiology [8]. Some of the few relatively replicated abnormalities in ME/CFS, which, if proven to be robust, may eventually provide direction for treatment development, include: higher brain lactate [9–15], increased usage of the IGHV3-30 gene by B cells [16–18], and reduced natural killer cell cytotoxicity [19].

While all of the findings noted above continue to be investigated in recent years, a further ME/CFS finding, namely abnormal prolactin response to neuroendocrine challenge, was observed in numerous studies, but has not been directly tested further in over 15 years. This review aims to outline what is known about prolactin response in ME/CFS, and to suggest research directions which may provide opportunities for deeper understanding of the mechanisms underlying this condition.

Neuroendocrine challenge testing

Studies of many health conditions have been undertaken to investigate the functioning of the nervous system through the use of neuroendocrine challenge tests. Such tests are performed by measuring the release of pituitary hormones after administration of drugs or other interventions known to affect hormonal release. As the pituitary gland is regulated, in part, by the brain, this testing strategy can be considered a relatively non-invasive "window" into central nervous system functioning [20].

In support of the potential for these tests to detect altered neural functioning, researchers have demonstrated changes in prolactin response following neurotoxic exposure. For example, studies have shown changes in the prolactin response to serotonin-related drugs following damage to serotonin neurons in monkeys [21] and rats [22, 23].

Prolactin

Prolactin is a protein produced within several tissues in the human body, but its main site of synthesis is the anterior pituitary gland. This hormone is best known for its essential

*Independent researcher. Correspondence to forestglip@tuta.io.

role in the development of mammary tissue and its ability to stimulate lactation. Additionally, prolactin exerts numerous other effects, which include the promotion of parental behavior, enhancement of immunity, modulation of lipid and glucose metabolism, and inhibition of the reproductive cycle during lactation. The most potent natural stimulant of prolactin secretion is suckling, however stress, sounds, and circadian light patterns also affect the release of this hormone [24, 25].

Unlike most hormones of the pituitary gland, prolactin is primarily regulated through inhibition instead of stimulation, whereby pituitary prolactin-secreting cells (lactotrophs) naturally secrete prolactin when hypothalamic influence is blocked. The main hormone involved in inhibiting prolactin is dopamine, which is released by neurons of the arcuate nucleus and periventricular nucleus of the hypothalamus, and which travels through the long or short portal veins to reach the lactotrophs. Additionally, several hormones can directly stimulate prolactin release by lactotrophs, and these include, but are not limited to, thyrotropin-releasing hormone (TRH), vasopressin, oxytocin, and vasoactive intestinal peptide (VIP) [25].

Prolactin response in ME/CFS

Buspirone

Strikingly, every single published study that has tested prolactin response to buspirone in ME/CFS has observed that patients exhibit an abnormally large increase in prolactin, which occurs about one to two hours after drug administration. The results of these studies are summarized below, as well as in Table 1.

Buspirone is marketed as a treatment for generalized anxiety disorder, though, unlike classic anxiety medications such as benzodiazepines, buspirone has no activity at GABA receptors. Instead, buspirone is an agonist of 5-HT_{1A} receptors, where it exhibits strong binding affinity, and also acts as an antagonist at dopamine D₂ receptors, albeit with weaker binding affinity [26].

The first study demonstrating increased prolactin response to buspirone in an ME/CFS-like condition was undertaken by Bakheit et al. in 1992. The findings from this study were reported both in Bakheit's thesis [27], as well as in a peer-reviewed paper [28]. The authors were studying postviral fatigue syndrome (PVFS), defined using criteria which closely resembled CDC [4] and Oxford [5] criteria for chronic fatigue syndrome. The study also required that patients' fatigue was preceded by a probable or definite viral infection, and was made worse by exercise and did not improve after bed rest. The authors analyzed males and females separately, finding, in both cases, that when compared to healthy controls or individuals with depression, those with PVFS exhibit a larger increase in plasma prolactin after administration of buspirone. Figure 1 illustrates the study results using the individual-level data provided in Bakheit's thesis. No-

tably, and as seen in virtually all other studies of this topic, baseline prolactin was not significantly altered in the patient group.

Further evidence of increased prolactin response to buspirone was provided by John Richardson in 1995. In this study, patients who fulfilled CDC and Oxford criteria for ME/CFS were compared to family members without ME/CFS. With prolactin response defined as the ratio of prolactin levels after to before buspirone administration, the response to buspirone (50 mg, orally) in the patients was, on average, over 3 times higher than that of the controls. The study also found a correlation between prolactin response in patients and their degree of sleep disturbance [29]. However, the sexes and ages of control participants were not described, raising the possibility of confounding related to these characteristics. A later study from Richardson and Costa again suggested very high prolactin responses to buspirone (50 mg, orally) in ME/CFS patients, however this study did not include a control group. The later paper also suggests possible selection bias, in that high prolactin response may have been part of the study's inclusion criteria [30].

In 1996, Sharpe et al. reported increased prolactin response to buspirone (0.5 mg/kg up to 45 mg, orally) in patients fulfilling Oxford criteria, and who did not also exhibit symptoms of depression. Notably, plasma levels of buspirone and its metabolite, 1-(2-Pyrimidinyl)piperazine (1-PP), were not significantly different between groups after administration of buspirone, suggesting that the difference in prolactin response was not caused by differences in drug metabolism. In contrast to prolactin response, the growth hormone response to buspirone did not significantly differ between groups [31].

Also in 1996, Tahir Majeed, under the supervision of Peter O. Behan (also an author on the 1992 PVFS paper), published a thesis testing hormonal responses to a variety of drug challenges in ME/CFS, providing further evidence of abnormal prolactin response. The study cohort included patients who fulfilled Oxford and Fukuda [6] criteria, and who had fatigue which worsened after exercise. The control group was made up of healthy, sex-matched volunteers. The ME/CFS patients exhibited a significantly larger increase in prolactin after administration of buspirone (60 mg, orally). Further findings in ME/CFS patients included normal growth hormone responses to bromocriptine and baclofen, a larger growth hormone response to pyridostigmine, smaller growth hormone responses to desipramine and dexamethasone, and a smaller adrenocorticotrophic hormone (ACTH) response to ipsapirone [32].

Another 1996 paper, authored by Behan, reported increased prolactin response to buspirone (60 mg, orally) in individuals with an ME/CFS-like illness (including myalgia, fatigue worsened by exercise, and with onset after a flu-like illness) and who had previously been exposed to organophosphates [33]. Note that while this paper describes the control group as 10 healthy males, later court testimony from Behan in an organophosphate trial indicated that there was a




Figure 1: Prolactin response to buspirone in PVFS, depression, and healthy controls. Plot created from data in A. M. O. Bakheit's thesis [27]. PVFS: Postviral fatigue syndrome.

mistake in the paper, and the actual control group was made up of 15 males and 15 females [34], with the control group data matching that of Majeed's 1996 thesis [32]. However, as Behan noted, the inclusion of females in the control group would not be expected to lead to a spuriously increased response in the patient group, as females have larger prolactin responses to buspirone than males [35].

Additionally, two studies from Racciatti et al. suggest that the authors have observed further evidence of increased prolactin response to buspirone in ME/CFS. However, one study did not include a healthy control group, and the study manuscript reported very minimal detail about the prolactin response finding [36]. The other study was reported in an abstract which we have not been able to access [37], and with the findings briefly mentioned in a later review [38].

Finally, a 2001 case report from Sharma et al. described a female patient with ME/CFS (Oxford criteria), who exhibited an abnormally large prolactin response to buspirone (30 mg, orally). After symptom improvement, which followed a graded exercise therapy intervention, the prolactin response was measured again on two occasions, and was found to be normal both times [39].

Non-buspirone probes

Additional evidence of abnormal prolactin response was provided by studies that tested fenfluramine, exercise, and insulin-induced hypoglycemia in ME/CFS patients. Negative or conflicting results have been observed after administra-

tion of m-chlorophenylpiperazine (m-CPP) and tryptophan. Details of these studies are summarized below and in Table 2.

Several studies of prolactin response to fenfluramine in ME/CFS have been published, though with varying results. Fenfluramine is a serotonin releasing agent, and, through its metabolites, agonizes dopamine and serotonin receptors [42], though the d-enantiomer formulation of fenfluramine selectively affects serotonin receptors [43]. One study tested prolactin response to 60 mg of racemic fenfluramine, with no significant difference being observed between the ME/CFS and healthy groups [44]. Three studies have also tested response to 30 mg of d-fenfluramine. In two of these studies, ME/CFS patients had an abnormally increased prolactin response [45, 46], while the third did not detect a significant difference between groups [47].

Studies have also reported significantly *blunted* prolactin responses after neuroendocrine challenge with exercise or hypoglycemia. Bearn et al., in the same study that did not find a difference after d-fenfluramine challenge, did detect a blunted prolactin response to insulin-induced hypoglycemia in ME/CFS patients [47]. Further, Ottenweller et al. reported a significantly smaller increase of prolactin in ME/CFS following a treadmill-based maximal exercise test [48].

Two studies of other probes produced negative or inconsistent findings. A 2001 study did not find a significant difference between ME/CFS patients and controls in prolactin response to the serotonin agonist, m-CPP (0.25 mg/kg) [49]. Most recently, a 2010 study reported a significantly increased prolactin response to the serotonin precursor, tryptophan, in

Table 1: Studies testing prolactin response to buspirone in ME/CFS-like conditions

Study	Participants	Probe	Prolactin response [*]	Notes
Bakheit, 1992 [27] (Thesis, supervised by Peter O. Behan) Bakheit et al., 1992 [28]	6F, 9M PVFS 6F, 7M depression 6F, 7M healthy	Buspirone (60 mg, orally)	PVFS: 8.2 (females) and 5.0 (males) Depression: 3.6 (females) and 2.1 (males) Healthy: 3.9 (females) and 2.5 (males)	Females tested during luteal phase. Only PVFS participants experienced excessive fatigue, lightheadedness, and nausea after buspirone. Minor data inconsistencies for some of the participants, likely due to transcription errors, have been noted [40].
Richardson, 1995 [29]	25F, 5M CFS/ME (Oxford) 30 healthy (sex ratio not reported)	Buspirone (50 mg, orally)	CFS/ME: 4.9 Healthy: 1.8	Females tested during luteal phase. Patients experienced more nausea after buspirone compared to controls.
Sharpe et al., 1996 [31]	11M CFS (Oxford) 11M healthy	Buspirone (0.5 mg/kg up to 45 mg, orally)	CFS: 3.4 Healthy: 2.6	Growth hormone response was not significantly different between groups. Patients had more nausea in response to buspirone, but this did not correlate with prolactin response. Plasma levels of buspirone and its major metabolite, 1-PP, were not significantly different between groups.
Majeed, 1996 [32] (Thesis, supervised by Peter O. Behan)	15F, 15M CFS (Fukuda) 15F, 15M healthy	Buspirone (60 mg, orally)	CFS: 2.9 Healthy: 2.0	Females tested during follicular phase. All participants had been medication-free for at least 3 months. No participants reported nausea after buspirone. Other studies within the thesis found: normal GH responses to bromocriptine and baclofen, larger GH response to pyridostigmine, smaller GH responses to desipramine and dexamethasone, and smaller ACTH response to ipsapirone.
Behan, 1996 [33]	10M CFS with previous exposure to organophosphates. 15F, 15M healthy from previous study [32]	Buspirone (60 mg, orally)	CFS: 4.9 Healthy: 2.0	The paper mistakenly described the control group as 10 males [34]. Data for controls instead matches that of Majeed's 1996 thesis [32]. Patients also had increased GH response to pyridostigmine and decreased GH response to dexamethasone.
Racciatti et al., 2001 [36]	3F, 2M CFS after toxic exposure 3F, 1M CFS after EBV 4F, 1M CFS with depression No healthy controls	Buspirone	"an abnormal increase of prolactine levels followed the buspirone challenge test [...] in all the three subgroups of patients."	Prolactin values and buspirone dosage not reported.
Sharma et al., 2001 [39]	1F CFS (Oxford)	Buspirone (30 mg, orally)	CFS: 10.7 After symptom improvement following graded exercise program: 12 months: 2.0, 18 months: 1.2	Case study.

^{*} Prolactin response was calculated as average prolactin at the timepoint when the group average was highest divided by average prolactin at timepoint 0. Data were approximated from plots using WebPlotDigitizer 5.2 [41] when exact values were not reported.

Bold text indicates that the referenced paper reported a significant difference between the highlighted group and the healthy control group for prolactin response. Note that the definition of prolactin response for the purpose of significance testing varied by study, and was not the same as the definition used in this table.

To avoid including a biased value, Richardson & Costa's study [30] was excluded from this table due to their paper suggesting that abnormal prolactin response was part of the inclusion criteria.

1-PP: 1-(2-pyrimidinyl)piperazine; CFS: Chronic fatigue syndrome; EBV: Epstein-Barr Virus; F: Females; Fukuda: Fukuda et al., 1994 [6]; GH: Growth hormone; Holmes: Holmes et al., 1988 [4]; M: Males; ME: Myalgic encephalomyelitis; Oxford: Sharpe et al., 1991 [5]; PVFS: Postviral fatigue syndrome;

Table 2: Studies testing prolactin response to non-buspirone probes in ME/CFS-like conditions

Study	Participants	Probe	Prolactin response*	Notes
Cleare et al., 1995 [46]	4F, 6M CFS (Holmes and Oxford) 4F, 6M depression 8F, 12M healthy	D-fenfluramine (30 mg, orally)	CFS: 1.5 Depression: 0.9 Healthy: 1.2	Females tested during follicular phase. Cortisol response did not differ from healthy group. When analyzing full cohort, baseline cortisol inversely correlated with prolactin response.
Bearn et al., 1995 [47]	4F, 5M CFS (Oxford and all but one fulfilled Holmes) 6F, 4M healthy	D-fenfluramine (30 mg, orally)	CFS: 1.4 Healthy: 1.7	Females tested on days 3-5 of their menstrual cycle. ACTH response was greater in patients. Cortisol response showed no group difference.
Bearn et al., 1995 [47]	4F, 5M CFS (Oxford and all but one fulfilled Holmes) 6F, 2M healthy	Insulin (0.1 or 0.15 u/kg)	CFS: 5.1 Healthy: 6.8	Females tested on days 3-5 of their menstrual cycle. Growth hormone response trended lower in patients. Cortisol and ACTH responses were not significantly different.
Yatham et al., 1995 [44]	8F, 3M CFS (Holmes) 8F, 3M healthy	DL-fenfluramine (60 mg, orally)	CFS: 1.5 Healthy: 2.4	Cortisol response did not significantly differ between groups.
Sharpe et al., 1997 [45]	10M CFS + neurasthenia 10M healthy	D-fenfluramine (30 mg, orally)	CFS: 1.6 Healthy: 1.1	
Ottenweller et al., 2001 [48]	17F CFS (Holmes and Fukuda) 14F healthy	Maximal exercise test on treadmill	CFS: 1.2 Healthy: 2.2	Females tested during luteal phase. ACTH, EPI, TSH, DHPG, DOPA, and DOPAC responses were lower, GH response was higher, and NE response not significantly different.
Vassallo et al., 2001 [49]	16F, 4M neurasthenia (all but one also fulfilled Oxford and CDC criteria for CFS) 15F, 5M healthy	m-CPP (0.25 mg/kg, orally)	Ratio could not be calculated because average prolactin values were not reported. No significant difference between groups.	Females tested during early follicular phase. Participants also underwent a placebo test which did not raise prolactin in either group.
Weaver et al., 2010 [50]	15F, 7M CFS (Fukuda) 8F, 3M CFS (Fukuda) + FM 10F, 6M healthy	L-tryptophan (120 mg/kg lean body mass)	CFS: 2.3 (females) and 2.2 (males) CFS+FM: 2.0 (females) and 3.0 (males) Healthy: 2.0 (females) and 2.2 (males)	Females tested during late follicular phase (days 7–14).

* Prolactin response was calculated as average prolactin at the timepoint when the group average was highest divided by average prolactin at timepoint 0. Data were approximated from plots using WebPlotDigitizer 5.2 [41] when exact values were not reported.

Bold text indicates that the referenced paper reported a significant difference between the highlighted group and the healthy control group for prolactin response. Note that the definition of prolactin response for the purpose of significance testing varied by study, and was not the same as the definition used in this table.

ACTH: Adrenocorticotrophic hormone; CFS: Chronic fatigue syndrome; EPI: Epinephrine; FM: Fibromyalgia; Fukuda: Fukuda et al., 1994 [6]; F: Females; Holmes: Holmes et al., 1988 [4]; GH: Growth hormone; m-CPP: m-chlorophenylpiperazine; M: Males; NE: Norepinephrine; Oxford: Sharpe et al., 1991 [5]; TSH: Thyroid-stimulating hormone;

females with ME/CFS. However, this difference was not detected in females suffering from both ME/CFS and fibromyalgia, or in either of the male patient groups [50].

Prolactin response in non-ME/CFS conditions

Abnormally increased prolactin response has been reported in several other conditions, such as migraine, irritable bowel syndrome (IBS), and gonadal disorders. The following section briefly outlines some examples of non-ME/CFS prolactin response research, but this should not be considered a comprehensive review of the literature.

Migraine

Abnormal prolactin response has been reported several times in migraine. Cassidy et al. have published two studies showing increased prolactin response to buspirone in migraine [51, 52]. The prolactin response to m-CPP was larger in a study that used a dosage of 0.5 mg/kg [53], but not in a study that used a smaller 0.25 mg/kg dosage of the drug [54]. A larger prolactin response was noted after administration of racemic fenfluramine [55]. Migraine patients may have an abnormally long return to baseline prolactin after administration of reserpine (detected only in males) [56] or benserazide (detected only in females) [57]. Studies have reported abnormally increased prolactin response in migraine patients to the D2 antagonists sulpiride [57, 58] and domperidone [58]. An abnormally large *decrease* in prolactin has been noted after administration of L-deprenyl [59]. In contrast, an abnormally *small* decrease was reported after administration of the dopamine reuptake inhibitor, nomifensine [58] and the dopamine precursor, L-DOPA [60].

Several studies have tested TRH challenges in migraine. A blunted prolactin response in migraine was seen after administration of 50 μ g and 200 μ g doses of TRH [61]. Another study found an increased prolactin response to TRH during migraine attacks [62]. Awaki et al. found an abnormally increased prolactin response after administration of a cocktail which included TRH, gonadotropin-releasing hormone (GnRH), and insulin [63].

Other studies of migraine did not find significant differences between groups, including after administration of domperidone [64], fenfluramine [57], lisuride [58], metoclopramide [65, 66], morphine [67], piribedil [68], and sumatriptan [69].

Gonadal disorders

Studies have also tested prolactin response in disorders related to testicular and ovarian function, with increased response being a commonly reported finding. Abnormalities in levels of hormones other than prolactin are also commonly reported in these studies.

For females with amenorrhea secondary to ovarian failure, abnormally large prolactin response to TRH has been reported both in a case report [70] and in a case-control study. In the latter study, the patients also had increased basal luteinizing hormone (LH) and follicle-stimulating hormone (FSH), decreased estrone and estradiol, and normal TSH response to TRH [71].

Similar observations of increased prolactin response have been reported in relation to testicular disorders, though findings have not been entirely consistent. In several studies from Spitz et al., males with oligo- or azoospermia were found to have larger prolactin responses to TRH [72–75] or to combined TRH and GnRH [76]. Baranowska et al. reported increased prolactin response to metoclopramide in oligo- or azoospermia [77]. Boucher et al. found increased prolactin response to combined TRH and GnRH in males with varicocele [78]. Studies have also reported blunted prolactin response to TRH in azoospermia [79], as well as normal prolactin response to TRH in oligospermia [80]. The latter study additionally did not detect an abnormal prolactin response to L-DOPA, pyridoxine, or metoclopramide [80]. A study from Winters et al. reported that early pubertal males with hypogonadotropic hypogonadism had smaller prolactin responses to the D2 antagonist chlorpromazine [81].

Other health conditions

Studies from Dinan et al. have reported abnormally increased prolactin response to buspirone in IBS [82] and in non-ulcer dyspepsia [83–86].

In patients with major depressive disorder, the prolactin response to various probes has been observed to be blunted in some [46, 87–94], but not all studies [28, 95–103]. Additionally, some studies of depression have reported increased prolactin response [104, 105].

Connection to sex hormones?

Females are more likely than males to be diagnosed with ME/CFS [1], which raises an intriguing possibility that the well-known influence of sex hormones on prolactin regulation, briefly discussed below, may relate to the abnormal prolactin responses observed in ME/CFS.

The clearest connection of prolactin response findings to the sex bias in ME/CFS comes from observations that females tend to have larger prolactin responses to various neuroendocrine probes than males. For example, healthy females have a larger prolactin response to buspirone [28, 35]. Similar sex differences in prolactin response magnitude have been observed after administration of d-fenfluramine [106], thyrotropin-releasing hormone (TRH) [107], phenothiazine [108], and possibly chlorpromazine [109].

Evidence of sex hormone influence on prolactin regulation is additionally provided by observations that the magnitude of prolactin response to various drugs differs throughout the menstrual cycle. O’Keane et al. observed that while

baseline prolactin levels in healthy females stayed relatively steady throughout the menstrual cycle, prolactin response to d-fenfluramine was highest at mid-cycle, followed by the luteal, then follicular phase, which mirrored the differing amounts of circulating estradiol measured at different phases [110]. Prolactin response to buspirone in females was found to be larger during the luteal phase than during the follicular phase or at mid-cycle [35]. Prolactin response to TRH may be somewhat larger during the follicular phase [111], though findings are inconsistent [112].

Direct evidence of sex hormone influence on prolactin secretion is provided by studies demonstrating the effect of exogenous estradiol on regulation of prolactin. TRH-induced prolactin response was seen to be substantially larger in females taking a short course of exogenous estradiol, as well as after subsequent short-term combined contraceptive use, but not in those taking long-term combined contraceptives [111]. In male rats, a prolactin response to TRH was only detectable when the rats had previously been administered estradiol [113]. Exogenous estradiol causes an increased prolactin response to ghrelin in postmenopausal women [114]. Administration of estradiol to ovariectomized rats resulted in substantially larger prolactin responses to the dopamine antagonist, thioproperazine [115].

Another interesting potential connection to sex hormones is based on Dinan et al. showing that healthy females had more symptoms of sedation in response to buspirone during the luteal phase than at other timepoints, with this also being the timepoint where prolactin response was found to be highest [35]. This may relate to observations that ME/CFS patients report more buspirone-induced side effects, such as fatigue and nausea, when compared to healthy controls [28, 29, 31].

Additionally, while increased prolactin response is a somewhat rarely observed finding in other health conditions, some of the few conditions in which this has been reported multiple times also suggest connections to sex hormones. As outlined in the preceding section, migraine patients were observed in several studies to have increased prolactin response to various neuroendocrine probes [51–53, 55, 57, 58]. Like ME/CFS [1], migraine appears to also be more prevalent in females than males [116, 117]. Several studies have also reported increased prolactin response in gonadal disorders such as primary testicular failure and primary ovarian failure [71–74, 76–78]. Gonadal organs are the primary sites of sex hormone synthesis, and as such, the authors of several of the gonadal disorder studies speculate that alterations in sex hormone pathways may mediate the changes in prolactin response seen in these conditions [71, 75, 77].

Importantly, a hypothesis which ties estradiol to the abnormal prolactin response seen in ME/CFS would need to involve a mechanism which is independent of blood estradiol levels, as studies have shown that blood levels of this hormone do not appear to be significantly increased in ME/CFS [118–121]. One alternative possibility was proposed in a study of increased prolactin response to TRH in primary

ovarian failure, where the authors speculated that the patients might have increased conversion of androgens to estrogens within the hypothalamus [71], based on an observation that such increased conversion was seen in castrated rabbits [122]. Studies have also shown that brain estrogen levels rise in response to various forms of brain damage [123], raising the possibility that some form of injury to brain tissue may underlie prolactin response findings in ME/CFS.

Future directions

The implications of abnormal prolactin response in ME/CFS are far from clear. While several of the ME/CFS prolactin response papers suggested that the findings were evidence of abnormalities in the serotonin system [28, 32, 45, 46], there is still too little direct evidence to make strong conclusions about specific pathways. Interestingly, a PET study of ME/CFS patients, which was undertaken to test a hypothesis of serotonin receptor upregulation, based partly on prolactin response findings, found the opposite of what was expected (the authors found reduced 5-HT1A binding) [124]. Serotonin-related drugs, which were used in most of the ME/CFS studies with significant results, are unlikely to be able to directly stimulate lactotrophs [125], and may instead exert at least part of their influence on lactotrophs through altering hypothalamic dopamine release [126]. Thus, while abnormal prolactin response could be explained by changes in the receptors directly affected by the neuroendocrine probes, such as 5-HT receptors, these findings could instead be highlighting any of a number of possible downstream abnormalities, such as alterations in dopamine functioning or sensitization of lactotrophs.

A straightforward next step would be testing prolactin response to other neuroendocrine probes. For example, if dopaminergic drugs and TRH both cause abnormal prolactin responses, this might suggest general changes in lactotroph sensitivity, as dopamine and TRH both directly affect lactotrophs [25]. On the other hand, if neither of these probes demonstrates altered responses, but responses to serotonin-related drugs are altered, this may indicate upstream abnormalities. It may be most fruitful to consider testing the probes which have already been reported to cause increased prolactin response in other health conditions. For example, in one study of migraine, the D2 receptor antagonist domperidone caused an increased prolactin response [58]. Domperidone does not appreciably cross the blood-brain barrier [127], allowing researchers to test direct lactotroph stimulation while limiting confounding due to central nervous system effects.

A finding related to levels of the prolactin regulating alpha-melanocyte-stimulating hormone (α -MSH) may also be valuable to replicate. One ME/CFS study has reported increased levels of α -MSH, albeit with substantial overlap between groups [128]. α -MSH is known to alter prolactin response, both in vitro [129, 130] and in vivo in non-human animals [131, 132]. To explore a possible connection, re-

searchers may wish to test for a correlation between α -MSH level and prolactin response in participants with ME/CFS.

Researchers may also wish to explore a potential connection to sex hormones, as discussed in the previous section. As one possible avenue, a study could test the hypothesis that menstrual phase, menopause status, or hormonal contraceptive use play a role in the magnitude of the difference between ME/CFS and controls. Research could also involve measuring the prolactin response to buspirone or another probe in ME/CFS patients and healthy controls both before and after administration of an estrogen receptor antagonist or aromatase inhibitor. If the difference between groups were found to be diminished after blocking estrogen pathways, this would provide evidence of sex hormone involvement in the abnormal prolactin response. It may be necessary to prescribe an extended course of these drugs prior to the second prolactin challenge, as estrogens are capable of inducing long term changes in prolactin-secreting cells, for example, through genomic regulation or by increasing the number of lactotrophs, as reviewed by Grattan et al. [24].

It may also be possible to explore abnormal prolactin regulation further through the use of PET scans, for example by measuring dopamine receptor D2 (DRD2) binding in the pituitary gland. If the ME/CFS abnormality does relate to sex hormones, then DRD2 levels may be altered, as is seen when estradiol is applied to anterior pituitary cells in vitro [133].

Controlling for confounders

To ensure interpretable results, and to increase the likelihood of detecting real differences, researchers should be mindful to control for relevant potential confounders when studying prolactin response in ME/CFS.

One of the most important factors to consider in studies of prolactin response is the sex of participants. As noted earlier, females demonstrate larger prolactin responses than males [28, 35, 106–109]. Thus, at minimum, groups in case-control studies should consist of the same ratio of females to males. However, it may be wise to also control for sex in an appropriate statistical model or analyze males and females separately to reduce the influence of variance associated with sex.

Additionally, as females exhibit variance in prolactin response throughout the menstrual cycle [35, 110], researchers should control for menstrual phase. Some of the clearest differences between ME/CFS and healthy control groups occurred in studies of females tested during the luteal phase [28, 29], while a study that included females tested during the follicular phase showed a smaller group difference [32]. While one can not make strong conclusions about menstrual phase effects from such limited data, this may suggest that the optimal time for testing females would be during the luteal phase. In any case, studies of females should ensure that controls are matched for menstrual phase.

Limited evidence suggests that sedentariness may also lead to altered prolactin response. Non-endurance trained

individuals, when compared to elite endurance-trained athletes, may exhibit an increased prolactin response to buspirone [134]. Cadegiani & Kater observed that non-athletes have smaller prolactin responses to insulin-induced hypoglycemia than athletes [135], which is directionally consistent with the results of an insulin study of ME/CFS [47]. Brooks et al. observed larger prolactin response to m-CPP in non-exercising subjects than in marathon runners [136], though in ME/CFS, a difference was not observed when testing a smaller dose of m-CPP [49]. In contrast, a study found no difference between endurance-trained and sedentary males during a d-fenfluramine prolactin challenge [137], and another study found that 9 weeks of exercise did not alter prolactin response to buspirone [138]. To explore the possibility that abnormal prolactin response findings in ME/CFS may be caused by deconditioning, researchers could include a control group matched for physical activity level, such as patients who are bed-bound for reasons unrelated to ME/CFS.

It may also be worthwhile to control for the effects of additional placebo-like factors on prolactin response [139]. Stress can cause changes in prolactin secretion [140, 141], and thus could potentially play a role in ME/CFS findings, for example through the stress associated with venipuncture. That being said, at least one study has tested prolactin response to placebo in ME/CFS, and found that patients did not exhibit an abnormally increased prolactin response following the placebo test [49].

Further, researchers should ensure that other significant medical conditions which could influence prolactin response are not present in the study cohort. For example, liver and kidney diseases affect metabolism of buspirone [142], and this may alter prolactin response to this drug. Even if disorders of metabolism are excluded, it may still be worthwhile to test for plasma concentrations of neuroendocrine probes. Also, as some studies have observed blunted prolactin response in depression [46, 87–94], it may be beneficial to exclude participants with comorbid depression, in order to reduce the likelihood of masking an ME/CFS-associated increase in prolactin response.

Limitations

While the prolactin response findings in ME/CFS studies are intriguing, some limitations limit the ability to make strong conclusions from the available data.

While five different published case-control studies have reported increased prolactin response to buspirone, these do not represent five different independent replications. Three of the five buspirone-prolactin studies shared an author/supervisor (P. O. Behan) [28, 32, 33], and all five studies were conducted in the United Kingdom. Additionally, the same control group data was used in two of these studies [32, 33]. Thus, findings may involve spurious effects specific to the few labs that have undertaken this research.

The sample size of each study was also relatively small,

with no study's ME/CFS group exceeding 30 cases. This raises the possibility of selection bias being responsible for the observations, as opposed to a finding generalizable to the ME/CFS population.

Studies testing prolactin challenge with exercise [48] and insulin-induced hypoglycemia [47] have observed a *blunted* response in ME/CFS, complicating a straightforward interpretation of over-responsivity of the prolactin system. Future research should explore whether these effects represent abnormalities in prolactin regulation, or instead relate to differences in exercise parameters and/or blood sugar regulation.

Study-specific limitations include weaknesses in describing control group characteristics. Richardson's 1995 paper does not report the sex ratio or ages of the healthy controls, and further indicates that some of the controls may have been parents of the ME/CFS patients [29]. Thus, sex- or age-related confounding may play a role in the study's findings. However, while sex ratio would be considered a clear confounder [35], several studies have failed to detect a significant correlation between age and prolactin response to buspirone in either healthy controls or individuals with depression [90, 96, 143]. Separately, Behan's 1996 paper of prolactin response included a significant error, describing the control group as 10 males instead of 30 individuals evenly split by sex [33]. As noted previously, an increased proportion of females in the control group would not be expected to lead to a spuriously increased prolactin response in cases [35], but the error still raises concerns about the study's data analysis procedures.

Further, as stress is known to increase prolactin [140, 141], differences related to drug side effects might explain the increased prolactin response seen in ME/CFS. Multiple papers have reported that ME/CFS patients demonstrate increased buspirone-induced side effects, including nausea and lightheadedness [28, 29, 31]. A greater degree of nausea could potentially lead to a larger prolactin response, by increasing patients' stress. However, even if side effects such as

nausea are responsible for the prolactin findings, the difference in side effects to buspirone would itself likely warrant further investigation.

Conclusion

Among the most replicated findings in ME/CFS research, abnormal prolactin response offers the opportunity to illuminate specific dysfunctional pathways underlying this condition. However, almost all studies into this phenomenon in ME/CFS were conducted in the 1990s, with no studies directly testing prolactin response in ME/CFS in over 15 years. The literature does not appear to include evidence that the ME/CFS prolactin findings were caused by spurious confounding. Instead, recent papers about ME/CFS continue to refer to these prolactin response findings as supporting evidence for their own claims [144, 145]

Duval et al. argue that neuroendocrine challenge testing, at least in the field of psychiatry, was largely abandoned in recent years for reasons which include the lack of standardization of neuroendocrine testing, difficulty in recruiting drug-naïve participants, and limited availability of drugs which selectively probe neurotransmitter systems [146]. One can imagine that prolactin response research in ME/CFS may have been a casualty of the pivot away from neuroendocrine research in other fields. The withdrawal of the serotonin-specific probe, d-fenfluramine, from the market in 1997 due to safety concerns [147] may also have played a role in slowing research. Still, as outlined in this review, numerous avenues are still available to continue this research.

Considering the massive societal impact of ME/CFS, both in terms of prevalence as well as the substantial suffering imposed on individuals, the importance of continuing this research is clear. Through careful consideration of study design and consultation with experts in endocrinology and neurology, future research into prolactin response may uncover important insights into the pathophysiology of ME/CFS, a condition which has, so far, eluded explanation.

References

- [1] E.-J. Lim, Y.-C. Ahn, E.-S. Jang, S.-W. Lee, S.-H. Lee, and C.-G. Son. "Systematic Review and Meta-Analysis of the Prevalence of Chronic Fatigue Syndrome/Myalgic Encephalomyelitis (CFS/ME)". In: *Journal of Translational Medicine* 18.1 (2020), p. 100. ISSN: 1479-5876. DOI: [10.1186/s12967-020-02269-0](https://doi.org/10.1186/s12967-020-02269-0). PMID: 32093722. URL: <https://link.springer.com/article/10.1186/s12967-020-02269-0>.
- [2] National Guideline Centre (UK). *Identifying and Diagnosing ME/CFS: Myalgic Encephalomyelitis (or Encephalopathy) / Chronic Fatigue Syndrome: Diagnosis and Management: Evidence Review D*. NICE Evidence Reviews Collection. London: National Institute for Health and Care Excellence (NICE), 2021. ISBN: 978-1-4731-4221-3. PMID: 35438857. URL: <http://www.ncbi.nlm.nih.gov/books/NBK579530/>.
- [3] *Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Redefining an Illness*. Washington, D.C.: National Academies Press, 2015, p. 19012. ISBN: 978-0-309-31689-7. DOI: [10.17226/19012](https://doi.org/10.17226/19012). URL: <http://www.nap.edu/catalog/19012>.

- [4] G. P. Holmes, J. E. Kaplan, N. M. Gantz, A. L. Komaroff, L. B. Schonberger, et al. "Chronic Fatigue Syndrome: A Working Case Definition." In: *Annals of internal medicine* 108.3 (1988), pp. 387–389. ISSN: 0003-4819. DOI: [10.7326/0003-4819-108-3-387](https://doi.org/10.7326/0003-4819-108-3-387). PMID: 2829679.
- [5] M. C. Sharpe, L. C. Archard, J. E. Banatvala, L. K. Borysiewicz, A. W. Clare, et al. "A Report—Chronic Fatigue Syndrome: Guidelines for Research." In: *Journal of the Royal Society of Medicine* 84.2 (1991), pp. 118–121. ISSN: 0141-0768. DOI: [10.1177/014107689108400224](https://doi.org/10.1177/014107689108400224). PMID: 1999813.
- [6] K. Fukuda, S. E. Straus, I. Hickie, M. C. Sharpe, J. G. Dobbins, and A. Komaroff. "The Chronic Fatigue Syndrome: A Comprehensive Approach to Its Definition and Study. International Chronic Fatigue Syndrome Study Group." In: *Annals of internal medicine* 121.12 (1994), pp. 953–959. ISSN: 0003-4819. DOI: [10.7326/0003-4819-121-12-199412150-00009](https://doi.org/10.7326/0003-4819-121-12-199412150-00009). PMID: 7978722.
- [7] B. M. Carruthers, A. K. Jain, K. L. De Meirleir, D. L. Peterson, N. G. Klimas, et al. "Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Clinical Working Case Definition, Diagnostic and Treatment Protocols". In: *Journal Of Chronic Fatigue Syndrome* 11.1 (2003), pp. 7–115. ISSN: 1057-3321. DOI: [10.1300/J092v11n01_02](https://doi.org/10.1300/J092v11n01_02). URL: https://www.tandfonline.com/doi/abs/10.1300/J092v11n01_02.
- [8] R. Maksoud, C. Magawa, N. Eaton-Fitch, K. Thapaliya, and S. Marshall-Gradisnik. "Biomarkers for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): A Systematic Review". In: *BMC Medicine* 21.1 (2023), p. 189. ISSN: 1741-7015. DOI: [10.1186/s12916-023-02893-9](https://doi.org/10.1186/s12916-023-02893-9). PMID: 37226227. URL: <https://link.springer.com/article/10.1186/s12916-023-02893-9>.
- [9] S. J. Mathew, X. Mao, K. A. Keegan, S. M. Levine, E. L. P. Smith, et al. "Ventricular Cerebrospinal Fluid Lactate Is Increased in Chronic Fatigue Syndrome Compared with Generalized Anxiety Disorder: An *in Vivo* 3.0 T¹ H MRS Imaging Study". In: *NMR in Biomedicine* 22.3 (2009), pp. 251–258. ISSN: 0952-3480, 1099-1492. DOI: [10.1002/nbm.1315](https://doi.org/10.1002/nbm.1315). URL: <https://analyticalsciencejournals.onlinelibrary.wiley.com/doi/10.1002/nbm.1315>.
- [10] J. W. Murrough, X. Mao, K. A. Collins, C. Kelly, G. Andrade, et al. "Increased Ventricular Lactate in Chronic Fatigue Syndrome Measured By¹ H MRS Imaging at 3.0 T. II: Comparison with Major Depressive Disorder". In: *NMR in Biomedicine* 23.6 (2010), pp. 643–650. ISSN: 0952-3480, 1099-1492. DOI: [10.1002/nbm.1512](https://doi.org/10.1002/nbm.1512). URL: <https://analyticalsciencejournals.onlinelibrary.wiley.com/doi/10.1002/nbm.1512>.
- [11] D. C. Shungu, N. Weiduschat, J. W. Murrough, X. Mao, S. Pillemer, et al. "Increased Ventricular Lactate in Chronic Fatigue Syndrome. III. Relationships to Cortical Glutathione and Clinical Symptoms Implicate Oxidative Stress in Disorder Pathophysiology". In: *NMR in biomedicine* 25.9 (2012), pp. 1073–1087. ISSN: 1099-1492. DOI: [10.1002/nbm.2772](https://doi.org/10.1002/nbm.2772). PMID: 22281935.
- [12] B. H. Natelson, X. Mao, A. J. Stegner, G. Lange, D. Vu, et al. "Multimodal and Simultaneous Assessments of Brain and Spinal Fluid Abnormalities in Chronic Fatigue Syndrome and the Effects of Psychiatric Comorbidity". In: *Journal of the neurological sciences* 375 (2017), pp. 411–416. ISSN: 0022-510X. DOI: [10.1016/j.jns.2017.02.046](https://doi.org/10.1016/j.jns.2017.02.046). PMID: 28320179. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5393352/>.
- [13] B. H. Natelson, D. Vu, J. D. Coplan, X. Mao, M. Blate, et al. "Elevations of Ventricular Lactate Levels Occur in Both Chronic Fatigue Syndrome and Fibromyalgia". In: *Fatigue: Biomedicine, Health & Behavior* 5.1 (2017), pp. 15–20. ISSN: 2164-1846, 2164-1862. DOI: [10.1080/21641846.2017.1280114](https://doi.org/10.1080/21641846.2017.1280114). URL: <https://www.tandfonline.com/doi/full/10.1080/21641846.2017.1280114>.
- [14] C. Mueller, J. C. Lin, S. Sheriff, A. A. Maudsley, and J. W. Younger. "Evidence of Widespread Metabolite Abnormalities in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Assessment with Whole-Brain Magnetic Resonance Spectroscopy." In: *Brain imaging and behavior* (2019), 10.1007/s11682-018-0029–4. ISSN: 1931-7557. DOI: [10.1007/s11682-018-0029-4](https://doi.org/10.1007/s11682-018-0029-4). PMID: 30617782. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6612467/>.
- [15] B. R. Godlewska, A. L. Sylvester, U. E. Emir, A. L. Sharpley, W. T. Clarke, et al. "Brain and Muscle Chemistry in Myalgic Encephalitis/Chronic Fatigue Syndrome (ME/CFS) and Long COVID: A 7T Magnetic Resonance Spectroscopy Study". In: *Molecular Psychiatry* 30.11 (2025), pp. 5215–5226. ISSN: 1476-5578. DOI: [10.1038/s41380-025-03108-8](https://doi.org/10.1038/s41380-025-03108-8). URL: <https://www.nature.com/articles/s41380-025-03108-8>.
- [16] A. A. Ryback and G. J. M. Cowan. "Deep Sequencing of BCR Heavy Chain Repertoires in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Frontiers in Immunology* 16 (2025), p. 1489312. ISSN: 1664-3224. DOI: [10.3389/fimmu.2025.1489312](https://doi.org/10.3389/fimmu.2025.1489312). URL: <https://www.frontiersin.org/articles/10.3389/fimmu.2025.1489312/full>.
- [17] M. Milivojevic, X. Che, L. Bateman, A. Cheng, B. A. Garcia, et al. "Plasma Proteomic Profiling Suggests an Association between Antigen Driven Clonal B Cell Expansion and ME/CFS". In: *PLOS ONE* 15.7 (2020), e0236148. ISSN: 1932-6203. DOI: [10.1371/journal.pone.0236148](https://doi.org/10.1371/journal.pone.0236148). URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0236148>.

- [18] W. Sato, H. Ono, T. Matsutani, M. Nakamura, I. Shin, et al. "Skewing of the B Cell Receptor Repertoire in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Brain, Behavior, and Immunity* 95 (2021), pp. 245–255. ISSN: 08891591. DOI: [10.1016/j.bbi.2021.03.023](https://doi.org/10.1016/j.bbi.2021.03.023). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0889159121001537>.
- [19] J. N. Baraniuk, N. Eaton-Fitch, and S. Marshall-Gradisnik. "Meta-Analysis of Natural Killer Cell Cytotoxicity in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome." In: *Frontiers in immunology* 15 (2024), p. 1440643. ISSN: 1664-3224. DOI: [10.3389/fimmu.2024.1440643](https://doi.org/10.3389/fimmu.2024.1440643). PMID: 39483457.
- [20] M. L. Gordon, R. B. Lipton, S. L. Brown, and H. M. van Praag. "The Neuroendocrine Challenge Paradigm in Headache Research". In: *Cephalalgia: An International Journal of Headache* 15.4 (1995), pp. 292–296. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.1995.1504292.x](https://doi.org/10.1046/j.1468-2982.1995.1504292.x). PMID: 7585926.
- [21] G. Hatzidimitriou, E. H. Tsai, U. D. McCann, and G. A. Ricaurte. "Altered Prolactin Response to M-chlorophenylpiperazine in Monkeys Previously Treated with 3,4-Methylenedioxymethamphetamine (MDMA) or Fenfluramine". In: *Synapse* 44.1 (2002), pp. 51–57. ISSN: 1098-2396. DOI: [10.1002/syn.10055](https://doi.org/10.1002/syn.10055). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/syn.10055>.
- [22] A. Quattrone, G. Schettini, L. Annunziato, and G. Di Renzo. "Pharmacological Evidence of Supersensitivity of Central Serotonergic Receptors Involved in the Control of Prolactin Secretion". In: *European Journal of Pharmacology* 76.1 (1981), pp. 9–13. ISSN: 00142999. DOI: [10.1016/0014-2999\(81\)90003-0](https://doi.org/10.1016/0014-2999(81)90003-0). URL: <https://linkinghub.elsevier.com/retrieve/pii/0014299981900030>.
- [23] C. M. Kuhn, R. A. Vogel, R. B. Mailman, R. A. Mueller, S. M. Schanberg, and G. R. Breese. "Effect of 5,7-Dihydroxytryptamine on Serotonergic Control of Prolactin Secretion and Behavior in Rats". In: *Psychopharmacology* 73.2 (1981), pp. 188–193. ISSN: 1432-2072. DOI: [10.1007/BF00429216](https://doi.org/10.1007/BF00429216). URL: <https://link.springer.com/article/10.1007/BF00429216>.
- [24] D. R. Grattan. "60 YEARS OF NEUROENDOCRINOLOGY: The Hypothalamo-Prolactin Axis". In: *Journal of Endocrinology* 226.2 (2015), T101–T122. ISSN: 0022-0795, 1479-6805. DOI: [10.1530/JOE-15-0213](https://doi.org/10.1530/JOE-15-0213). URL: <http://journals.bioscientifica.com/joe/article/226/2/T101-T122/22808>.
- [25] M. E. Freeman, B. Kanyicska, A. Lerant, and G. Nagy. "Prolactin: Structure, Function, and Regulation of Secretion". In: *Physiological Reviews* 80.4 (2000), pp. 1523–1631. ISSN: 0031-9333, 1522-1210. DOI: [10.1152/physrev.2000.80.4.1523](https://doi.org/10.1152/physrev.2000.80.4.1523). URL: <https://www.physiology.org/doi/10.1152/physrev.2000.80.4.1523>.
- [26] C. Loane and M. Politis. "Buspirone: What Is It All About?" In: *Brain Research* 1461 (2012), pp. 111–118. ISSN: 0006-8993. DOI: [10.1016/j.brainres.2012.04.032](https://doi.org/10.1016/j.brainres.2012.04.032). URL: <https://www.sciencedirect.com/science/article/pii/S0006899312007263>.
- [27] A. M. O. Bakheit. "Hypothalamic Dysfunction and Neurotransmitter Abnormalities in the Postviral Fatigue Syndrome". MD. Ann Arbor : ProQuest Dissertations & Theses, 1992. 215 pp. URL: <https://theses.gla.ac.uk/75589/>.
- [28] A. M. O. Bakheit, P. O. Behan, T. G. Dinan, C. E. Gray, and V. O'Keane. "Possible Upregulation of Hypothalamic 5-Hydroxytryptamine Receptors in Patients with Postviral Fatigue Syndrome." In: *BMJ : British Medical Journal* 304.6833 (1992), pp. 1010–1012. ISSN: 0959-8138. PMID: 1586780. URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1881733/>.
- [29] J. Richardson. "Disturbance of Hypothalamic Function and Evidence for Persistent Enteroviral Infection in Patients with Chronic Fatigue Syndrome". In: *Journal Of Chronic Fatigue Syndrome* 1.2 (1995), pp. 59–66. ISSN: 1057-3321. DOI: [10.1300/J092v01n02_05](https://doi.org/10.1300/J092v01n02_05). URL: https://www.tandfonline.com/doi/abs/10.1300/J092v01n02_05.
- [30] J. Richardson and D. C. Costa. "Relationship Between SPECT Scans and Buspirone Tests in Patients with ME/CFS". In: *Journal of Chronic Fatigue Syndrome* 4.3 (1998), pp. 23–38. ISSN: 1057-3321, 1547-0660. DOI: [10.1300/J092v04n03_04](https://doi.org/10.1300/J092v04n03_04). URL: http://www.tandfonline.com/doi/full/10.1300/J092v04n03_04.
- [31] M. Sharpe, A. Clements, K. Hawton, A. H. Young, P. Sargent, and P. J. Cowen. "Increased Prolactin Response to Buspirone in Chronic Fatigue Syndrome". In: *Journal of Affective Disorders* 41.1 (1996), pp. 71–76. ISSN: 0165-0327. DOI: [10.1016/0165-0327\(96\)00075-4](https://doi.org/10.1016/0165-0327(96)00075-4). URL: <https://www.sciencedirect.com/science/article/pii/0165032796000754>.
- [32] T. Majeed. "Neuroendocrine Alterations in Chronic Fatigue Syndrome". PhD thesis. Ann Arbor : ProQuest Dissertations & Theses, 1996. 281 pp. URL: <https://theses.gla.ac.uk/75796/>.
- [33] P. O. Behan. "Chronic Fatigue Syndrome as a Delayed Reaction to Chronic Low-dose Organophosphate Exposure". In: *Journal of Nutritional & Environmental Medicine* 6.4 (1996), pp. 341–350. ISSN: 1359-0847. DOI: [10.3109/13590849609007262](https://doi.org/10.3109/13590849609007262). URL: <https://www.tandfonline.com/doi/abs/10.3109/13590849609007262>.

- [34] *Professor's Future in Doubt. University Inquiry Blames Academic's Sloppy Preparation for Court Fiasco*. The Herald. 1997. URL: <https://www.heraldscotland.com/news/12294532.professors-future-in-doubt-university-inquiry-blames-academics-sloppy-preparation-for-court-fiasco/>.
- [35] T. G. Dinan, S. Barry, L. N. Yatham, M. Mobayed, and M. O'Hanlon. "The Reproducibility of the Prolactin Response to Buspirone: Relationship to the Menstrual Cycle". In: *International Clinical Psychopharmacology* 5.2 (1990), pp. 119–123. ISSN: 0268-1315. DOI: [10.1097/00004850-199004000-00006](https://doi.org/10.1097/00004850-199004000-00006). PMID: [2380543](https://pubmed.ncbi.nlm.nih.gov/2380543/).
- [36] D. Racciatti, J. Vecchiet, A. Ceccomancini, F. Ricci, and E. Pizzigallo. "Chronic Fatigue Syndrome Following a Toxic Exposure". In: *Science of The Total Environment*. Environment, Human Health and Immune System (Selected Papers from International Symposium on Occupational and Environmental Allergy and Immune Diseases, Chieti (Italy), April 27–30, 1999 270.1 (2001), pp. 27–31. ISSN: 0048-9697. DOI: [10.1016/S0048-9697\(00\)00777-4](https://doi.org/10.1016/S0048-9697(00)00777-4). URL: <https://www.sciencedirect.com/science/article/pii/S0048969700007774>.
- [37] D. Racciatti, A. Barberio, E. Pizzigallo, et al. "Neuroendocrine Perspective of Chronic Fatigue Syndrome". In: *Journal of Musculoskeletal Pain* 6.S2 (1998), p. 152.
- [38] D. Racciatti, A. Barberio, J. Vecchiet, and E. Pizzigallo. "Clinical and Pathogenetical Characterization of 238 Patients of a Chronic Fatigue Syndrome Italian Center". In: *Journal Of Chronic Fatigue Syndrome* 5.3–4 (1999), pp. 61–70. ISSN: 1057-3321. DOI: [10.1300/J092v05n03_05](https://doi.org/10.1300/J092v05n03_05). URL: https://www.tandfonline.com/doi/abs/10.1300/J092v05n03_05.
- [39] A. Sharma, F. Oyeboode, M. J. Kendall, and D. A. Jones. "Recovery from Chronic Fatigue Syndrome Associated with Changes in Neuroendocrine Function". In: *Journal of the Royal Society of Medicine* 94.1 (2001), pp. 26–27. ISSN: 0141-0768, 1758-1095. DOI: [10.1177/014107680109400107](https://doi.org/10.1177/014107680109400107). URL: <https://journals.sagepub.com/doi/10.1177/014107680109400107>.
- [40] *Forum Thread - Hypothalamic Dysfunction and Neurotransmitter Abnormalities in the Postviral Fatigue Syndrome, 1992, Bakheit*. Science for ME. 2026. URL: <https://s4me.info/threads/hypothalamic-dysfunction-and-neurotransmitter-abnormalities-in-the-postviral-fatigue-syndrome-1992-bakheit.51528/post-709853>.
- [41] A. Rohatgi. *WebPlotDigitizer*. Version 5.2. URL: <https://automeris.io>.
- [42] L. E. Vivero, P. O. Anderson, and R. F. Clark. "A Close Look at Fenfluramine and Dexfenfluramine". In: *The Journal of Emergency Medicine* 16.2 (1998), pp. 197–205. ISSN: 0736-4679. DOI: [10.1016/S0736-4679\(97\)00289-8](https://doi.org/10.1016/S0736-4679(97)00289-8). URL: <https://www.sciencedirect.com/science/article/pii/S0736467997002898>.
- [43] K. A. Bever and P. J. Perry. "Dexfenfluramine Hydrochloride: An Anorexigenic Agent". In: *American Journal of Health-System Pharmacy* 54.18 (1997), pp. 2059–2072. ISSN: 1079-2082. DOI: [10.1093/ajhp/54.18.2059](https://doi.org/10.1093/ajhp/54.18.2059). URL: <https://academic.oup.com/ajhp/article-abstract/54/18/2059/5158304>.
- [44] L. N. Yatham, R. L. Morehouse, B. T. Chisholm, D. A. Haase, D. D. MacDonald, and T. J. Marrie. "Neuroendocrine Assessment of Serotonin (5-HT) Function in Chronic Fatigue Syndrome". In: *Canadian Journal of Psychiatry. Revue Canadienne De Psychiatrie* 40.2 (1995), pp. 93–96. ISSN: 0706-7437. DOI: [10.1177/070674379504000207](https://doi.org/10.1177/070674379504000207). PMID: [7788624](https://pubmed.ncbi.nlm.nih.gov/7788624/). URL: <https://journals.sagepub.com/doi/abs/10.1177/070674379504000207>.
- [45] M. Sharpe, K. Hawton, A. Clements, and P. Cowen. "Increased Brain Serotonin Function in Men with Chronic Fatigue Syndrome". In: *BMJ* 315.7101 (1997), pp. 164–165. ISSN: 0959-8138, 1468-5833. DOI: [10.1136/bmj.315.7101.164](https://doi.org/10.1136/bmj.315.7101.164). URL: <https://www.bmj.com/lookup/doi/10.1136/bmj.315.7101.164>.
- [46] A. J. Cleare, J. Bearn, T. Allain, A. McGregor, S. Wessely, R. M. Murray, and V. O'Keane. "Contrasting Neuroendocrine Responses in Depression and Chronic Fatigue Syndrome". In: *Journal of Affective Disorders* 34.4 (1995), pp. 283–289. ISSN: 0165-0327. DOI: [10.1016/0165-0327\(95\)00026-J](https://doi.org/10.1016/0165-0327(95)00026-J). URL: <https://www.sciencedirect.com/science/article/pii/S016503279500026J>.
- [47] J. Bearn, T. Allain, P. Coskeran, N. Munro, J. Butler, A. McGregor, and S. Wessely. "Neuroendocrine Responses to D-Fenfluramine and Insulin-Induced Hypoglycemia in Chronic Fatigue Syndrome". In: *Biological Psychiatry* 37.4 (1995), pp. 245–252. ISSN: 00063223. DOI: [10.1016/0006-3223\(94\)00121-I](https://doi.org/10.1016/0006-3223(94)00121-I). URL: <https://linkinghub.elsevier.com/retrieve/pii/S000632239400121I>.
- [48] J. E. Ottenweller, S. A. Sisto, R. C. McCarty, and B. H. Natelson. "Hormonal Responses to Exercise in Chronic Fatigue Syndrome". In: *Neuropsychobiology* 43.1 (2001), pp. 34–41. ISSN: 0302-282X, 1423-0224. DOI: [10.1159/000054863](https://doi.org/10.1159/000054863). URL: <https://karger.com/article/doi/10.1159/000054863>.
- [49] C. M. Vassallo, E. Feldman, T. Peto, L. Castell, A. L. Sharpley, and P. J. Cowen. "Decreased Tryptophan Availability but Normal Post-Synaptic 5-HT_{2c} Receptor Sensitivity in Chronic Fatigue Syndrome". In: *Psychological Medicine* 31.4 (2001), pp. 585–591. ISSN: 0033-2917, 1469-8978. DOI: [10.1017/S0033291701003580](https://doi.org/10.1017/S0033291701003580). PMID: [11352361](https://pubmed.ncbi.nlm.nih.gov/11352361/). URL: <https://www.cambridge.org/core/journals/psychological-medicine/article/abs/decreased-tryptophan-availability-but-normal-postsynaptic-5ht2c-receptor-sensitivity-in-chronic-fatigue-syndrome/7444BBC820CA52CC5434346D6F53CCD5>.

- [50] S. A. Weaver, M. N. Janal, N. Aktan, J. E. Ottenweller, and B. H. Natelson. "Sex Differences in Plasma Prolactin Response to Tryptophan in Chronic Fatigue Syndrome Patients With and Without Comorbid Fibromyalgia". In: *Journal of Women's Health* 19.5 (2010), pp. 951–958. ISSN: 1540-9996. DOI: [10.1089/jwh.2009.1697](https://doi.org/10.1089/jwh.2009.1697). URL: <https://journals.sagepub.com/doi/abs/10.1089/jwh.2009.1697>.
- [51] E. M. Cassidy, E. Tomkins, T. Dinan, O. Hardiman, and V. O'Keane. "Central 5-HT Receptor Hypersensitivity in Migraine without Aura". In: *Cephalalgia: An International Journal of Headache* 23.1 (2003), pp. 29–34. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.2003.00441.x](https://doi.org/10.1046/j.1468-2982.2003.00441.x). PMID: [12534577](https://pubmed.ncbi.nlm.nih.gov/12534577/).
- [52] E. M. Cassidy, E. Tomkins, N. Sharifi, T. Dinan, O. Hardiman, and V. O'Keane. "Differing Central Amine Receptor Sensitivity in Different Migraine Subtypes? A Neuroendocrine Study Using Buspirone". In: *Pain* 101.3 (2003), pp. 283–290. ISSN: 0304-3959. DOI: [10.1016/S0304-3959\(02\)00335-4](https://doi.org/10.1016/S0304-3959(02)00335-4). PMID: [12583871](https://pubmed.ncbi.nlm.nih.gov/12583871/).
- [53] M. Leone, A. Attanasio, D. Croci, A. Ferraris, D. D'Amico, et al. "5-HT_{1A} Receptor Hypersensitivity in Migraine Is Suggested by the m-Chlorophenylpiperazine Test". In: *Neuroreport* 9.11 (1998), pp. 2605–2608. ISSN: 0959-4965. DOI: [10.1097/00001756-199808030-00033](https://doi.org/10.1097/00001756-199808030-00033). PMID: [9721941](https://pubmed.ncbi.nlm.nih.gov/9721941/).
- [54] M. L. Gordon, R. B. Lipton, S.-L. Brown, C. Nakraseive, M. Russell, et al. "Headache and Cortisol Responses to M-Chlorophenylpiperazine Are Highly Correlated". In: *Cephalalgia* 13.6 (1993), pp. 400–405. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.1993.1306400.x](https://doi.org/10.1046/j.1468-2982.1993.1306400.x). URL: <https://journals.sagepub.com/doi/10.1046/j.1468-2982.1993.1306400.x>.
- [55] V. Glover, F. Ahmed, N. Hussein, J. Jarman, and R. Peatfield. "Central 5-Hydroxytryptamine Supersensitivity in Migraine". In: *Migraine: Pharmacology and Genetics*. Chapman & Hall, 1996, pp. 117–123. ISBN: 1-86036-006-8. URL: <https://archive.org/details/migrainepharmac0000unse/page/116/>.
- [56] G. Nappi, F. Savoldi, G. Bono, and E. Martignoni. "Reserpine - Headache and PRL Release in Migraine". In: *Headache: The Journal of Head and Face Pain* 19.5 (1979), pp. 273–277. ISSN: 1526-4610. DOI: [10.1111/j.1526-4610.1979.hed1905273.x](https://doi.org/10.1111/j.1526-4610.1979.hed1905273.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1526-4610.1979.hed1905273.x>.
- [57] A. Polleri, G. Nappi, P. Masturzo, E. Martignoni, G. Murialdo, et al. "Neuroendocrine Approach to Headache". In: *Advances in Neurology*. Vol. 33. 1982, pp. 173–182. URL: <https://archive.org/details/headachephysiopa0000unse/page/172/mode/2up/>.
- [58] G. Murialdo, E. Martignoni, A. D. Maria, M. L. Bonura, G. Sances, G. Bono, and A. Polleri. "Changes in the Dopaminergic Control of Prolactin Secretion and in Ovarian Steroids in Migraine". In: *Cephalalgia* 6.1 (1986), pp. 43–49. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.1986.0601043.x](https://doi.org/10.1046/j.1468-2982.1986.0601043.x).
- [59] P. Calabresi, M. Silvestrini, F. Stratta, L. Cupini, G. Argiro, G. Atzei, and G. Bernardi. "L-Deprenyl Test in Migraine: Neuroendocrinological Aspects". In: *Cephalalgia* 13.6 (1993), pp. 406–409. ISSN: 1468-2982. DOI: [10.1046/j.1468-2982.1993.1306406.x](https://doi.org/10.1046/j.1468-2982.1993.1306406.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1468-2982.1993.1306406.x>.
- [60] G. Nattero, M. Corno, L. Savi, G. Isaia, C. Priolo, and M. Mussetta. "Prolactin and Migraine: Effect of L-Dopa on Plasma Prolactin Levels in Migraineurs and Normals". In: *Headache: The Journal of Head and Face Pain* 26.1 (1986), pp. 9–12. ISSN: 1526-4610. DOI: [10.1111/j.1526-4610.1986.hed2601009.x](https://doi.org/10.1111/j.1526-4610.1986.hed2601009.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1526-4610.1986.hed2601009.x>.
- [61] G. Murialdo, P. Masturzo, U. Filippi, D. De Palma, D. Balbi, et al. "Thyroid-Stimulating Hormone and Prolactin Responses to Thyrotropin-Releasing Hormone in Common Migraine". In: *Cephalalgia* 7.4 (1987), pp. 267–272. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.1987.0704267.x](https://doi.org/10.1046/j.1468-2982.1987.0704267.x). URL: <https://journals.sagepub.com/doi/10.1046/j.1468-2982.1987.0704267.x>.
- [62] Y. Papakostas, M. Daras, M. Markianos, and C. Stefanis. "Increased Prolactin Response to Thyrotropin Releasing Hormone during Migraine Attacks." In: *Journal of Neurology, Neurosurgery & Psychiatry* 50.7 (1987), pp. 927–928. ISSN: 0022-3050, 1468-330X. DOI: [10.1136/jnnp.50.7.927](https://doi.org/10.1136/jnnp.50.7.927). PMID: [3114432](https://pubmed.ncbi.nlm.nih.gov/3114432/). URL: <https://jnnp.bmj.com/content/50/7/927>.
- [63] E. Awaki, T. Takeshima, and K. Takahashi. "A Neuroendocrinological Study in Female Migraineurs: Prolactin and Thyroid Stimulating Hormone Responses". In: *Cephalalgia* 9.3 (1989), pp. 187–193. ISSN: 0333-1024. DOI: [10.1046/j.1468-2982.1989.0903187.x](https://doi.org/10.1046/j.1468-2982.1989.0903187.x). URL: <https://journals.sagepub.com/doi/abs/10.1046/j.1468-2982.1989.0903187.x>.
- [64] J. L. Capellan, C. V. Dacosta, J. G. Garcia, P. R. Servan, and J. A. Cermeno. "Tuberoinfundibular Dopaminergic Tonus in Common Migraine". In: *Headache: The Journal of Head and Face Pain* 30.5 (1990), pp. 282–284. ISSN: 1526-4610. DOI: [10.1111/j.1526-4610.1990.hed3005282.x](https://doi.org/10.1111/j.1526-4610.1990.hed3005282.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1526-4610.1990.hed3005282.x>.
- [65] A. Klimek. "Growth Hormone and Prolactin Release in Migraine Patients". In: *Updating in Headache*. Ed. by V. Pfaffenrath, P.-O. Lundberg, and O. Sjaastad. Berlin, Heidelberg: Springer, 1985, pp. 241–248. ISBN: 978-3-642-88581-5. DOI: [10.1007/978-3-642-88581-5_40](https://doi.org/10.1007/978-3-642-88581-5_40).

- [66] A. Klimek. "Metoclopramide Tests in Headache Patients". In: *Cephalalgia* 5 (3_suppl 1985), pp. 354–355. ISSN: 0333-1024. DOI: [10.1177/03331024850050S3138](https://doi.org/10.1177/03331024850050S3138). URL: <https://journals.sagepub.com/doi/10.1177/03331024850050S3138>.
- [67] G. Bussone, A. Boiardi, L. L. Mantia, F. Frediani, A. Vescovi, and E. A. Parati. "Long-Acting Ergot: Clinical and Neuroendocrinological Aspects in Migraine Patients". In: *Cephalalgia* 3 (1_suppl 1983), pp. 163–167. ISSN: 0333-1024. DOI: [10.1177/03331024830030S127](https://doi.org/10.1177/03331024830030S127). URL: <https://journals.sagepub.com/doi/10.1177/03331024830030S127>.
- [68] G. Bussone, F. Frediani, E. Lamperti, L. LaMantia, A. Vescovi, C. Peccarisi, and A. Boiardi. "Piribedil Test in Migraine: Neuroendocrinological Aspects". In: *Headache: The Journal of Head and Face Pain* 26.9 (1986), pp. 482–485. ISSN: 1526-4610. DOI: [10.1111/j.1526-4610.1986.hed2609482.x](https://doi.org/10.1111/j.1526-4610.1986.hed2609482.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1526-4610.1986.hed2609482.x>.
- [69] I. Rainero, W. Valfrè, L. Savi, S. Gentile, L. Pinessi, et al. "Neuroendocrine Effects of Subcutaneous Sumatriptan in Patients with Migraine". In: *Journal of Endocrinological Investigation* 24.5 (2001), pp. 310–314. ISSN: 0391-4097. DOI: [10.1007/BF03343866](https://doi.org/10.1007/BF03343866). PMID: [11407649](https://pubmed.ncbi.nlm.nih.gov/11407649/).
- [70] J. Leer, B. Patel, M. Innes, and D. P. Cameron. "Secondary Amenorrhoea Due to Autoimmune Ovarian Failure". In: *Australian and New Zealand Journal of Obstetrics and Gynaecology* 20.3 (1980), pp. 177–179. ISSN: 1479-828X. DOI: [10.1111/j.1479-828X.1980.tb02902.x](https://doi.org/10.1111/j.1479-828X.1980.tb02902.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1479-828X.1980.tb02902.x>.
- [71] D. Hochner-Celnikier, E. Zylber-Haran, S. Shilo, Z. Palti, and I. M. Spitz. "Increased Prolactin Response to Thyrotropin-Releasing Hormone in Primary Ovarian Failure". In: *Obstetrics and Gynecology* 59.3 (1982), pp. 280–284. ISSN: 0029-7844. PMID: [6804899](https://pubmed.ncbi.nlm.nih.gov/6804899/).
- [72] I. M. Spitz, E. Zylber, H. Cohen, U. Almaliach, and D. Leroith. "Impaired Prolactin Response to Thyrotropin-Releasing Hormone in Isolated Gonadotropin Deficiency and Exaggerated Response in Primary Testicular Failure". In: *The Journal of Clinical Endocrinology and Metabolism* 48.6 (1979), pp. 941–945. ISSN: 0021-972X. DOI: [10.1210/jcem-48-6-941](https://doi.org/10.1210/jcem-48-6-941). PMID: [109459](https://pubmed.ncbi.nlm.nih.gov/109459/).
- [73] I. M. Spitz, D. Leroith, Y. Livshin, E. Zylber-Haran, S. Trestian, et al. "Exaggerated Prolactin Response to Thyrotropin-Releasing Hormone and Metoclopramide in Primary Testicular Failure". In: *Fertility and Sterility* 34.6 (1980), pp. 573–580. ISSN: 0015-0282. DOI: [10.1016/S0015-0282\(16\)45198-8](https://doi.org/10.1016/S0015-0282(16)45198-8). URL: <https://www.sciencedirect.com/science/article/pii/S0015028216451988>.
- [74] I. M. Spitz, P. Lavie, N. Calderon, C. R. Gordon, A. Oksenberg, et al. "Dissociation between Sleep-Related and TRH-induced Prolactin Secretion in Seminiferous Tubule Failure". In: *Metabolism* 31.1 (1982), pp. 10–13. ISSN: 0026-0495. DOI: [10.1016/0026-0495\(82\)90020-8](https://doi.org/10.1016/0026-0495(82)90020-8). URL: <https://www.sciencedirect.com/science/article/pii/0026049582900208>.
- [75] D. Leroith, G. Potashnik, J. Dunn, and I. M. Spitz. "The Exaggerated Prolactin Response to Thyrotropin-Releasing Hormone and Metoclopramide in 1,2-Dibromo-3-Chloropropane-Induced Azoospermia". In: *The Journal of Clinical Endocrinology and Metabolism* 52.1 (1981), pp. 38–41. ISSN: 0021-972X. DOI: [10.1210/jcem-52-1-38](https://doi.org/10.1210/jcem-52-1-38). PMID: [6778892](https://pubmed.ncbi.nlm.nih.gov/6778892/).
- [76] I. M. Spitz, B. Levitt, E. Zylber-Haran, M. Haas, and Y. Livshin. "The Dissociation of the Exaggerated Prolactin and Thyrotropin Responses in Seminiferous Tubule Failure Following the Administration of a Double-Pulse of Thyrotropin-Releasing Hormone". In: *Journal of Endocrinological Investigation* 6.4 (1983), pp. 273–276. ISSN: 0391-4097. DOI: [10.1007/BF03347589](https://doi.org/10.1007/BF03347589). PMID: [6417218](https://pubmed.ncbi.nlm.nih.gov/6417218/).
- [77] B. Baranowska, W. Jeske, A. Niewiadomska, G. Rozbicka, L. Walczak, and S. Zgliczynski. "Enhanced Serum Prolactin Concentration after Metoclopramide Stimulation in Idiopathic Oligozoospermia and Azoospermia". In: *Andrologia* 15.S1 (1983), pp. 554–559. ISSN: 03034569, 14390272. DOI: [10.1111/j.1439-0272.1983.tb00211.x](https://doi.org/10.1111/j.1439-0272.1983.tb00211.x). URL: <https://onlinelibrary.wiley.com/doi/10.1111/j.1439-0272.1983.tb00211.x>.
- [78] D. Boucher, G. Grizard, J. Hermabessiere, G. Gaillard, and J. Pays. "Pituitary Gonadal Function in Infertile Men with Varicocele". In: *Andrologia* 15.1 (1983), pp. 78–89. ISSN: 03034569, 14390272. DOI: [10.1111/j.1439-0272.1983.tb00120.x](https://doi.org/10.1111/j.1439-0272.1983.tb00120.x). URL: <https://onlinelibrary.wiley.com/doi/10.1111/j.1439-0272.1983.tb00120.x>.
- [79] G. Goldhaber, Z. Zuckerman, J. Shani, and J. J. Cohen. "Effect of Thyrotrophin-Releasing Hormone on Serum Prolactin Levels in Men with Azoospermia". In: *Journal of Reproduction and Fertility* 49.1 (1977), pp. 135–137. ISSN: 0022-4251. DOI: [10.1530/jrf.0.0490135](https://doi.org/10.1530/jrf.0.0490135). URL: <https://academic.oup.com/reproduction/article-abstract/49/1/135/8289647?redirectedFrom=fulltext>.
- [80] A. Masala, G. Delitala, S. Alagna, L. Devilla, P. P. Rovasio, and G. Lotti. "Dynamic Evaluation of Prolactin Secretion in Patients with Oligospermia: Effects of Treatment with Metergoline". In: *Fertility and Sterility* 31.1 (1979), pp. 63–67. ISSN: 0015-0282. DOI: [10.1016/s0015-0282\(16\)43762-3](https://doi.org/10.1016/s0015-0282(16)43762-3). PMID: [105938](https://pubmed.ncbi.nlm.nih.gov/105938/).

- [81] S. J. Winters, R. E. Johnsonbaugh, and R. J. Sherins. "The Response of Prolactin to Chlorpromazine Stimulation in Men with Hypogonadotrophic Hypogonadism and Early Pubertal Boys: Relationship to Sex Steroid Exposure". In: *Clinical Endocrinology* 16.4 (1982), pp. 321–330. ISSN: 1365-2265. DOI: [10.1111/j.1365-2265.1982.tb00723.x](https://doi.org/10.1111/j.1365-2265.1982.tb00723.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2265.1982.tb00723.x>.
- [82] S. O'Mahony, A. S. B. Chua, E. M. M. Quigley, G. Clarke, F. Shanahan, P. W. N. Keeling, and T. G. Dinan. "Evidence of an Enhanced Central 5HT Response in Irritable Bowel Syndrome and in the Rat Maternal Separation Model". In: *Neurogastroenterology and Motility* 20.6 (2008), pp. 680–688. ISSN: 1365-2982. DOI: [10.1111/j.1365-2982.2007.01065.x](https://doi.org/10.1111/j.1365-2982.2007.01065.x). PMID: [18194152](https://pubmed.ncbi.nlm.nih.gov/18194152/).
- [83] A. Chua, J. Keating, D. Hamilton, P. W. Keeling, and T. G. Dinan. "Central Serotonin Receptors and Delayed Gastric Emptying in Non-Ulcer Dyspepsia". In: *BMJ* 305.6848 (1992), pp. 280–282. ISSN: 0959-8138. DOI: [10.1136/bmj.305.6848.280](https://doi.org/10.1136/bmj.305.6848.280). PMID: [1392859](https://pubmed.ncbi.nlm.nih.gov/1392859/).
- [84] T. G. Dinan, A. S. Chua, and P. W. Keeling. "Serotonin and Physical Illness: Focus on Non-Ulcer Dyspepsia". In: *Journal of Psychopharmacology* 7.1 (1993), pp. 126–130. ISSN: 0269-8811. DOI: [10.1177/026988119300700104](https://doi.org/10.1177/026988119300700104). PMID: [22289664](https://pubmed.ncbi.nlm.nih.gov/22289664/).
- [85] T. G. Dinan, N. Mahmud, O. Rathore, J. Thakore, L. V. Scott, et al. "A Double-Blind Placebo-Controlled Study of Buspirone-Stimulated Prolactin Release in Non-Ulcer Dyspepsia—Are Central Serotonergic Responses Enhanced?" In: *Alimentary Pharmacology & Therapeutics* 15.10 (2001), pp. 1613–1618. ISSN: 1365-2036. DOI: [10.1046/j.1365-2036.2001.01090.x](https://doi.org/10.1046/j.1365-2036.2001.01090.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2036.2001.01090.x>.
- [86] T. G. Dinan, L. N. Yatham, S. Barry, A. Chua, and P. W. Keeling. "Serotonin Supersensitivity: The Pathophysiologic Basis of Non-Ulcer Dyspepsia? A Preliminary Report of Buspirone/Prolactin Responses". In: *Scandinavian Journal of Gastroenterology* 25.5 (1990), pp. 541–544. ISSN: 0036-5521. DOI: [10.3109/00365529009095527](https://doi.org/10.3109/00365529009095527). PMID: [2359983](https://pubmed.ncbi.nlm.nih.gov/2359983/).
- [87] R. N. Golden, J. K. Hsiao, E. Lane, D. Ekstrom, S. Rogers, R. Hicks, and W. Z. Potter. "Abnormal Neuroendocrine Responsivity to Acute i.v. Clomipramine Challenge in Depressed Patients". In: *Psychiatry Research* 31.1 (1990), pp. 39–47. ISSN: 0165-1781. DOI: [10.1016/0165-1781\(90\)90107-g](https://doi.org/10.1016/0165-1781(90)90107-g). PMID: [2156276](https://pubmed.ncbi.nlm.nih.gov/2156276/).
- [88] I. M. Anderson, C. J. Ware, J. M. da Roza Davis, and P. J. Cowen. "Decreased 5-HT-mediated Prolactin Release in Major Depression". In: *The British Journal of Psychiatry: The Journal of Mental Science* 160 (1992), pp. 372–378. ISSN: 0007-1250. DOI: [10.1192/bjp.160.3.372](https://doi.org/10.1192/bjp.160.3.372). PMID: [1562864](https://pubmed.ncbi.nlm.nih.gov/1562864/).
- [89] J. V. Lucey, V. O'Keane, G. Butcher, A. W. Clare, and T. G. Dinan. "Cortisol and Prolactin Responses to D-Fenfluramine in Non-Depressed Patients with Obsessive-Compulsive Disorder: A Comparison with Depressed and Healthy Controls". In: *The British Journal of Psychiatry: The Journal of Mental Science* 161 (1992), pp. 517–521. ISSN: 0007-1250. DOI: [10.1192/bjp.161.4.517](https://doi.org/10.1192/bjp.161.4.517). PMID: [1393339](https://pubmed.ncbi.nlm.nih.gov/1393339/).
- [90] F. G. Moeller, J. L. Steinberg, M. Fulton, G. Kramer, and F. Petty. "A Preliminary Neuroendocrine Study with Buspirone in Major Depression". In: *Neuropsychopharmacology* 10.2 (1994), pp. 75–83. ISSN: 1740-634X. DOI: [10.1038/npp.1994.9](https://doi.org/10.1038/npp.1994.9). URL: <https://www.nature.com/articles/npp19949>.
- [91] T. Kapitany, M. Schindl, S. D. Schindler, B. Heßelmann, T. Füreder, et al. "The Citalopram Challenge Test in Patients with Major Depression and in Healthy Controls". In: *Psychiatry Research* 88.2 (1999), pp. 75–88. ISSN: 0165-1781. DOI: [10.1016/S0165-1781\(99\)00082-7](https://doi.org/10.1016/S0165-1781(99)00082-7). URL: <https://www.sciencedirect.com/science/article/pii/S0165178199000827>.
- [92] H. Corrêa, F. Duval, M.-C. Mokrani, P. Bailey, F. Trémeau, et al. "Prolactin Response to D-Fenfluramine and Suicidal Behavior in Depressed Patients". In: *Psychiatry Research* 93.3 (2000), pp. 189–199. ISSN: 0165-1781. DOI: [10.1016/S0165-1781\(00\)00114-1](https://doi.org/10.1016/S0165-1781(00)00114-1). URL: <https://www.sciencedirect.com/science/article/pii/S0165178100001141>.
- [93] M. Markianos, J. Hatzimanolis, and L. Lykouras. "Relationship between Prolactin Responses to ECT and Dopaminergic and Serotonergic Responsivity in Depressed Patients". In: *European Archives of Psychiatry and Clinical Neuroscience* 252.4 (2002), pp. 166–171. ISSN: 1433-8491. DOI: [10.1007/s00406-002-0377-2](https://doi.org/10.1007/s00406-002-0377-2). URL: <https://link.springer.com/article/10.1007/s00406-002-0377-2>.
- [94] L. J. Siever, D. L. Murphy, S. Slater, E. de la Vega, and S. Lipper. "Plasma Prolactin Changes Following Fenfluramine in Depressed Patients Compared to Controls: An Evaluation of Central Serotonergic Responsibility in Depression". In: *Life Sciences* 34.11 (1984), pp. 1029–1039. ISSN: 0024-3205. DOI: [10.1016/0024-3205\(84\)90016-X](https://doi.org/10.1016/0024-3205(84)90016-X). URL: <https://www.sciencedirect.com/science/article/pii/002432058490016X>.
- [95] R. Navinés, E. Gómez-Gil, R. Martín-Santos, M. J. M. de Osaba, G. Escolar, and C. Gastó. "Hormonal Response to Buspirone Is Not Impaired in Major Depression". In: *Human Psychopharmacology: Clinical and Experimental* 22.6 (2007), pp. 389–395. ISSN: 1099-1077. DOI: [10.1002/hup.862](https://doi.org/10.1002/hup.862). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/hup.862>.

- [96] H. Y. Meltzer and M. Maes. "Effects of Buspirone on Plasma Prolactin and Cortisol Levels in Major Depressed and Normal Subjects". In: *Biological Psychiatry* 35.5 (1994), pp. 316–323. ISSN: 0006-3223. DOI: [10.1016/0006-3223\(94\)90035-3](https://doi.org/10.1016/0006-3223(94)90035-3). URL: <https://www.sciencedirect.com/science/article/pii/0006322394900353>.
- [97] F. Duval, M.-C. Mokrani, H. Correa, P. Bailey, M. Valdebenito, et al. "Lack of Effect of HPA Axis Hyperactivity on Hormonal Responses to D-Fenfluramine in Major Depressed Patients: Implications for Pathogenesis of Suicidal Behaviour". In: *Psychoneuroendocrinology* 26.5 (2001), pp. 521–537. ISSN: 0306-4530. DOI: [10.1016/S0306-4530\(01\)00011-7](https://doi.org/10.1016/S0306-4530(01)00011-7). URL: <https://www.sciencedirect.com/science/article/pii/S0306453001000117>.
- [98] H. McPherson, A. Walsh, and T. Silverstone. "Growth Hormone and Prolactin Response to Apomorphine in Bipolar and Unipolar Depression". In: *Journal of Affective Disorders* 76.1–3 (2003), pp. 121–125. ISSN: 0165-0327. DOI: [10.1016/S0165-0327\(02\)00077-0](https://doi.org/10.1016/S0165-0327(02)00077-0). PMID: 12943941.
- [99] J. Krogh, M. Nordentoft, M. Mohammad-Nezhad, and Å. Westrin. "Growth Hormone, Prolactin and Cortisol Response to Exercise in Patients with Depression". In: *Journal of Affective Disorders* 125.1–3 (2010), pp. 189–197. ISSN: 01650327. DOI: [10.1016/j.jad.2010.01.009](https://doi.org/10.1016/j.jad.2010.01.009). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0165032710000352>.
- [100] W. J. Cudala and J. Landowski. "Prolactin Response to Buspirone Is Not Impaired in Drug-Naïve First Episode Patients with Major Depressive Disorder". In: *Journal of Affective Disorders* 152–154 (2014), pp. 468–473. ISSN: 0165-0327. DOI: [10.1016/j.jad.2013.08.005](https://doi.org/10.1016/j.jad.2013.08.005). URL: <https://www.sciencedirect.com/science/article/pii/S0165032713005958>.
- [101] R. H. McAllister-Williams, H. A. Alhaj, A. Massey, J. Pankiv, and U. Reckermann. "Somatodendritic 5-hydroxytryptamine_{1A} (5-HT_{1A}) Autoreceptor Function in Major Depression as Assessed Using the Shift in Electroencephalographic Frequency Spectrum with Buspirone". In: *Psychological Medicine* 44.4 (2014), pp. 767–777. ISSN: 1469-8978. DOI: [10.1017/S0033291713001475](https://doi.org/10.1017/S0033291713001475). PMID: 23809646.
- [102] J. A. Monreal, F. Duval, M.-C. Mokrani, S. Fattah, and D. Palao. "Differences in Multihormonal Responses to the Dopamine Agonist Apomorphine between Unipolar and Bipolar Depressed Patients". In: *Journal of Psychiatric Research* 112 (2019), pp. 18–22. ISSN: 1879-1379. DOI: [10.1016/j.jpsychires.2019.02.009](https://doi.org/10.1016/j.jpsychires.2019.02.009). PMID: 30836201.
- [103] G. M. Asnis, J. Eisenberg, H. M. van Praag, C. Z. Lemus, J. M. Harkavy Friedman, and A. H. Miller. "The Neuroendocrine Response to Fenfluramine in Depressives and Normal Controls". In: *Biological Psychiatry* 24.1 (1988), pp. 117–120. ISSN: 0006-3223. DOI: [10.1016/0006-3223\(88\)90133-3](https://doi.org/10.1016/0006-3223(88)90133-3). URL: <https://www.sciencedirect.com/science/article/pii/0006322388901333>.
- [104] W. Verbeeck, M. Berk, J. Paiker, and B. Jersky. "The Prolactin Response to Sulpiride in Major Depression: The Role of the D₂ Receptor in Depression". In: *European Neuropsychopharmacology* 11.3 (2001), pp. 215–220. ISSN: 0924977X. DOI: [10.1016/S0924-977X\(01\)00086-4](https://doi.org/10.1016/S0924-977X(01)00086-4). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0924977X01000864>.
- [105] M. Maes, M. Vandewoude, C. Schotte, L. Maes, M. Martin, and P. Blockx. "Sex-Linked Differences in Cortisol, ACTH and Prolactin Responses to 5-Hydroxy-Tryptophan in Healthy Controls and Minor and Major Depressed Patients". In: *Acta Psychiatrica Scandinavica* 80.6 (1989), pp. 584–590. ISSN: 1600-0447. DOI: [10.1111/j.1600-0447.1989.tb03030.x](https://doi.org/10.1111/j.1600-0447.1989.tb03030.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1600-0447.1989.tb03030.x>.
- [106] P. Monteleone, A. La Rocca, A. Fuschino, F. Bortolotti, and M. Maj. "Seasonal Variation in Plasma Prolactin Response to D-Fenfluramine in Healthy Subjects". In: *Psychoneuroendocrinology* 24.2 (1999), pp. 201–208. ISSN: 0306-4530. DOI: [10.1016/S0306-4530\(98\)00074-2](https://doi.org/10.1016/S0306-4530(98)00074-2). URL: <https://www.sciencedirect.com/science/article/pii/S0306453098000742>.
- [107] G. L. Noel, R. C. Dimond, L. Wartofsky, J. M. Earll, and A. G. Frantz. "Studies of Prolactin and TSH Secretion by Continuous Infusion of Small Amounts of Thyrotropin-Releasing Hormone (TRH)". In: *The Journal of Clinical Endocrinology & Metabolism* 39.1 (1974), pp. 6–17. ISSN: 0021-972X. DOI: [10.1210/jcem-39-1-6](https://doi.org/10.1210/jcem-39-1-6). URL: <https://academic.oup.com/jcem/article-abstract/39/1/6/2685411>.
- [108] M. T. Buckman and G. T. Peak. "Estrogen Potentiation of Phenothiazine-Induced Prolactin Secretion in Man". In: *The Journal of Clinical Endocrinology & Metabolism* 37.6 (1973), pp. 977–980. ISSN: 0021-972X. DOI: [10.1210/jcem-37-6-977](https://doi.org/10.1210/jcem-37-6-977). URL: <https://academic.oup.com/jcem/article-abstract/37/6/977/2686299>.
- [109] G. Schwinn, A. von zur Mühlen, J. Köbberling, E. Halves, K. W. Wenzel, and H. Meinhold. "Plasma Prolactin Levels after TRH and Chlorpromazine in Normal Subjects and Patients with Impaired Pituitary Function". In: *Acta Endocrinologica* 79.4 (1975), pp. 663–676. ISSN: 0001-5598. DOI: [10.1530/acta.0.0790663](https://doi.org/10.1530/acta.0.0790663). PMID: 808064.
- [110] V. O'Keane, M. O'Hanlon, M. Webb, and T. Dinan. "D-Fenfluramine/Prolactin Response throughout the Menstrual Cycle: Evidence for an Oestrogen-Induced Alteration". In: *Clinical Endocrinology* 34.4 (1991), pp. 289–292. ISSN: 0300-0664. DOI: [10.1111/j.1365-2265.1991.tb03768.x](https://doi.org/10.1111/j.1365-2265.1991.tb03768.x). PMID: 1879060.

- [111] M. Reymond and Th. Lemarchand-Béraud. “Effects of Oestrogens on Prolactin and Thyrotrophin Responses to TRH in Women during the Menstrual Cycle and under Oral Contraceptive Treatment”. In: *Clinical Endocrinology* 5.5 (1976), pp. 429–437. ISSN: 0300-0664, 1365-2265. DOI: [10.1111/j.1365-2265.1976.tb01973.x](https://doi.org/10.1111/j.1365-2265.1976.tb01973.x). URL: <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2265.1976.tb01973.x>.
- [112] J. E. Tyson and H. G. Friesen. “Factors Influencing the Secretion of Human Prolactin and Growth Hormone in Menstrual and Gestational Women”. In: *American Journal of Obstetrics and Gynecology* 116.3 (1973), pp. 377–387. ISSN: 0002-9378. DOI: [10.1016/S0002-9378\(15\)31297-7](https://doi.org/10.1016/S0002-9378(15)31297-7). URL: <https://www.sciencedirect.com/science/article/pii/S0002937815312977>.
- [113] A. De Léan, L. Ferland, J. Drouin, P. Kelly, and F. Labrie. “Modulation of Pituitary Thyrotropin Releasing Hormone Receptor Levels by Estrogens and Thyroid Hormones”. In: *Endocrinology* 100.6 (1977), pp. 1496–1504. ISSN: 0013-7227. DOI: [10.1210/endo-100-6-1496](https://doi.org/10.1210/endo-100-6-1496). URL: <https://academic.oup.com/endo/article-abstract/100/6/1496/2617792>.
- [114] C. I. Messina, M. Malandri, G. Anifandis, K. Dafopoulos, P. Georgoulas, et al. “Submaximal Doses of Ghrelin Do Not Inhibit Gonadotrophin Levels but Stimulate Prolactin Secretion in Postmenopausal Women”. In: *Clinical Endocrinology* 87.1 (2017), pp. 44–50. ISSN: 1365-2265. DOI: [10.1111/cen.13349](https://doi.org/10.1111/cen.13349). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/cen.13349>.
- [115] F. Labrie, L. Ferland, F. Denizeau, and M. Beaulieu. “Sex Steroids Interact with Dopamine at the Hypothalamic and Pituitary Levels to Modulate Prolactin Secretion”. In: *Journal of Steroid Biochemistry* 12 (1980), pp. 323–330. ISSN: 0022-4731. DOI: [10.1016/0022-4731\(80\)90287-3](https://doi.org/10.1016/0022-4731(80)90287-3). URL: <https://www.sciencedirect.com/science/article/pii/0022473180902873>.
- [116] W. F. Stewart, R. B. Lipton, D. D. Celentano, and M. L. Reed. “Prevalence of Migraine Headache in the United States: Relation to Age, Income, Race, and Other Sociodemographic Factors”. In: *JAMA* 267.1 (1992), pp. 64–69. ISSN: 0098-7484. DOI: [10.1001/jama.1992.03480010072027](https://doi.org/10.1001/jama.1992.03480010072027). URL: <https://jamanetwork.com/journals/jama/article-abstract/394233>.
- [117] T. Victor, X. Hu, J. Campbell, D. Buse, and R. Lipton. “Migraine Prevalence by Age and Sex in the United States: A Life-Span Study”. In: *Cephalalgia* 30.9 (2010), pp. 1065–1072. ISSN: 0333-1024, 1468-2982. DOI: [10.1177/0333102409355601](https://doi.org/10.1177/0333102409355601). URL: <https://journals.sagepub.com/doi/10.1177/0333102409355601>.
- [118] R. Cevik, A. Gur, S. Acar, K. Nas, and A. J. Sarac. “Hypothalamic-Pituitary-Gonadal Axis Hormones and Cortisol in Both Menstrual Phases of Women with Chronic Fatigue Syndrome and Effect of Depressive Mood on These Hormones”. In: *BMC Musculoskeletal Disorders* 5.1 (2004), p. 47. ISSN: 1471-2474. DOI: [10.1186/1471-2474-5-47](https://doi.org/10.1186/1471-2474-5-47). URL: <https://link.springer.com/article/10.1186/1471-2474-5-47>.
- [119] J. Studd and N. Panay. “Chronic Fatigue Syndrome”. In: *The Lancet* 348.9038 (1996), p. 1384. ISSN: 0140-6736, 1474-547X. DOI: [10.1016/S0140-6736\(05\)65448-7](https://doi.org/10.1016/S0140-6736(05)65448-7). PMID: 8918295. URL: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(05\)65448-7/abstract](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(05)65448-7/abstract).
- [120] N. Thomas, S. J. K. A. Ubhayasekera, C. W. Armstrong, K. Huang, and J. Bergquist. “Steroid Dynamics in Myalgic Encephalomyelitis / Chronic Fatigue Syndrome: A Case-Control Study Using Ultra Performance Supercritical Fluid Chromatography Tandem Mass Spectrometry”. In: *Journal of Translational Medicine* 23.1 (2025), p. 829. ISSN: 1479-5876. DOI: [10.1186/s12967-025-06841-4](https://doi.org/10.1186/s12967-025-06841-4). URL: <https://link.springer.com/article/10.1186/s12967-025-06841-4>.
- [121] V. B. Wyller, J. A. Evang, K. Godang, K. K. Solhjell, and J. Bollerslev. “Hormonal Alterations in Adolescent Chronic Fatigue Syndrome”. In: *Acta Paediatrica* 99.5 (2010), pp. 770–773. ISSN: 1651-2227. DOI: [10.1111/j.1651-2227.2010.01701.x](https://doi.org/10.1111/j.1651-2227.2010.01701.x). URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1651-2227.2010.01701.x>.
- [122] V. V. Reddy, F. Naftolin, and K. J. Ryan. “Aromatization in the Central Nervous System of Rabbits: Effects of Castration and Hormone Treatment”. In: *Endocrinology* 92.2 (1973), pp. 589–594. ISSN: 0013-7227. DOI: [10.1210/endo-92-2-589](https://doi.org/10.1210/endo-92-2-589). PMID: 4682870.
- [123] C. J. Saldanha, K. A. Duncan, and B. J. Walters. “Neuroprotective Actions of Brain Aromatase”. In: *Frontiers in Neuroendocrinology* 30.2 (2009), pp. 106–118. ISSN: 00913022. DOI: [10.1016/j.yfrne.2009.04.016](https://doi.org/10.1016/j.yfrne.2009.04.016). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0091302209000107>.
- [124] A. J. Cleare, C. Messa, E. A. Rabiner, and P. M. Grasby. “Brain 5-HT_{1A} Receptor Binding in Chronic Fatigue Syndrome Measured Using Positron Emission Tomography and [11C]WAY-100635”. In: *Biological Psychiatry* 57.3 (2005), pp. 239–246. ISSN: 00063223. DOI: [10.1016/j.biopsych.2004.10.031](https://doi.org/10.1016/j.biopsych.2004.10.031). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0006322304011114>.
- [125] S. W. J. Lamberts and R. M. Macleod. “The Interaction of the Serotonergic and Dopaminergic Systems on Prolactin Secretion in the Rat: The Mechanism of Action of the “Specific” Serotonin Receptor Antagonist, Methysergide”. In: *Endocrinology* 103.1 (1978), pp. 287–295. ISSN: 0013-7227. DOI: [10.1210/endo-103-1-287](https://doi.org/10.1210/endo-103-1-287). URL: <https://academic.oup.com/endo/article-abstract/103/1/287/2618571>.

- [126] C. O. Lynch, M. D. Johnson, and W. R. Crowley. “Effects of the Serotonin Agonist, Quipazine, on Luteinizing Hormone and Prolactin Release: Evidence for Serotonin-Catecholamine Interactions”. In: *Life Sciences* 35.14 (1984), pp. 1481–1487. ISSN: 0024-3205. DOI: [10.1016/0024-3205\(84\)90165-6](https://doi.org/10.1016/0024-3205(84)90165-6). URL: <https://www.sciencedirect.com/science/article/pii/0024320584901656>.
- [127] S. C. Reddymasu, I. Soykan, and R. W. McCallum. “Domperidone: Review of Pharmacology and Clinical Applications in Gastroenterology”. In: *The American Journal of Gastroenterology* 102.9 (2007), pp. 2036–2045. ISSN: 0002-9270. DOI: [10.1111/j.1572-0241.2007.01255.x](https://doi.org/10.1111/j.1572-0241.2007.01255.x). PMID: [17488253](https://pubmed.ncbi.nlm.nih.gov/17488253/).
- [128] N. Shishioh-Ikejima, T. Ogawa, K. Yamaguti, Y. Watanabe, H. Kuratsune, and H. Kiyama. “The Increase of Alpha-Melanocyte-Stimulating Hormone in the Plasma of Chronic Fatigue Syndrome Patients.” In: *BMC neurology* 10 (2010), p. 73. ISSN: 1471-2377. DOI: [10.1186/1471-2377-10-73](https://doi.org/10.1186/1471-2377-10-73). PMID: [20731841](https://pubmed.ncbi.nlm.nih.gov/20731841/).
- [129] J. B. Hill, G. M. Nagy, and L. S. Frawley. “Suckling Unmasks the Stimulatory Effect of Dopamine on Prolactin Release: Possible Role for α -Melanocyte-Stimulating Hormone as a Mammatrope Responsiveness Factor”. In: *Endocrinology* 129.2 (1991), pp. 843–847. ISSN: 0013-7227, 1945-7170. DOI: [10.1210/endo-129-2-843](https://doi.org/10.1210/endo-129-2-843). URL: <https://academic.oup.com/endo/article-lookup/doi/10.1210/endo-129-2-843>.
- [130] L. Nuñez and L. S. Frawley. “ α -MSH Potentiates the Responsiveness of Mammatropes by Increasing Ca^{2+} Entry”. In: *American Journal of Physiology-Endocrinology and Metabolism* 274.6 (1998), E971–E977. ISSN: 0193-1849, 1522-1555. DOI: [10.1152/ajpendo.1998.274.6.E971](https://doi.org/10.1152/ajpendo.1998.274.6.E971). URL: <https://www.physiology.org/doi/10.1152/ajpendo.1998.274.6.E971>.
- [131] O. Khorram, J. C. Bedran De Castro, and S. M. McCann. “Physiological Role of Alpha-Melanocyte-Stimulating Hormone in Modulating the Secretion of Prolactin and Luteinizing Hormone in the Female Rat.” In: *Proceedings of the National Academy of Sciences* 81.24 (1984), pp. 8004–8008. ISSN: 0027-8424, 1091-6490. DOI: [10.1073/pnas.81.24.8004](https://doi.org/10.1073/pnas.81.24.8004). URL: <https://pnas.org/doi/full/10.1073/pnas.81.24.8004>.
- [132] C. B. Newman, S. L. Wardlaw, and A. G. Frantz. “Suppression of Basal and Stress-Induced Prolactin Release and Stimulation of Luteinizing Hormone Secretion by α -Melanocyte-Stimulating Hormone”. In: *Life Sciences* 36.17 (1985), pp. 1661–1668. ISSN: 00243205. DOI: [10.1016/0024-3205\(85\)90369-8](https://doi.org/10.1016/0024-3205(85)90369-8). URL: <https://linkinghub.elsevier.com/retrieve/pii/0024320585903698>.
- [133] C. Pasqualini, F. Bojda, and B. Kerdellhué. “Direct Effect of Estradiol on the Number of Dopamine Receptors in the Anterior Pituitary of Ovariectomized Rats”. In: *Endocrinology* 119.6 (1986), pp. 2484–2489. ISSN: 0013-7227. DOI: [10.1210/endo-119-6-2484](https://doi.org/10.1210/endo-119-6-2484). URL: <https://academic.oup.com/endo/article-abstract/119/6/2484/2540572>.
- [134] P. Jakeman, J. Hawthorne, S. Maxwell, M. Kendall, and G. Holder. “Evidence for Downregulation of Hypothalamic 5-hydroxytryptamine Receptor Function in Endurance-trained Athletes”. In: *Experimental Physiology* 79.3 (1994), pp. 461–464. ISSN: 0958-0670, 1469-445X. DOI: [10.1113/expphysiol.1994.sp003780](https://doi.org/10.1113/expphysiol.1994.sp003780). URL: <https://physoc.onlinelibrary.wiley.com/doi/10.1113/expphysiol.1994.sp003780>.
- [135] F. A. Cadegiani and C. E. Kater. “Hormonal Response to a Non-Exercise Stress Test in Athletes with Overtraining Syndrome: Results from the Endocrine and Metabolic Responses on Overtraining Syndrome (EROS) – EROS-STRESS”. In: *Journal of Science and Medicine in Sport* 21.7 (2018), pp. 648–653. ISSN: 1440-2440, 1878-1861. DOI: [10.1016/j.jsams.2017.10.033](https://doi.org/10.1016/j.jsams.2017.10.033). PMID: [29157780](https://pubmed.ncbi.nlm.nih.gov/29157780/). URL: [https://www.jsams.org/article/S1440-2440\(17\)31746-2/abstract](https://www.jsams.org/article/S1440-2440(17)31746-2/abstract).
- [136] A. Broocks, T. Meyer, A. George, U. Hillmer-Vogel, D. Meyer, et al. “Decreased Neuroendocrine Responses to Meta-Chlorophenylpiperazine (m-CPP) but Normal Responses to Ipsapirone in Marathon Runners”. In: *Neuropsychopharmacology* 20.2 (1999), pp. 150–161. ISSN: 1740-634X. DOI: [10.1016/S0893-133X\(98\)00056-6](https://doi.org/10.1016/S0893-133X(98)00056-6). URL: <https://www.nature.com/articles/1395239>.
- [137] A. T. Strachan and R. J. Maughan. “The Hormonal Response to a D-Fenfluramine Challenge in Trained and Sedentary Men.” in: *Medicine & Science in Sports & Exercise* 31.4 (1999), pp. 547–553. ISSN: 0195-9131. DOI: [10.1097/00005768-199904000-00009](https://doi.org/10.1097/00005768-199904000-00009). URL: <https://www.ovid.com/00005768-199904000-00009>.
- [138] D. Dwyer and J. Flynn. “Short Term Aerobic Exercise Training in Young Males Does Not Alter Sensitivity to a Central Serotonin Agonist”. In: *Experimental Physiology* 87.1 (2002), pp. 83–89. ISSN: 0958-0670. DOI: [10.1113/eph8702176](https://doi.org/10.1113/eph8702176). PMID: [11805862](https://pubmed.ncbi.nlm.nih.gov/11805862/).
- [139] P. A. Thompson, M. Maes, and H. Y. Meltzer. “Effect of the Placebo Control Condition in Neuroendocrine Challenge Studies”. In: *Psychiatry Research* 52.3 (1994), pp. 317–326. ISSN: 0165-1781. DOI: [10.1016/0165-1781\(94\)90077-9](https://doi.org/10.1016/0165-1781(94)90077-9). PMID: [7991724](https://pubmed.ncbi.nlm.nih.gov/7991724/).
- [140] A.-K. Lennartsson and I. H. Jonsdottir. “Prolactin in Response to Acute Psychosocial Stress in Healthy Men and Women”. In: *Psychoneuroendocrinology* 36.10 (2011), pp. 1530–1539. ISSN: 0306-4530. DOI: [10.1016/j.psyneuen.2011.04.007](https://doi.org/10.1016/j.psyneuen.2011.04.007). URL: <https://www.sciencedirect.com/science/article/pii/S0306453011001314>.

- [141] R. R. Gala. “The Physiology and Mechanisms of the Stress-Induced Changes in Prolactin Secretion in the Rat”. In: *Life Sciences* 46.20 (1990), pp. 1407–1420. ISSN: 0024-3205. DOI: [10.1016/0024-3205\(90\)90456-2](https://doi.org/10.1016/0024-3205(90)90456-2). URL: <https://www.sciencedirect.com/science/article/pii/0024320590904562>.
- [142] R. E. Gammans, R. F. Mayol, and J. A. Labudde. “Metabolism and Disposition of Buspirone”. In: *The American Journal of Medicine*. Pharmacology, Efficacy, and Safety of a Non-Benzodiazepine Anxiolytic 80 (3, Supplement 2 1986), pp. 41–51. ISSN: 0002-9343. DOI: [10.1016/0002-9343\(86\)90331-1](https://doi.org/10.1016/0002-9343(86)90331-1). URL: <https://www.sciencedirect.com/science/article/pii/0002934386903311>.
- [143] L. Sevinçok and A. Erol. “The Prolactin Response to Buspirone in Poststroke Depression: A Preliminary Report”. In: *Journal of Affective Disorders* 59.2 (2000), pp. 169–173. ISSN: 0165-0327. DOI: [10.1016/S0165-0327\(99\)00143-3](https://doi.org/10.1016/S0165-0327(99)00143-3). URL: <https://www.sciencedirect.com/science/article/pii/S0165032799001433>.
- [144] A. L. Komaroff and W. I. Lipkin. “ME/CFS and Long COVID Share Similar Symptoms and Biological Abnormalities: Road Map to the Literature”. In: *Frontiers in Medicine* 10 (2023). ISSN: 2296-858X. DOI: [10.3389/fmed.2023.1187163](https://doi.org/10.3389/fmed.2023.1187163). URL: <https://www.frontiersin.org/journals/medicine/articles/10.3389/fmed.2023.1187163/full>.
- [145] M. Hulens, W. Dankaerts, R. Rasschaert, F. Bruyninckx, P. De Mulder, and C. Bervoets. “The Link Between Empty Sella Syndrome, Fibromyalgia, and Chronic Fatigue Syndrome: The Role of Increased Cerebrospinal Fluid Pressure”. In: *Journal of Pain Research* Volume 16 (2023), pp. 205–219. ISSN: 1178-7090. DOI: [10.2147/JPR.S394321](https://doi.org/10.2147/JPR.S394321). URL: <https://www.tandfonline.com/doi/full/10.2147/JPR.S394321>.
- [146] F. Duval, M.-C. Mokrani, and M.-A. Crocq. “What Future for Neuroendocrinology in Psychiatry?” In: *Psychoneuroendocrinology* 38.8 (2013), pp. 1213–1219. ISSN: 0306-4530. DOI: [10.1016/j.psyneuen.2013.04.012](https://doi.org/10.1016/j.psyneuen.2013.04.012). URL: <https://www.sciencedirect.com/science/article/pii/S0306453013001509>.
- [147] “Anti-Obesity Drugs Recalled from Global Market”. In: *The Lancet* 350.9081 (1997), p. 867. ISSN: 0140-6736, 1474-547X. DOI: [10.1016/S0140-6736\(97\)23038-2](https://doi.org/10.1016/S0140-6736(97)23038-2). URL: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(97\)23038-2/abstract](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(97)23038-2/abstract).

Acknowledgments

Thank you to members of the [Science for ME](#) forum for help with finding and analyzing studies related to prolactin response, as well as for helping me to nurture my interest in rigorous science.

Updates

- Version 1 — August 20, 2026.
- Version 2 — August 21, 2026: Fixed citation for sentence about “increased conversion of androgens to estrogens within the hypothalamus”. Minor clarifications and style improvements.
- Version 3 — August 27, 2026: Added tables, sections about confounding and neuroendocrine challenge testing, expanded conclusion, and made other minor style improvements.
- Version 4 — August 29, 2026: Added email address, improved figure, and other minor improvements.
- Version 5 — September 4, 2026: Significant update included improving structure and style, adding abstract and expanding introduction, improving figure, and adding additional relevant citations.
- Version 6 — September 5, 2026: Minor typo and style fixes. Added limitations section.
- Version 7 — September 19, 2026: Minor style and clarity improvements. Noted matching control data for two ME/CFS studies. Replaced one citation for blunted prolactin response in depression (replaced review with original research) and added citations for normal and increased response in depression. Improved description of Duval et al.’s claims in conclusion. Added speculation that brain damage may underlie prolactin response findings. Added note about fenfluramine being removed from market. Re-calculated prolactin response values using higher resolution plot images, while also correcting the response value for Sharpe et al., 1996, as the wrong timepoint for baseline was previously used.

See all versions at <https://github.com/ruvilonix/prolactin-response/>.