

Abnormal prolactin response in ME/CFS

Introduction

Abnormal prolactin response in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a consistently replicated finding, but the implications of this finding have only been minimally explored. Further research into this phenomenon could potentially illuminate whether hormonal, neural, or other pathways are involved in the pathophysiology of ME/CFS.

Prolactin

Prolactin, a protein encoded by a gene on chromosome 6, is produced within several tissues in the human body, but its main site of synthesis is the anterior pituitary gland. Prolactin is best known for its essential role in the development of mammary tissue and its ability to stimulate lactation. Additionally, prolactin exerts numerous other effects, which include the promotion of parental behavior, enhancement of immunity, modulation of lipid and glucose metabolism, and inhibition of the reproductive cycle during lactation. Suckling is the most potent natural stimulant of prolactin secretion, however stress, sounds, and circadian light patterns also affect the release of prolactin. (1,2)

Unlike most hormones of the pituitary gland, prolactin is primarily regulated through inhibition instead of stimulation, whereby pituitary prolactin-secreting cells (lactotrophs) naturally secrete prolactin when hypothalamic influence is blocked. The main hormone involved in inhibiting prolactin is dopamine, which is released by neurons of the arcuate nucleus and periventricular nucleus of the hypothalamus, and which travels through the long or short portal veins to reach the lactotrophs. However, several other hormones appear to also have the ability to directly stimulate prolactin release by lactotrophs, and these include, but are not limited to, thyrotropin-releasing hormone (TRH), vasopressin, oxytocin, and vasoactive intestinal peptide (VIP). (2)

Prolactin response to buspirone in ME/CFS

Strikingly, every single study that has tested prolactin response to buspirone in ME/CFS has reported that patients exhibit an abnormally large prolactin release about one to two hours after administration.

Buspirone is marketed as a treatment for generalized anxiety disorder, though unlike classical anxiety medications such as benzodiazepines, buspirone has no activity at GABA receptors. Instead, buspirone is an agonist of 5HT_{1A} receptors, where it exhibits strong binding affinity, and also acts as an antagonist at D₂ dopamine receptors, albeit with weaker binding affinity. (3)

The first study of prolactin response in ME/CFS came from Bakheit et al. in 1992. The findings from this study were reported both in Bakheit's thesis (4), as well as in a peer-reviewed paper (5). The authors were studying postviral fatigue syndrome (PVFS), whereby the criteria used for the study closely resembled the chronic fatigue syndrome criteria proposed by Holmes et al. (6) and Sharpe et al. (7). The criteria also required that the fatigue was preceded by a probable or definite viral infection, and was made worse by exercise and did not improve after bed rest. The authors found that the patient group had a larger increase in plasma prolactin after administration of buspirone, when the patients were compared to healthy controls or individuals with depression. The same effect was observed in both males and females.

Figure 1 illustrates the study results using the individual level data provided in Bakheit's thesis. Notably, and as seen in virtually all other studies of this topic, baseline prolactin was not significantly altered in the patient group.




Figure 1: Prolactin response, based on data from Bakheit 1992. PVFS: Postviral fatigue syndrome

Further evidence was provided in a study from John Richardson in 1995. In this study, patients who fulfilled the CDC and Oxford criteria for ME/CFS were compared to family members without ME/CFS. With prolactin response defined as the ratio of prolactin levels after to before buspirone administration, the response to buspirone (50 mg, orally) in the patients was, on average, over 3 times higher than that of the controls. The study also found a correlation between prolactin response in patients and their degree of sleep disturbance (8). A later study from Richardson and Costa again suggested very high prolactin responses to buspirone (50 mg, orally) in ME/CFS patients, however this study did not include a control group, and it is unclear if high prolactin response was a criteria for inclusion in the study (9).

In 1996, Sharpe et al. reported increased prolactin response to buspirone (0.5 mg/kg orally, up to 45 mg

in patients fulfilling Oxford criteria for ME/CFS (7), but who did not also exhibit symptoms of depression. Notably, plasma levels of buspirone and its metabolite, 1-(2-Pyrimidinyl)piperazine (1-PP), were not significantly different between groups after administration of buspirone, suggesting that the difference in prolactin response was not caused by differences in drug metabolism. The growth hormone response to buspirone did not significantly differ between groups (10).

Also in 1996, Tahir Majeed, under the supervision of Peter O. Behan, published a thesis testing hormonal responses to a variety of drug challenges. The study cohort included patients who fulfilled Fukuda (11) and Oxford (7) criteria, and who had fatigue which worsened after exercise. The

control group was made up of healthy, sex-matched volunteers. The ME/CFS patients exhibited a significantly larger increase in prolactin after administration of buspirone (60 mg, orally). Further findings in ME/CFS patients included a normal growth hormone response to bromocriptine and baclofen, a larger growth hormone response to pyridostigmine, smaller growth hormone responses to desipramine and dexamethasone, and a smaller adrenocorticotrophic hormone (ACTH) response to ipsapirone (12).

Another 1996 paper, authored by Behan, reported increased prolactin response to buspirone (60 mg, orally) in individuals with an ME/CFS-like illness (including fatigue worsened by exercise, myalgia, and with onset after a flu-like illness) and who had previously been exposed to organophosphates (13). Note that while this paper indicates that the study

included 10 male healthy controls, later court testimony from Behan in an organophosphate trial indicated that there was a mistake in the paper, and the actual control group was made up of 15 males and 15 females (14). However, as Behan noted in court, the inclusion of females in the control group would not be expected to lead to spurious differences between groups, and would instead more likely have led to a smaller difference, as females have larger prolactin responses to buspirone than males (15).

Additionally, two studies from Racciatti et al. suggest that the authors have observed further evidence of increased prolactin response to buspirone in ME/CFS. However one study did not include a healthy control group (16), and the other was reported in an abstract which we have not been able to access (17), and with the findings briefly mentioned in a later review (18).

Finally, a 2001 case report included details of a female patient with ME/CFS, according to Oxford criteria (7), who exhibited an abnormally large prolactin response to buspirone (30 mg, orally). After symptom improvement, which followed a graded exercise therapy intervention, the prolactin response was measured again on two occasions, and was found to be normal both times (19).

Prolactin response to other challenges in ME/CFS

Additional evidence of abnormal prolactin response is provided by studies which tested fenfluramine, exercise, and insulin-induced hypoglycemia in ME/CFS patients. Negative or conflicting results have been observed after administration of m-chlorophenylpiperazine (mCPP) and tryptophan.

In 1995, Bearn et al. tested the prolactin response in ME/CFS to two different neuroendocrine challenges. The serotonin agonist, d-fenfluramine, was tested, but prolactin response did not significantly differ between groups. On the other hand, inducing hypoglycemia through administration of insulin led to a significantly *smaller* increase of prolactin in the ME/CFS group (20).

Ottenweller et al. tested a treadmill-based maximal exercise challenge, and again reported a significantly smaller increase of prolactin in ME/CFS patients (21).

Yatham et al. published a study of response to racemic fenfluramine in ME/CFS. In this study, the groups did not significantly differ in the magnitude of prolactin response (22).

Two later studies from separate groups tested the prolactin response to d-fenfluramine, and both of these studies reported *increased* prolactin response in the ME/CFS patients (23,24).

A 2001 study did not find a significant difference between ME/CFS patients and controls in prolactin response to the 5-HT_{2c} receptor agonist, mCPP (25).

Finally, a 2010 study reported a significantly increased prolactin response to tryptophan, but only in females with ME/CFS, and not in females with ME/CFS+fibromyalgia, or in either of the male patient groups (26).

Other conditions

Abnormally increased prolactin response has been reported in several other conditions, such as migraine, irritable bowel syndrome (IBS), hormonal disorders, and gonadal disorders.

Migraine

Abnormal prolactin response has been reported several times in migraine. Migraine patients may have an abnormally long return to baseline prolactin after administration of reserpine (seen in males) (27) or benserazide (seen in females) (28). Studies have reported abnormally increased prolactin response in migraine patients to the D₂ antagonists sulpiride (28,29) and domperidone (29). An abnormally large *decrease* in prolactin has been noted after administration of L-deprenyl (30). In contrast, an abnormally *small* decrease was reported after administration of the dopamine reuptake inhibitor, nomifensine (29) and the dopamine precursor, L-DOPA (31). Cassidy et al. have published two studies showing increased prolactin response to buspirone in

migraine (32,33). The prolactin response to mCPP was larger in a study that used a dosage of 0.5 mg/kg (34), but not in a study that used a smaller 0.25 mg/kg dosage of the drug (35). A larger prolactin response was noted after administration of racemic fenfluramine (36).

Several studies have tested TRH challenges in migraine. A blunted prolactin response in migraine was seen after administration of 50 µg and 200 µg doses of TRH (37). Another study found an increased prolactin response to TRH during migraine attacks (38). Awaki et al. found an abnormally increased prolactin response after administration of a cocktail which included TRH, gonadotropin-releasing hormone (GnRH), and insulin (39).

However, other studies of migraine did not find significant differences between groups, including after administration of domperidone (40), fenfluramine (28), lisuride (29), metaclopramide (41,42), morphine (43), piribedil (44), and sumatriptan (45).

Other disorders

Increased prolactin response has been reported in only a small number of other conditions. The following section includes some examples of these, but this should not be considered exhaustive.

Several studies from the same research group have reported abnormally increased prolactin response to buspirone in IBS (46), and in non-ulcer dyspepsia (47–50).

An increased prolactin response was seen in primary testicular failure after administration of TRH (51) and metoclopramide (52). Prolactin response to TRH was also significantly larger in females with primary ovarian failure (53). An increased prolactin response to TRH was also reported in patients with polyendocrine metabolic ovarian syndrome (PMOS) (54).

Additionally, studies have reported observing health conditions in which patients demonstrated *blunted* prolactin responses. For example, the prolactin response to buspirone or fenfluramine has been

observed to be blunted in depression in some (23,55), but not all studies (56).

Connection to sex hormones?

As females are more likely than males to have ME/CFS (57), it may be useful to consider whether sex hormones influence prolactin response in a way that may explain the observations of increased prolactin response to buspirone and d-fenfluramine in ME/CFS.

Females have a higher prolactin response to buspirone than males (15). Similar sex differences have been observed after administration of thyrotropin-releasing hormone (TRH), phenothiazine, and possibly chlorpromazine (58–60).

Prolactin response to various drugs differs throughout the menstrual cycle. While baseline prolactin levels stayed relatively steady throughout the menstrual cycle, it was observed that prolactin response to d-fenfluramine was highest at mid-cycle, followed by luteal, then follicular phases. This mirrored the differing amounts of circulating estradiol present across the menstrual cycle (61). Prolactin response to buspirone in females was found to be larger during the luteal phase than during the follicular phase or mid-cycle (15). Prolactin response to TRH may be somewhat larger during the follicular phase (62), but findings are inconsistent (63).

Exogenous estradiol can modulate prolactin response. TRH-induced prolactin response is substantially larger in females taking a short course of exogenous estradiol, as well as after subsequent short term combined contraceptive use, but not in those taking long term combined contraceptives (62). In male rats, a prolactin response to TRH was only detectable when the rats had previously been administered estradiol (64). Exogenous estradiol also causes an increased prolactin response to ghrelin in postmenopausal women (65). Further, administration of estradiol to ovariectomized rats caused substantially larger prolactin responses to the dopamine antagonist, thioproperazine. (66)

Dinan et al. also reported that females had more symptoms of sedation in response to buspirone during the luteal phase than at other timepoints, with this being the timepoint where prolactin response was found to be highest (15). This may relate to the finding that ME/CFS patients appear to have more buspirone-induced side effects, such as fatigue and nausea, when compared to healthy controls (5,8).

Like ME/CFS (57), migraine appears to be more prevalent in females than males (67,68). As noted in the “Migraine” section, this is one of the few other health conditions in which several studies have reported increased prolactin response to various neuroendocrine challenges. These commonalities may suggest similar underlying pathways related to sex hormones in these health conditions.

An important point to keep in mind is that a hypothesis which ties estradiol to the abnormal prolactin response seen in ME/CFS would need to be explained by a mechanism which is independent of blood estradiol levels, as studies have shown that blood levels of this hormone do not appear to be significantly increased in ME/CFS (69–72). One alternative possibility was proposed in a study of increased prolactin response to TRH in primary ovarian failure, where the authors speculated that the patients might have increased conversion of androgens to estrogens within the hypothalamus (73). This was not based on direct evidence, so should be interpreted cautiously, but it may be worth considering this possibility.

One could test the estradiol hypothesis by measuring the prolactin response to buspirone in ME/CFS and healthy controls both before and after administration of estrogen receptor antagonists or aromatase inhibitors. If the difference between groups were found to be diminished after blocking the influence of estrogens, this would provide evidence of sex hormone involvement in the abnormal prolactin response. It may be necessary to prescribe an extended course of these drugs prior to the second prolactin challenge, as estrogens are capable of inducing long term changes in prolactin-secreting cells, for example

through genomic regulation or by increasing the number of lactotrophs, as reviewed in (1).

Where to go from here?

The implications of abnormal prolactin response in ME/CFS are far from clear. While several of the ME/CFS prolactin response studies suggested that the findings were evidence of abnormalities in the serotonin system (5,12,23,24), there is too little direct evidence for this hypothesis to make strong conclusions. Indeed, a PET study of serotonin receptors in ME/CFS resulted in the opposite of what the authors had expected based on prior prolactin response findings (they found downregulation instead of the upregulation of 5-HT_{1A} receptors) (74). While abnormal prolactin response could involve the receptors directly affected by the neuroendocrine probes, such as 5-HT_{1A}, this phenomenon could instead be highlighting any of a number of possible downstream abnormalities, such as altered dopaminergic tone or sensitized lactotrophs. For example, serotonin-related drugs, which were used in most of the ME/CFS studies with significant results, are unlikely to be able to directly stimulate lactotrophs (75), and may instead exert at least part of their influence on lactotrophs through altering hypothalamic dopamine release (76).

What is clear is the importance of further exploration of these findings. ME/CFS is a condition with notoriously few well-replicated, specific findings. Some of other relatively replicated findings include female sex bias (57) and decreased NK cell cytotoxicity (77), but these findings have not yet led to usable insights into pathophysiology or treatment development. With so few clues into the underlying pathophysiology, it is important to thoroughly investigate every available avenue.

A straightforward next step would be testing prolactin response to other neuroendocrine probes. For example, if dopaminergic drugs and TRH both lead to abnormal prolactin responses, this would suggest abnormally sensitized lactotrophs, as dopamine and TRH both directly affect lactotrophs (2). On the other hand, if

neither of these probes demonstrates altered responses, but responses to serotonin-related drugs are altered, this may indicate upstream abnormalities. It may be most fruitful to consider testing the probes which have already been reported to cause increased prolactin response in other health conditions, such as the D2 antagonist domperidone, which has demonstrated an increased prolactin response in one study of migraine (29), and which can modulate prolactin release from the pituitary gland without appreciably crossing the blood-brain barrier (78), thus limiting the potential for confounding due to central nervous system effects.

One ME/CFS study has reported increased levels of alpha-melanocyte-stimulating hormone (α -MSH), albeit with substantial overlap between groups (79). α -MSH is known to alter prolactin response, both in vitro and in vivo in non-human animals (80–83). Thus, it may be worthwhile to attempt to replicate increased α -MSH in ME/CFS, and to test for a correlation to prolactin response in these patients.

It may also be possible to explore this finding further using PET scans, for example by measuring dopamine receptor D2 (DRD2) binding in the pituitary gland. If the ME/CFS abnormality does relate to sex hormones, then DRD2 levels may be altered, as is seen when estradiol is applied to anterior pituitary cells *in vitro* (73).

It is not clear why research into abnormal prolactin response in ME/CFS has appeared to stall, with almost all studies of this phenomenon being conducted in the 1990s, and with no studies directly testing prolactin response in ME/CFS in over 15 years. Through careful consideration of study design and consultation with experts in endocrinology and neurology, future research into prolactin response may uncover important insights about the pathophysiology of ME/CFS, a condition that has, so far, eluded explanation.

References

1. Grattan DR. 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-prolactin axis. *Journal of Endocrinology*. 2015 Aug;226(2):T101–22. doi:10.1530/JOE-15-0213
2. Freeman ME, Kanyicska B, Lerant A, Nagy G. Prolactin: Structure, Function, and Regulation of Secretion. *Physiological Reviews*. 2000 Jan 10;80(4):1523–631. doi:10.1152/physrev.2000.80.4.1523
3. Loane C, Politis M. Buspirone: What is it all about? *Brain Research*. 2012 Jun 21;1461:111–8. doi:10.1016/j.brainres.2012.04.032
4. Bakheit AMO. Hypothalamic Dysfunction and Neurotransmitter Abnormalities in the Postviral Fatigue Syndrome [MD] [Internet]. [Ann Arbor :]: ProQuest Dissertations & Theses,; 1992 [cited 2026 Jul 21]. Available from: <https://theses.gla.ac.uk/75589/>
5. Bakheit AM, Behan PO, Dinan TG, Gray CE, O’Keane V. Possible upregulation of hypothalamic 5-hydroxytryptamine receptors in patients with postviral fatigue syndrome. *BMJ*. 1992 Apr 18;304(6833):1010–2. PubMed PMID: 1586780; PubMed Central PMCID: PMC1881733.

6. Holmes GP, Kaplan JE, Gantz NM, Komaroff AL, Schonberger LB, Straus SE, et al. Chronic fatigue syndrome: a working case definition. *Ann Intern Med.* 1988 Mar;108(3):387–9. doi:10.7326/0003-4819-108-3-387 PubMed PMID: 2829679.
7. Sharpe MC. A Report—Chronic Fatigue Syndrome: Guidelines for Research. *J R Soc Med.* 1991 Feb 1;84(2):118–21. doi:10.1177/014107689108400224
8. Richardson J. Disturbance of Hypothalamic Function and Evidence for Persistent Enteroviral Infection in Patients with Chronic Fatigue Syndrome. *Journal Of Chronic Fatigue Syndrome.* 1995 Jan 1;1(2):59–66. doi:10.1300/J092v01n02_05
9. Richardson J, Costa DC. Relationship Between SPECT Scans and Buspirone Tests in Patients with ME/CFS. *Journal of Chronic Fatigue Syndrome.* 1998 Jan;4(3):23–38. doi:10.1300/J092v04n03_04
10. Sharpe M, Clements A, Hawton K, Young AH, Sargent P, Cowen PJ. Increased prolactin response to buspirone in chronic fatigue syndrome. *Journal of Affective Disorders.* 1996 Nov 4;41(1):71–6. doi:10.1016/0165-0327(96)00075-4
11. Fukuda K, Straus SE, Hickie I, Sharpe MC, Dobbins JG, Komaroff A. The chronic fatigue syndrome: a comprehensive approach to its definition and study. International Chronic Fatigue Syndrome Study Group. *Ann Intern Med.* 1994 Dec 15;121(12):953–9. doi:10.7326/0003-4819-121-12-199412150-00009 PubMed PMID: 7978722.
12. Majeed T. Neuroendocrine Alterations in Chronic Fatigue Syndrome [PhD] [Internet]. [Ann Arbor :]: ProQuest Dissertations & Theses,; 1996 [cited 2026 May 15]. Available from: <https://theses.gla.ac.uk/75796/>
13. Behan PO. Chronic Fatigue Syndrome as a Delayed Reaction to Chronic Low-dose Organophosphate Exposure. *Journal of Nutritional & Environmental Medicine.* 1996 Jan 1;6(4):341–50. doi:10.3109/13590849609007262
14. The Herald [Internet]. 1997 [cited 2026 Aug 7]. Professor’s future in doubt. University inquiry blames academic’s sloppy preparation for court fiasco. Available from: <https://www.heraldsotland.com/news/12294532.professors-future-in-doubt-university-inquiry-blames-academics-sloppy-preparation-for-court-fiasco/>
15. Dinan TG, Barry S, Yatham LN, Mobayed M, O’Hanlon M. The reproducibility of the prolactin response to buspirone: relationship to the menstrual cycle. *Int Clin Psychopharmacol.* 1990 Apr;5(2):119–23. doi:10.1097/00004850-199004000-00006 PubMed PMID: 2380543; PubMed Central PMCID: 2380543.
16. Racciatti D, Vecchiet J, Ceccomancini A, Ricci F, Pizzigallo E. Chronic fatigue syndrome following a toxic exposure. *Science of The Total Environment.* 2001 Apr 10;Environment, Human Health and Immune System (selected papers from International Symposium on Occupational and Environmental Allergy and Immune Diseases,Chieti(Italy),April 27-30,1999270(1):27–31. doi:10.1016/S0048-9697(00)00777-4

17. Racciatti D, Barberio A, Pizzigallo E, others. Neuroendocrine perspective of chronic fatigue syndrome. *Journal of Musculoske Pain*. 1998;6(S2):152.
18. Racciatti D, Barberio A, Vecchiet J, Pizzigallo E. Clinical and Pathogenetical Characterization of 238 Patients of a Chronic Fatigue Syndrome Italian Center. *Journal Of Chronic Fatigue Syndrome*. 1999 Jan 1;5(3–4):61–70. doi:10.1300/J092v05n03_05
19. Sharma A, Oyeboode F, Kendall MJ, Jones DA. Recovery from chronic fatigue syndrome associated with changes in neuroendocrine function. *J R Soc Med*. 2001 Jan;94(1):26–7. doi:10.1177/014107680109400107
20. Bearn J, Allain T, Coskeran P, Munro N, Butler J, McGregor A, et al. Neuroendocrine responses to d-fenfluramine and insulin-induced hypoglycemia in chronic fatigue syndrome. *Biological Psychiatry*. 1995 Feb;37(4):245–52. doi:10.1016/0006-3223(94)00121-I
21. Ottenweller JE, Sisto SA, McCarty RC, Natelson BH. Hormonal Responses to Exercise in Chronic Fatigue Syndrome. *Neuropsychobiology*. 2001;43(1):34–41. doi:10.1159/000054863
22. Yatham LN, Morehouse RL, Chisholm BT, Haase DA, MacDonald DD, Marrie TJ. Neuroendocrine assessment of serotonin (5-HT) function in chronic fatigue syndrome. *Can J Psychiatry*. 1995 Mar;40(2):93–6. doi:10.1177/070674379504000207 PubMed PMID: 7788624.
23. Cleare AJ, Bearn J, Allain T, McGregor A, Wessely S, Murray RM, et al. Contrasting neuroendocrine responses in depression and chronic fatigue syndrome. *Journal of Affective Disorders*. 1995 Aug 18;34(4):283–9. doi:10.1016/0165-0327(95)00026-J
24. Sharpe M, Hawton K, Clements A, Cowen P. Increased brain serotonin function in men with chronic fatigue syndrome. *BMJ*. 1997 Jul 19;315(7101):164–5. doi:10.1136/bmj.315.7101.164
25. Vassallo CM, Feldman E, Peto T, Castell L, Sharpley AL, Cowen PJ. Decreased tryptophan availability but normal post-synaptic 5-HT_{2c} receptor sensitivity in chronic fatigue syndrome. *Psychol Med*. 2001 May;31(4):585–91. doi:10.1017/S0033291701003580
26. Weaver SA, Janal MN, Aktan N, Ottenweller JE, Natelson BH. Sex Differences in Plasma Prolactin Response to Tryptophan in Chronic Fatigue Syndrome Patients With and Without Comorbid Fibromyalgia. *Journal of Women's Health*. 2010 May 1;19(5):951–8. doi:10.1089/jwh.2009.1697
27. Nappi G, Savoldi F, Bono G, Martignoni E. Reserpine - Headache and PRL Release in Migraine. *Headache: The Journal of Head and Face Pain*. 1979;19(5):273–7. doi:10.1111/j.1526-4610.1979.hed1905273.x
28. Polleri A, Nappi G, Masturzo P, Martignoni E, Murialdo G, Bono G, et al. Neuroendocrine approach to headache. *Advances in neurology*. 1982;33:173–82. Available from: <https://archive.org/details/headachephysiopa0000unse/page/172/mode/2up/>
29. Murialdo G, Martignoni E, Maria AD, Bonura ML, Sances G, Bono G, et al. Changes in the Dopaminergic Control of Prolactin Secretion and in Ovarian Steroids in Migraine. *Cephalalgia*. 1986 Mar 1;6(1):43–9. doi:10.1046/j.1468-2982.1986.0601043.x

30. Calabresi P, Silvestrini M, Stratta F, Cupini L, Argiro G, Atzei G, et al. L-Deprenyl test in migraine: neuroendocrinological aspects. *Cephalalgia*. 1993;13(6):406–9. doi:10.1046/j.1468-2982.1993.1306406.x
31. Nattero G, Corno M, Savi L, Isaia G c., Priolo C, Mussetta M. Prolactin and Migraine: Effect of L-Dopa on Plasma Prolactin Levels in Migraineurs and Normals. *Headache: The Journal of Head and Face Pain*. 1986;26(1):9–12. doi:10.1111/j.1526-4610.1986.hed2601009.x
32. Cassidy EM, Tomkins E, Dinan T, Hardiman O, O’Keane V. Central 5-HT receptor hypersensitivity in migraine without aura. *Cephalalgia*. 2003 Feb;23(1):29–34. doi:10.1046/j.1468-2982.2003.00441.x PubMed PMID: 12534577.
33. Cassidy EM, Tomkins E, Sharifi N, Dinan T, Hardiman O, O’Keane V. Differing central amine receptor sensitivity in different migraine subtypes? A neuroendocrine study using buspirone. *Pain*. 2003 Feb;101(3):283–90. doi:10.1016/S0304-3959(02)00335-4 PubMed PMID: 12583871.
34. Leone M, Attanasio A, Croci D, Ferraris A, D’Amico D, Grazzi L, et al. 5-HT_{1A} receptor hypersensitivity in migraine is suggested by the m-chlorophenylpiperazine test. *Neuroreport*. 1998 Aug 3;9(11):2605–8. doi:10.1097/00001756-199808030-00033 PubMed PMID: 9721941.
35. Gordon ML, Lipton RB, Brown SL, Nakraseive C, Russell M, Pollack SZ, et al. Headache and Cortisol Responses to M-Chlorophenylpiperazine are Highly Correlated. *Cephalalgia*. 1993 Dec 1;13(6):400–5. doi:10.1046/j.1468-2982.1993.1306400.x
36. Glover V, Ahmed F, Hussein N, Jarman J, Peatfield R. Central 5-hydroxytryptamine supersensitivity in migraine. In: *Migraine: Pharmacology and genetics* [Internet]. Chapman & Hall; 1996. p. 117–23. Available from: <https://archive.org/details/migraine-pharmacology-and-genetics/page/116/>
37. Murialdo G, Masturzo P, Filippi U, De Palma D, Balbi D, Polleri A, et al. Thyroid-Stimulating Hormone and Prolactin Responses to Thyrotropin-Releasing Hormone in Common Migraine. *Cephalalgia*. 1987 Dec 1;7(4):267–72. doi:10.1046/j.1468-2982.1987.0704267.x
38. Papakostas Y, Daras M, Markianos M, Stefanis C. Increased prolactin response to thyrotropin releasing hormone during migraine attacks. *Journal of Neurology, Neurosurgery & Psychiatry*. 1987 Jul 1;50(7):927–8. doi:10.1136/jnnp.50.7.927 PubMed PMID: 3114432.
39. Awaki E, Takeshima T, Takahashi K. A Neuroendocrinological Study in Female Migraineurs: Prolactin and Thyroid Stimulating Hormone Responses. *Cephalalgia*. 1989 Sep 1;9(3):187–93. doi:10.1046/j.1468-2982.1989.0903187.x
40. Capellan JIL, Dacosta CV, Garcia JMG, Servan PR, Cermeño JCA. Tuberoinfundibular Dopaminergic Tonus in Common Migraine. *Headache: The Journal of Head and Face Pain*. 1990;30(5):282–4. doi:10.1111/j.1526-4610.1990.hed3005282.x
41. Klimek A. Growth Hormone and Prolactin Release in Migraine Patients. In: Pfaffenrath V, Lundberg PO, Sjaastad O, editors. *Updating in Headache*. Berlin, Heidelberg: Springer; 1985. p. 241–8. doi:10.1007/978-3-642-88581-5_40

42. Klimek A. Metoclopramide Tests in Headache Patients. *Cephalalgia*. 1985 Jul 1;5(3_suppl):354–5. doi:10.1177/03331024850050S3138
43. Bussone G, Boiardi A, Mantia LL, Frediani F, Vescovi A, Parati EA. Long-acting ergot: clinical and neuroendocrinological aspects in migraine patients. *Cephalalgia*. 1983 Aug 1;3(1_suppl):163–7. doi:10.1177/03331024830030S127
44. Bussone G, Frediani F, Lamperti E, LaMantia L, Vescovi A, Peccarisi C, et al. Piribedil test in migraine: neuroendocrinological aspects. *Headache: The Journal of Head and Face Pain*. 1986;26(9):482–5. doi:10.1111/j.1526-4610.1986.hed2609482.x
45. Rainero I, Valfrè W, Savi L, Gentile S, Pinessi L, Gianotti L, et al. Neuroendocrine effects of subcutaneous sumatriptan in patients with migraine. *J Endocrinol Invest*. 2001 May;24(5):310–4. doi:10.1007/BF03343866 PubMed PMID: 11407649.
46. O’Mahony S, Chua ASB, Quigley EMM, Clarke G, Shanahan F, Keeling PWN, et al. Evidence of an enhanced central 5HT response in irritable bowel syndrome and in the rat maternal separation model. *Neurogastroenterol Motil*. 2008 Jun;20(6):680–8. doi:10.1111/j.1365-2982.2007.01065.x PubMed PMID: 18194152.
47. Chua A, Keating J, Hamilton D, Keeling PW, Dinan TG. Central serotonin receptors and delayed gastric emptying in non-ulcer dyspepsia. *BMJ*. 1992 Aug 1;305(6848):280–2. doi:10.1136/bmj.305.6848.280 PubMed PMID: 1392859; PubMed Central PMCID: PMC1882744.
48. Dinan TG, Chua AS, Keeling PW. Serotonin and physical illness: focus on non-ulcer dyspepsia. *J Psychopharmacol*. 1993 Jan;7(1):126–30. doi:10.1177/026988119300700104 PubMed PMID: 22289664.
49. Dinan TG, Mahmud N, Rathore O, Thakore J, Scott LV, Carr E, et al. A double-blind placebo-controlled study of buspirone-stimulated prolactin release in non-ulcer dyspepsia—are central serotonergic responses enhanced? *Alimentary Pharmacology & Therapeutics*. 2001;15(10):1613–8. doi:10.1046/j.1365-2036.2001.01090.x
50. Dinan TG, Yatham LN, Barry S, Chua A, Keeling PW. Serotonin supersensitivity: the pathophysiologic basis of non-ulcer dyspepsia? A preliminary report of buspirone/prolactin responses. *Scand J Gastroenterol*. 1990 May;25(5):541–4. doi:10.3109/00365529009095527 PubMed PMID: 2359983.
51. Spitz IM, Zylber E, Cohen H, Almaliach U, Leroith D. Impaired prolactin response to thyrotropin-releasing hormone in isolated gonadotropin deficiency and exaggerated response in primary testicular failure. *J Clin Endocrinol Metab*. 1979 Jun;48(6):941–5. doi:10.1210/jcem-48-6-941 PubMed PMID: 109459.
52. Spitz IM, Leroith D, Livshin Y, Zylber-Haran E, Trestian S, Laufer N, et al. Exaggerated Prolactin Response to Thyrotropin-Releasing Hormone and Metoclopramide in Primary Testicular Failure*. *Fertility and Sterility*. 1980 Dec 1;34(6):573–80. doi:10.1016/S0015-0282(16)45198-8

53. Hochner-Celnikier D, Zylber-Haran E, Shilo S, Palti Z, Spitz IM. Increased prolactin response to thyrotropin-releasing hormone in primary ovarian failure. *Obstet Gynecol.* 1982 Mar;59(3):280–4. PubMed PMID: 6804899.
54. Szilágyi A, Csermely T, Csaba IF. Increased prolactin response to TRH in polycystic ovary syndrome with low basal prolactin values. *Gynecological Endocrinology.* 1988 Jan;2(4):313–7. doi:10.3109/09513598809107654
55. Power AC, Cowen PJ. Neuroendocrine challenge tests: Assessment of 5-HT function in anxiety and depression. *Molecular Aspects of Medicine.* 1992 Jan 1;13(3):205–20. doi:10.1016/0098-2997(92)90010-W
56. Navinés R, Gómez-Gil E, Martín-Santos R, de Osaba MJM, Escolar G, Gastó C. Hormonal response to buspirone is not impaired in major depression. *Human Psychopharmacology: Clinical and Experimental.* 2007;22(6):389–95. doi:10.1002/hup.862
57. Lim EJ, Ahn YC, Jang ES, Lee SW, Lee SH, Son CG. Systematic review and meta-analysis of the prevalence of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME). *J Transl Med.* 2020 Feb 24;18(1):100. doi:10.1186/s12967-020-02269-0
58. Buckman MT, Peak GT. Estrogen Potentiation of Phenothiazine-Induced Prolactin Secretion in Man. *J Clin Endocrinol Metab.* 1973 Dec 1;37(6):977–80. doi:10.1210/jcem-37-6-977
59. Noel GL, Dimond RC, Wartofsky L, Earll JM, Frantz AG. Studies of Prolactin and TSH Secretion by Continuous Infusion of Small Amounts of Thyrotropin-Releasing Hormone (TRH). *J Clin Endocrinol Metab.* 1974 Jul 1;39(1):6–17. doi:10.1210/jcem-39-1-6
60. Schwinn G, von zur Mühlen A, Köbberling J, Halves E, Wenzel KW, Meinhold H. Plasma prolactin levels after TRH and chlorpromazine in normal subjects and patients with impaired pituitary function. *Acta Endocrinol (Copenh).* 1975 Aug;79(4):663–76. doi:10.1530/acta.0.0790663 PubMed PMID: 808064.
61. O’Keane V, O’Hanlon M, Webb M, Dinan T. d-fenfluramine/prolactin response throughout the menstrual cycle: evidence for an oestrogen-induced alteration. *Clin Endocrinol (Oxf).* 1991 Apr;34(4):289–92. doi:10.1111/j.1365-2265.1991.tb03768.x PubMed PMID: 1879060.
62. Reymond M, Lemarchand-Béraud Th. Effects of oestrogens on prolactin and thyrotrophin responses to TRH in women during the menstrual cycle and under oral contraceptive treatment. *Clinical Endocrinology.* 1976 Sep;5(5):429–37. doi:10.1111/j.1365-2265.1976.tb01973.x
63. Tyson JE, Friesen HG. Factors influencing the secretion of human prolactin and growth hormone in menstrual and gestational women. *American Journal of Obstetrics and Gynecology.* 1973 Jan 1;116(3):377–87. doi:10.1016/S0002-9378(15)31297-7
64. De Léan, A, Ferland L, Drouin J, Kelly PA, Labrie F. Modulation of Pituitary Thyrotropin Releasing Hormone Receptor Levels by Estrogens and Thyroid Hormones. *Endocrinology.* 1977 Jun 1;100(6):1496–504. doi:10.1210/endo-100-6-1496

65. Messini CI, Malandri M, Anifandis G, Dafopoulos K, Georgoulas P, Sveronis G, et al. Submaximal doses of ghrelin do not inhibit gonadotrophin levels but stimulate prolactin secretion in postmenopausal women. *Clinical Endocrinology*. 2017;87(1):44–50. doi:10.1111/cen.13349
66. Labrie F, Ferland L, Denizeau F, Beaulieu M. Sex steroids interact with dopamine at the hypothalamic and pituitary levels to modulate prolactin secretion. *Journal of Steroid Biochemistry*. 1980 Jan 1;12:323–30. doi:10.1016/0022-4731(80)90287-3
67. Stewart WF, Lipton RB, Celentano DD, Reed ML. Prevalence of Migraine Headache in the United States: Relation to Age, Income, Race, and Other Sociodemographic Factors. *JAMA*. 1992 Jan 1;267(1):64–9. doi:10.1001/jama.1992.03480010072027
68. Victor T, Hu X, Campbell J, Buse D, Lipton R. Migraine prevalence by age and sex in the United States: A life-span study. *Cephalalgia*. 2010 Sep;30(9):1065–72. doi:10.1177/0333102409355601
69. Cevik R, Gur A, Acar S, Nas K, Sarac AJ. Hypothalamic-pituitary-gonadal axis hormones and cortisol in both menstrual phases of women with chronic fatigue syndrome and effect of depressive mood on these hormones. *BMC Musculoskelet Disord*. 2004 Dec 8;5(1):47. doi:10.1186/1471-2474-5-47
70. Studd J, Panay N. Chronic fatigue syndrome. *The Lancet*. 1996 Nov 16;348(9038):1384. doi:10.1016/S0140-6736(05)65448-7 PubMed PMID: 8918295.
71. Thomas N, Ubhayasekera SJKA, Armstrong CW, Huang K, Bergquist J. Steroid dynamics in myalgic encephalomyelitis / chronic fatigue syndrome: a case-control study using ultra performance supercritical fluid chromatography tandem mass spectrometry. *J Transl Med*. 2025 Jul 25;23(1):829. doi:10.1186/s12967-025-06841-4
72. Wyller VB, Evang JA, Godang K, Solhjell KK, Bollerslev J. Hormonal alterations in adolescent chronic fatigue syndrome. *Acta Paediatrica*. 2010;99(5):770–3. doi:10.1111/j.1651-2227.2010.01701.x
73. Pasqualini C, Bojda F, Kerdelhué B. Direct Effect of Estradiol on the Number of Dopamine Receptors in the Anterior Pituitary of Ovariectomized Rats. *Endocrinology*. 1986 Dec 1;119(6):2484–9. doi:10.1210/endo-119-6-2484
74. Cleare AJ, Messa C, Rabiner EA, Grasby PM. Brain 5-HT_{1A} receptor binding in chronic fatigue syndrome measured using positron emission tomography and [¹¹C]WAY-100635. *Biological Psychiatry*. 2005 Feb;57(3):239–46. doi:10.1016/j.biopsych.2004.10.031
75. Lamberts SWJ, Macleod RM. The Interaction of the Serotonergic and Dopaminergic Systems on Prolactin Secretion in the Rat: The Mechanism of Action of the “Specific” Serotonin Receptor Antagonist, Methysergide. *Endocrinology*. 1978 Jul 1;103(1):287–95. doi:10.1210/endo-103-1-287
76. Lynch CO, Johnson MD, Crowley WR. Effects of the serotonin agonist, quipazine, on luteinizing hormone and prolactin release: Evidence for serotonin-catecholamine interactions. *Life Sciences*. 1984 Oct 1;35(14):1481–7. doi:10.1016/0024-3205(84)90165-6

77. Baraniuk JN, Eaton-Fitch N, Marshall-Gradisnik S. Meta-analysis of natural killer cell cytotoxicity in myalgic encephalomyelitis/chronic fatigue syndrome. *Front Immunol.* 2024;15:1440643. doi:10.3389/fimmu.2024.1440643 PubMed PMID: 39483457; PubMed Central PMCID: PMC11524851.
78. Reddymasu SC, Soykan I, McCallum RW. Domperidone: review of pharmacology and clinical applications in gastroenterology. *Am J Gastroenterol.* 2007 Sep;102(9):2036–45. doi:10.1111/j.1572-0241.2007.01255.x PubMed PMID: 17488253.
79. Shishioh-Ikejima N, Ogawa T, Yamaguti K, Watanabe Y, Kuratsune H, Kiyama H. The increase of alpha-melanocyte-stimulating hormone in the plasma of chronic fatigue syndrome patients. *BMC Neurol.* 2010 Aug 23;10:73. doi:10.1186/1471-2377-10-73 PubMed PMID: 20731841; PubMed Central PMCID: PMC2933583.
80. Hill JB, Nagy GM, Frawley LS. Suckling Unmasks the Stimulatory Effect of Dopamine on Prolactin Release: Possible Role for α -Melanocyte- Stimulating Hormone as a Mammatrope Responsiveness Factor*. *Endocrinology.* 1991 Aug;129(2):843–7. doi:10.1210/endo-129-2-843
81. Khorram O, Bedran De Castro JC, McCann SM. Physiological role of alpha-melanocyte-stimulating hormone in modulating the secretion of prolactin and luteinizing hormone in the female rat. *Proc Natl Acad Sci USA.* 1984 Dec;81(24):8004–8. doi:10.1073/pnas.81.24.8004
82. Newman CB, Wardlaw SL, Frantz AG. Suppression of basal and stress-induced prolactin release and stimulation of luteinizing hormone secretion by α -melanocyte-stimulating hormone. *Life Sciences.* 1985 Apr;36(17):1661–8. doi:10.1016/0024-3205(85)90369-8
83. Nuñez L, Frawley LS. α -MSH potentiates the responsiveness of mammatropes by increasing Ca^{2+} entry. *American Journal of Physiology-Endocrinology and Metabolism.* 1998 Jun 1;274(6):E971–7. doi:10.1152/ajpendo.1998.274.6.E971