

# Alpha-taxonomy in world mammals

R. van Mazijk VMZRUA001@myuct.ac.za

2017-08-24

BSc Hons Biological Sciences, Species & speciation module deliverable

For A/Prof A.M. Muasya



Herein I discuss the progress of mammalian alpha-taxonomy from 1975 to 2005. First, I focus on the methods employed to diagnose and delimit these mammal species and how these have changes over the the past 30 to 40 years. Second, I discuss the new species and groups discovered in this period. Third, I will attempt to draw conclusions, if possible, about the speciation mechanisms at work in mammal populations based on the evidence use to delimit those mammal species. I will frame these discussions in the context of the species concepts (or lack thereof) typically used in mammalogy and mammalian taxonomy.

*Mammal Species of the World, 3rd Edition* (MSW3) (Wilson & Reeder 2005) is a reputable database of mammalian taxa (species, sub-specific, and super-specific) and information concerning their treatments (e.g. scientific name, authors' name and year described, original publication citation, distribution, etc. (Wilson & Reeder 2005)) from 1702 to 2005 AD. This data product is a boon to the discussions here. Using the .csv file from the MSW3 webpage (<https://www.departments.bucknell.edu/biology/resources/msw3/>), I explored this dataset in R (R Core Team 2017), primarily using the `tidyverse` suite of packages for data exploration and visualisation (Wickham 2017). Not only does MSW3 enable exploration of mammal taxonomic discovery through time, it also facilitated easier literature searches, providing common journals in which mammal descriptions are published.

There is a subset of mammal order that are most taxonomically “active” (Figure 1), such that there are only eight orders in which  $\geq 10$  new species were described in the period 1975—2005. This is worth bearing in mind when discussing mammalian taxonomy; there is a taxonomic bias to, either, the rate of discovery in certain taxa, or the species richness of certain taxa. Either way, species concepts have bearing on these numbers, discussed below.

## Mammalogical species concepts

The species concept or criteria a taxonomist follows can affect the number of species delimited. Supposing there is reality to species, and that two lineages being different species is grounded in reality, species concepts enable us to estimate when this occurs. Following the reasoning of De Queiroz (2011), different




Figure 1: Bar-plots of the number of new species (*sensu stricto*, excluding sub-specific additions) described each year in the eight most taxonomically active orders of mammal over the period 1975—2005. Data are from MSW3 (see text) (Wilson & Reeder 2005). Taxonomically “active” here implies that  $\geq 10$  new species were described for an order over the 1975—2005 period. These eight orders are those that meet this criterion, such 533 species’ discoveries are illustrated here. For reference, Dasyuromorphs are commonly referred to as Tasmanian devils and numbats, Didelphimorphs as opossums, Diprotodonts as koalas and wombats, and Soricomorpha as moles and shrews

species concepts each deal with different lines of empirical evidence. These each describe different aspects of the speciation process, such that in each the criteria for delimiting two different species is based on some property of each lineage that would differ between them if they were separate species (De Queiroz 2011). Consequently, different species concepts each encapsulate *some* aspect of dissimilarity between two lineages as a consequence of speciation. De Queiroz (2011) argues that should any one criterion be met, it is parsimonious to assume speciation has begun, and that the bifurcation of the lineages be appropriately regarded with a species delineation. This aside, consistency in species concepts in taxonomic research is still desirable (if perhaps unattainable). The case with mammal taxonomists is summarised well:

*Mammalogists, for the most part, have avoided debate on species concepts, but most investigators follow a particular species definition.*

Baker & Bradley (2006) make this somewhat alarming statement about the *modus operandi* of mammal taxonomists. It appears that mammal taxonomy, like many fields of biology, has agreed-upon nomenclatural standards, but has little consensus on which species concepts to use when delineating species. This precludes ready comparison of, say, mice species vs bovid species, as there is no guarantee that “species” represent similarly biologically important units.

For the most part, mammalogists historically followed (if only loosely) the Biological Species Concept (BSC). This is intuitive as mammals are sexually reproducing eukaryotes (???; Baker & Bradley 2006). The Phylogenetic Species Concept (PSC) contrasts with the BSC in focussing on an explicitly evolutionary, lineage based criterion. The PSC thus now dominates in mammalogy, as the most common species concept used or referred to. The PSC solves the problem of the BSC wherein entities widely regarded as ideally being two species fail to be differentiated reproductively. Baker & Bradley (2006) themselves put forward the Genetic Species Concept (GSC), wherein a species represents:

*[...] a group of genetically compatible interbreeding natural populations that is genetically isolated from other such groups.*

This does not necessarily imply reproductive isolation either, as can be seen in the case of *Thomomys* pocket gophers (Baker & Bradley 2006). The transition from majority usage of the BSC to PSC in mammal taxonomy is not so simple, however. There is little explicit consideration of species concepts by the majority of mammal taxonomists. And, there is debate. Groves (2012) in particular advocate more strongly for the PSC, even in place of the GSC, in that species defined by the PSC are:

*[...] 100% diagnosable: they have fixed heritable differences between them [...] [and the PSC] offer[s] a repeatable, falsifiable definition of species [...]*

This is evident in the widespread application of the PSC in mammal taxonomy. But nonetheless, the lack of explicit concern for species criteria is troubling. Do mammal taxonomists employ similar methodologies

when delimiting species? I discuss this below, despite the nuanced inconsistencies that differing species concepts among mammalogists may lead to.

## Lines of evidence

Morphology was the dominant method in mammal species delimitation prior to 1985 (Baker & Bradley 2006). Indeed, morphological differences between taxa are still influential in their taxonomy. Furthermore, allopatric populations of mammal that are morphologically distinct (whatever that may entail) are regularly treated as separate species. This concurs with the usage of the PSC above, where non-reproductively isolated taxa (e.g. grizzly and polar bears; lions and tigers) are treated as separate species. Phenetics still common practice in mammal taxonomy, wherein morphological divergence between populations is numerically described. Species are then diagnosed as clusters in this quantitative trait-space. Phenetics and morphology are often also the starting points of alpha-taxonomic research (e.g. Lew et al. 2006), as it is more amenable to efficient, cheap study than molecular systematics.

Early molecular work used in mammal taxonomy employed karyotype studies (Baker & Bradley 2006)—to delineate species on the basis of chromosomal traits (e.g. diploid number, chromosome structure, arrangement). Allozymes and DNA sequencing methods arrived later. These molecular methods were costly in their infancy (Baker & Bradley 2006), but soon became much more practical. The benefits of these methods is their greater objectivity, in that qualitatively scored morphological traits are more subject to researchers' idiosyncrasies. The other benefit of characters divorced from observable morphology is that they can allow cryptic species to be uncovered (Zimmermann & Radespiel 2014). This does, however, contribute to taxonomic inflation. As Rylands et al. (2012) point out (in reference to primates, but this easily pertains to many biota), taxonomic inflation may indeed represent a greater understanding of existing diversity. Furthermore, molecular systematic methods may reveal morphologically distinct populations to in fact be polymorphs of a single species (Zimmermann & Radespiel 2014). In fact,

## Mammal discoveries, 1975—2005

Discoveries of new mammal species over this 30 year period are dominated by rodents and bats (Figure 1). It is thus interesting that, in the Cape, only one new mammal species was described in the the period 1975 to 2005 (Meester & Dippenaar 1978): *Myosorex longicaudatus*, from the order Soricomorpha, discovered around Knysna. This shrew species was first delimited on the basis of its morphology and geographic range, distinct from those of its relatives (Meester & Dippenaar 1978).

South African mammal species over this period include *Myosorex longicaudatus*, the rodent *Mastomys awashensis*, and the golden mole *Amblysomus robustus* (Afrosoricidimorpha).

- Africa
  - Primates, Rodents, Soricom.
- Global?
  -
- e.g. cytochrome-b (mtDNA)
  - Carvalho et al. 2009. Phylogeny of Thylamys (Didelphimorphia, Didelphidae) species, with special reference to Thylamys karimii
- cytochrome-b in rodents
  - Re: GSC sensu Baker & Bradley 2006 (J. of Mammal. 87(4):643–662), Bradley & Baker 2001 (J. of Mammal. 82(4):960–973)
- multi-locus work in Afro-tropical shrews
  - cytochrome-b, among others,
  - Jacquet et al. 2012. Zool. J. Linn. Soc. 166(3):672–687

## Modes of mammalian speciation

## Concluding remarks

- Molecular species delimitation is relevant to PSC/ESC/BSC
- Ecologically relevant speciation processes (e.g. bats) are case-by-case
- Little generality to be drawn about speciation from the alpha-taxonomic methods
- Genetic sequencing & PSC (or at least ESC)
- Good taxonomic inflation!
- Genomics is up-and-coming! e.g. Murphy et al. 2004

---

## References

- Baker, R.J. & Bradley, R.D. 2006. Speciation in Mammals and the Genetic Species Concept. *J. Mammal.* 87(4):643–662. DOI: 10.1644/06-MAMM-F-038R2.1.
- De Queiroz, K. 2011. Branches in the lines of descent: Charles Darwin and the evolution of the species concept. *Biological Journal of the Linnean Society.* 103:19–35. DOI: 10.1111/j.1095-8312.2011.01634.x.
- Groves, C. 2012. Species Concept in Primates. *American Journal of Primatology.* 74(8):687–691. DOI: 10.1002/ajp.22035.
- Lew, D., Pérez-Hernandez, R. & Ventura, J. 2006. Two New Species of Philander (Didelphimorphia, Didelphidae) from Northern South America. *Journal of Mammalogy.* 87(2):224–237. DOI: 10.1644/05-MAMM-A-065R2.1.
- Meester, J. & Dippenaar, N. 1978. A New Species Of Myosorex From Knysna, South Africa (Mammalia: Soricidae). *Annals of the Transvaal Museum.* 31(4):30–42. Available: [http://journals.co.za.ezproxy.uct.ac.za/docserver/fulltext/nfi\\_annalstm/](http://journals.co.za.ezproxy.uct.ac.za/docserver/fulltext/nfi_annalstm/)

31/4/201.pdf.

- R Core Team. 2017. *R: A Language and Environment for Statistical Computing*. 2017. Vienna, Austria: R Foundation for Statistical Computing. Available: <https://www.r-project.org/>.
- Rylands, A.B., Mittermeier, R.A. & Silva, J.S. 2012. Neotropical primates: Taxonomy and recently described species and subspecies. *International Zoo Yearbook*. 46(1):11–24. DOI: 10.1111/j.1748-1090.2011.00152.x.
- Wickham, H. 2017. *tidyverse: Easily Install and Load 'Tidyverse' Packages*. 2017. Available: <https://cran.r-project.org/package=tidyverse>.
- Wilson, D. & Reeder, D.-A. Eds. 2005. *Mammal Species of the World. A Taxonomic and Geographic Reference*. 3rd edition. 2005. Baltimore: John Hopkins University Press. Available: <https://www.departments.bucknell.edu/biology/resources/msw3/>.
- Zimmermann, E. & Radespiel, U. 2014. Species concepts, diversity, and evolution in primates: Lessons to be learned from mouse lemurs. *Evolutionary Anthropology*. 23(1):11–14. DOI: 10.1002/evan.21388.