Dynamic CO-CIN report to SAGE and NERVTAG (Recent cases)

Dynamic content updated: 2020-11-25 03:38:03.


Executive summary


Includes patients admitted after 01 August 2020

There are 20006 patients included in CO-CIN. Of these, 2666 patient(s) have died and 2608 required ICU. 10338 have been discharged home.


Admission

Figure 1






Number with symptom onset occuring after admission to hospital Figure 1C





Proportion with symptom onset occuring after hospital admission

Figure 1D - 7-day rolling percentage. n = 14428



Patients with outcome stratified by age, and sex Figure 1E






Location by CCG / Healthboard

Figure 2

Click and drag on map to zoom into area. Reset via toolbar at top of map.





Symptoms and comorbidity

Figure 3A




Figure 3B - Symptoms by Age

Symptoms on presentation to hospital Stratifed by age






Figure 3C

Correlation of symtpoms in all pages. Note clusters, top right to bottom left, flu-like, coryzal, abdominal, respiratory, neurocutaneous.




Symptoms (diagnostic criteria)

Figure 4A n = 17580




Symptoms (most common)

Figure 4B n = 17580




Comorbidity (most common)

Figure 4C n = 20006



Medication prior to illness

Figure 5




Preadmission treatment

Figure 6

Pre-admission treatment

Anti-infectives for illness episode (left) immunosuppressants including oral (not



Patient flow

Figure 7A - All patients

N = 17962





Figure 7B - Patients admitted >=14 days and <=28 days ago

N = 3369



Oxygen requirement

Figure 8A - All patients

N = 17123






Figure 8B - Patients admitted >=14 days and <=28 days ago

In-hospital medical treatment

Figure 9



Treatment




Status in patients admitted >=14 days from today


Figure 11

All: status in patients admitted >=14 days ago


Figure 11A



ICU/HDU admissions: status in patients admitted >=14 days ago




Invasive ventilation: status in patients admitted >=14 days ago Figure 11c



Length of stay stratified by age

Figure 12

Length of stay stratified by age and mortality Proportion who reach outcome by day 14 shown



Predictors of death: logistic regression multivariable model

Logistic regression model only includes patients admitted >14 days ago from today.

| Dependent: death | | No | Yes | OR (univariable) | OR (multivariable) |
|--------------------------|-------|-------------|------------|---------------------------|---------------------------|
| Age on admission (years) | <50 | 2838 (97.9) | 61 (2.1) | • | • |
| | 50-69 | 3580 (89.5) | 421 (10.5) | 5.47 (4.20-7.25, p<0.001) | 4.61 (3.42-6.36, p<0.001) |


| Dependent: death | | No | Yes | OR (univariable) | OR (multivariable) |
|--|------------------|-----------------|----------------|---------------------------------|---------------------------------|
| | 70-79 | 2238 (74.6) | 764 (25.4) | 15.88 (12.27-20.94, p<0.001) | 11.58 (8.61-15.93, p<0.001) |
| | 80+ | 2414 (63.7) | 1378 (36.3) | 26.56 (20.62-34.87, p<0.001) | 18.76 (13.96-25.79, p<0.001) |
| Sex at Birth | Male | 5837 (78.5) | 1596 (21.5) | • | • |
| | Female | 5236 (83.6) | 1028 (16.4) | 0.72 (0.66-0.78, p<0.001) | 0.67 (0.61-0.75, p<0.001) |
| | Not specified | 0 (NaN) | 0 (NaN) | • | • |
| Chronic cardiac disease | NO | 7906 (86.0) | 1292 (14.0) | • | • |
| | YES | 2562 (68.7) | 1167 (31.3) | 2.79 (2.55-3.05, p<0.001) | 1.31 (1.17-1.46, p<0.001) |
| Chronic pulmonary disease (not asthma) | NO | 9051 (83.4) | 1806 (16.6) | • | • |
| | YES | 1418 (68.3) | 657 (31.7) | 2.32 (2.09-2.58, p<0.001) | 1.51 (1.33-1.72, p<0.001) |
| Chronic neurological disorder | NO | 9522 (81.9) | 2104 (18.1) | • | • |
| | YES | 902 (73.5) | 326 (26.5) | 1.64 (1.43-1.87, p<0.001) | 1.10 (0.93-1.30, p=0.259) |
| Chronic hematologic disease | NO | 10047 (81.5) | 2279 (18.5) | • | • |
| | YES | 369 (70.7) | 153 (29.3) | 1.83 (1.50-2.21, p<0.001) | 1.29 (1.01-1.63, p=0.037) |
| Chronic kidney disease | NO | 9235 (83.7) | 1799 (16.3) | • | • |
| | YES | 1222 (65.4) | 647 (34.6) | 2.72 (2.44-3.03, p<0.001) | 1.45 (1.26-1.65, p<0.001) |
| Dementia | NO | 9651 (83.1) | 1968 (16.9) | • | • |
| | YES | 791 (63.7) | 450 (36.3) | 2.79 (2.46-3.16, p<0.001) | 1.34 (1.15-1.57, p<0.001) |
| Obesity (as defined by clinical staff) | NO | 7885 (80.9) | 1867 (19.1) | • | • |
| | YES | 1399 (85.0) | 247 (15.0) | 0.75 (0.64-0.86, p<0.001) | 1.05 (0.88-1.24, p=0.588) |
| Malignant neoplasm | NO | 9652 (82.6) | 2034 (17.4) | • | • |
| | YES | 773 (65.6) | 406 (34.4) | 2.49 (2.19-2.83, p<0.001) | 1.61 (1.37-1.89, p<0.001) |

Number in dataframe = 18460, Number in model = 10942, Missing = 7518, AIC = 8714.9, C-statistic = 0.775, H&L = Chi-sq(8) 18.38 (p=0.019)

Figure 13 - Adjusted odds ratio plot

| Death | | | |
|---------------------------------------|-----------------|------------------------------|--------------------------------|
| Age on admission (years) | <50 | - | • |
| | 50-69 | 4.61 (3.42-6.36, p<0.001) | ⊢■ |
| | 70-79 | 11.58 (8.61-15.93, p<0.001) | ⊢ ■ |
| | 80+ | 18.76 (13.96-25.79, p<0.001) | ⊢■ |
| Sex at Birth | Male | - | • |
| | Female | 0.67 (0.61-0.75, p<0.001) | |
| Not s | pecified | - | • |
| Chronic cardiac disease | YES | 1.31 (1.17-1.46, p<0.001) | ⊢⊞ + |
| Chronic pulmonary disease (not as | thm ¥ #S | 1.51 (1.33-1.72, p<0.001) | + ■ + |
| Chronic neurological disorder | YES | 1.10 (0.93-1.30, p=0.259) | + ■ - |
| Chronic hematologic disease | YES | 1.29 (1.01-1.63, p=0.037) | ▶ |
| Chronic kidney disease | YES | 1.45 (1.26-1.65, p<0.001) | +■- |
| Dementia | YES | 1.34 (1.15-1.57, p<0.001) | ⊢= ⊣ |
| Obesity (as defined by clinical staff |) YES | 1.05 (0.88-1.24, p=0.588) | - |
| Malignant neoplasm | YES | 1.61 (1.37-1.89, p<0.001) | ⊢■ → |
| | | | Odds ratio (95% CI, log scale) |


Figure 14 - ROC



Survival models

Kaplan-Meier plots for survival from symptom onset stratified by age (left) and sex (right)

Figure 15
P-value is log-rank test.



Cox proportional hazards model

The methodology for this is now up and running, but models are still being explored. **What is presented here is not a final model, but to demonstrate methodology**. The results are correct, but important variables have not yet been included.

Time from symptom onset.

| Dependent: Surv(time, status) | | all | HR (univariable) | HR (multivariable) |
|--|-------------|--------------|------------------------------|------------------------------|
| Age on admission (years) | <50 | 3811 (19.3) | • | • |
| | 50-69 | 5749 (29.0) | 5.33 (4.06-7.00, p<0.001) | 4.20 (3.04-5.79, p<0.001) |
| | 70-79 | 4454 (22.5) | 14.14 (10.86-18.43, p<0.001) | 10.85 (7.93-14.85, p<0.001) |
| | 80+ | 5783 (29.2) | 22.06 (17.00-28.62, p<0.001) | 16.39 (11.97-22.43, p<0.001) |
| Sex at Birth | Male | 10840 (54.8) | • | • |
| | Female | 8950 (45.2) | 0.75 (0.70-0.81, p<0.001) | 0.76 (0.69-0.83, p<0.001) |
| qSOFA score on admission | 0 | 7871 (51.2) | • | • |
| | 1 | 6476 (42.1) | 1.80 (1.64-1.97, p<0.001) | 1.97 (1.78-2.18, p<0.001) |
| | 2 | 985 (6.4) | 4.38 (3.86-4.96, p<0.001) | 3.79 (3.28-4.38, p<0.001) |
| | 3 | 51 (0.3) | 7.98 (5.32-11.98, p<0.001) | 9.42 (6.08-14.60, p<0.001) |
| Symptomatic at presentation | No symptoms | 2381 (13.4) | • | • |
| | Symptoms | 15435 (86.6) | 1.51 (1.33-1.72, p<0.001) | • |
| Chronic cardiac disease | NO | 12110 (70.9) | • | • |
| | YES | 4978 (29.1) | 2.49 (2.30-2.70, p<0.001) | 1.29 (1.17-1.42, p<0.001) |
| Chronic kidney disease | NO | 14565 (85.4) | • | • |
| | YES | 2493 (14.6) | 2.40 (2.19-2.62, p<0.001) | 1.34 (1.19-1.49, p<0.001) |
| Moderate or severe liver disease | NO | 16626 (98.2) | • | • |
| | YES | 305 (1.8) | 1.77 (1.40-2.24, p<0.001) | 1.52 (1.11-2.08, p=0.009) |
| Chronic neurological disorder | NO | 15313 (90.2) | • | • |
| | YES | 1662 (9.8) | 1.51 (1.34-1.70, p<0.001) | • |
| Malignant neoplasm | NO | 15386 (90.5) | • | • |
| | YES | 1615 (9.5) | 2.15 (1.94-2.39, p<0.001) | 1.60 (1.41-1.82, p<0.001) |
| Chronic hematologic disease | NO | 16287 (96.0) | • | • |
| | YES | 671 (4.0) | 1.68 (1.43-1.98, p<0.001) | • |
| Obesity (as defined by clinical staff) | NO | 12984 (85.9) | • | • |
| | YES | 2126 (14.1) | 0.77 (0.67-0.88, p<0.001) | 1.01 (0.87-1.17, p=0.901) |

| Dependent: Surv(time, status) | | all | HR (univariable) | HR (multivariable) |
|--------------------------------|-----|--------------|---------------------------|---------------------------|
| Diabetes without complications | NO | 14060 (87.5) | • | • |
| | YES | 2003 (12.5) | 1.21 (1.08-1.36, p=0.001) | • |
| Rheumatologic disorder | NO | 14821 (87.4) | • | • |
| | YES | 2141 (12.6) | 1.31 (1.17-1.46, p<0.001) | • |
| Dementia | NO | 15283 (89.8) | • | • |
| | YES | 1728 (10.2) | 2.41 (2.18-2.67, p<0.001) | 1.13 (0.99-1.29, p=0.066) |
| Malnutrition | NO | 15470 (98.0) | • | • |
| | YES | 323 (2.0) | 1.78 (1.41-2.25, p<0.001) | • |
| smoking_mhyn_2levels | NO | 8853 (50.3) | • | • |
| | YES | 949 (5.4) | 0.99 (0.82-1.19, p=0.876) | • |
| | N/K | 7813 (44.4) | 1.31 (1.21-1.42, p<0.001) | • |

Number in dataframe = 19895, Number in model = 12565, Missing = 7330, Number of events = 1807, Concordance = 0.779 (SE = 0.005), R-squared = 0.136 (Max possible = 0.929), Likelihood ratio test = 1843.083 (df = 13, p = 0.000)

Figure 16a - Multivariable Cox proportional hazards model





Figure 16b - Multivariable Cox proportional hazards model (age, sex, comorbidities only)



ROC = 0.7806839

Figure 17 - Predictions calibration plot




Figure 18 - Prognostic model predictions

Again, for demonstration of methods.

Prognostic model for survival 7 days after onset qSOFA score on admission




Figure 19 - Death by severity (NEWS) on admission

Number of deaths by NEWS score at admission




Figure 20 - Death by severity (NEWS) on admission stratified by age





Figure 21 - Death by severity (qSOFA) on admission stratified by age

Number of deaths by qSOFA score at admission Stratified by age



Healthcare workers

| Employed as a healthcare worker? | | NO | YES | р |
|----------------------------------|--------------|--------------|------------|--------|
| Total N (%) | | 16139 (97.6) | 389 (2.4) | |
| NEWS score on admission | Median (IQR) | 3.0 (4.0) | 3.0 (3.5) | 0.170 |
| Death | No | 9273 (79.1) | 294 (97.0) | <0.001 |
| | Yes | 2444 (20.9) | 9 (3.0) | |