

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \hat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ where A_{BR} is 0×0 , B_B has 0 rows, C_B has 0 rows
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \hat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \hat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \hat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \hat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	Determine block size b $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{c c c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ \hline A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$ where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows
6	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{20}^T B_2 + \hat{C}_0 \\ A_{21}^T B_2 + \hat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \hat{C}_2 \end{array} \right) \right\}$
8	$C_0 := A_{00} B_0 + A_{10}^T B_1 + A_{20}^T B_2 + C_0$ $C_1 := A_{10} B_0 + A_{11} B_1 + A_{21}^T B_2 + C_1$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{10}^T B_1 + A_{20}^T B_2 + \hat{C}_0 \\ A_{10} B_0 + A_{11} B_1 + A_{21}^T B_2 + \hat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \hat{C}_2 \end{array} \right) \right\}$
5b	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{c c c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ \hline A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \hat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \hat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \hat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \hat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \hat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	{
4	
	where
2	{
3	while do
2,3	{
	\wedge
5a	Determine block size b
	where
6	{
8	$A_{00}B_0 + A_{20}^T B_2 + A_{21}^T B_1 +$ $C_2 := A_{20}B_0 + A_{21}B_1 + A_{22}B_2 + C_2$
7	{
5b	
2	{
	endwhile
2,3	{
	$\wedge \neg($
1b	{

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	{ $C = \hat{C}$ }
4	where
2	{
3	while do
2,3	{ \wedge }
5a	Determine block size b where
6	{
8	$A_{00}B_0 + A_{20}^T B_2 + A_{21}^T B_1 +$ $C_2 := A_{20}B_0 + A_{21}B_1 + A_{22}B_2 + C_2$
7	{
5b	
2	{
	endwhile
2,3	{ $\wedge \neg($) }
1b	{ $C = AB + \hat{C}$ }

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	
	where
2	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \widehat{C}_T}{A_{BL} B_T + A_{BR} B_B + \widehat{C}_B} \right) \right\}$
3	while do
2,3	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \widehat{C}_T}{A_{BL} B_T + A_{BR} B_B + \widehat{C}_B} \right) \wedge \right\}$
5a	Determine block size b
	where
6	$\left\{ \right.$
8	$A_{00} B_0 + \quad A_{20}^T B_2 +$ $A_{21}^T B_1 +$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \right.$
5b	
2	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \widehat{C}_T}{A_{BL} B_T + A_{BR} B_B + \widehat{C}_B} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \widehat{C}_T}{A_{BL} B_T + A_{BR} B_B + \widehat{C}_B} \right) \wedge \neg(\quad) \right\}$
1b	$\{C = AB + \widehat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \hat{C}$
4	
	where
2	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \hat{C}_T}{A_{BL} B_T + A_{BR} B_B + \hat{C}_B} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \hat{C}_T}{A_{BL} B_T + A_{BR} B_B + \hat{C}_B} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	Determine block size b
	where
6	$\left\{ \right.$
8	$A_{00} B_0 + \quad A_{20}^T B_2 +$ $A_{21}^T B_1 +$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \right.$
5b	
2	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \hat{C}_T}{A_{BL} B_T + A_{BR} B_B + \hat{C}_B} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\frac{C_T}{C_B} \right) = \left(\frac{A_{BL}^T B_B + \hat{C}_T}{A_{BL} B_T + A_{BR} B_B + \hat{C}_B} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \hat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ <p style="color: red;">where A_{BR} is 0×0, B_B has 0 rows, C_B has 0 rows</p>
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	<p>Determine block size b</p> <p style="color: blue;">where</p>
6	$\left\{ \right.$
8	$A_{00}B_0 + A_{20}^T B_2 + A_{21}^T B_1 + C_2 := A_{20}B_0 + A_{21}B_1 + A_{22}B_2 + C_2$
7	$\left\{ \right.$
5b	
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \widehat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ where A_{BR} is 0×0 , B_B has 0 rows, C_B has 0 rows
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	Determine block size b $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{c c c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$ where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows
6	$\left\{ \right.$
8	$A_{00} B_0 +$ $A_{20}^T B_2 +$ $A_{21}^T B_1 +$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \right.$
5b	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{c c c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \widehat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ where A_{BR} is 0×0 , B_B has 0 rows, C_B has 0 rows
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	Determine block size b $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$ where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows
6	$\left\{ \begin{pmatrix} C_0 \\ C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} A_{20}^T B_2 + \widehat{C}_0 \\ A_{21}^T B_2 + \widehat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \widehat{C}_2 \end{pmatrix} \right\}$
8	$A_{00} B_0 +$ $A_{20}^T B_2 +$ $A_{21}^T B_1 +$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \right\}$
5b	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \widehat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ where A_{BR} is 0×0 , B_B has 0 rows, C_B has 0 rows
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	Determine block size b $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$ where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows
6	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{20}^T B_2 + \widehat{C}_0 \\ A_{21}^T B_2 + \widehat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \widehat{C}_2 \end{array} \right) \right\}$
8	$A_{00} B_0 +$ $A_{20}^T B_2 +$ $A_{21}^T B_1 +$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{10}^T B_1 + A_{20}^T B_2 + \widehat{C}_0 \\ A_{10} B_0 + A_{11} B_1 + A_{21}^T B_2 + \widehat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \widehat{C}_2 \end{array} \right) \right\}$
5b	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \widehat{C}$

Step	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
1a	$\{C = \widehat{C}$
4	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ C_B \end{array} \right)$ <p>where A_{BR} is 0×0, B_B has 0 rows, C_B has 0 rows</p>
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
3	while $m(A_{BR}) < m(A)$ do
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge m(A_{BR}) < m(A) \right\}$
5a	<p>Determine block size b</p> $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$ <p>where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows</p>
6	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{20}^T B_2 + \widehat{C}_0 \\ A_{21}^T B_2 + \widehat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \widehat{C}_2 \end{array} \right) \right\}$
8	$C_0 := A_{00} B_0 + A_{10}^T B_1 + A_{20}^T B_2 + C_0$ $C_1 := A_{10} B_0 + A_{11} B_1 + A_{21}^T B_2 + C_1$ $C_2 := A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + C_2$
7	$\left\{ \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right) = \left(\begin{array}{c} A_{10}^T B_1 + A_{20}^T B_2 + \widehat{C}_0 \\ A_{10} B_0 + A_{11} B_1 + A_{21}^T B_2 + \widehat{C}_1 \\ A_{20} B_0 + A_{21} B_1 + A_{22} B_2 + \widehat{C}_2 \end{array} \right) \right\}$
5b	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ C_1 \\ C_2 \end{array} \right)$
2	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \right\}$
	endwhile
2,3	$\left\{ \left(\begin{array}{c} C_T \\ C_B \end{array} \right) = \left(\begin{array}{c} A_{BL}^T B_B + \widehat{C}_T \\ A_{BL} B_T + A_{BR} B_B + \widehat{C}_B \end{array} \right) \wedge \neg(m(A_{BR}) < m(A)) \right\}$
1b	$\{C = AB + \widehat{C}$

	Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part
	$A \rightarrow \left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right)$ <p>where A_{BR} is 0×0, B_B has 0 rows, C_B has 0 rows</p>
	while $m(A_{BR}) < m(A)$ do
	<p>Determine block size b</p> $\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ \hline B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ \hline C_1 \\ C_2 \end{array} \right)$ <p>where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows</p>
	$C_0 := A_{00}B_0 + A_{10}^T B_1 + A_{20}^T B_2 + C_0$ $C_1 := A_{10}B_0 + A_{11}B_1 + A_{21}^T B_2 + C_1$ $C_2 := A_{20}B_0 + A_{21}B_1 + A_{22}B_2 + C_2$
	$\left(\begin{array}{c c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{cc c} A_{00} & A_{01} & A_{02} \\ \hline A_{10} & A_{11} & A_{12} \\ A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ \hline B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ \hline C_1 \\ C_2 \end{array} \right)$
	endwhile

Algorithm: $C := AB + C$ where A is symmetric and stored in the lower triangular part

$$A \rightarrow \left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right), B \rightarrow \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right), C \rightarrow \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right)$$

where A_{BR} is 0×0 , B_B has 0 rows, C_B has 0 rows

while $m(A_{BR}) < m(A)$ do

Determine block size b

$$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \rightarrow \left(\begin{array}{cc|c} A_{00} & A_{01} & A_{02} \\ A_{10} & A_{11} & A_{12} \\ \hline A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right) \rightarrow \left(\begin{array}{c} B_0 \\ \hline B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right) \rightarrow \left(\begin{array}{c} C_0 \\ \hline C_1 \\ C_2 \end{array} \right)$$

where A_{11} is $b \times b$, B_1 has b rows, C_1 has b rows

$$\begin{aligned} C_0 &:= A_{10}^T B_1 + C_0 \\ C_1 &:= A_{10} B_0 + A_{11} B_1 + C_1 \end{aligned}$$

$$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array} \right) \leftarrow \left(\begin{array}{cc|c} A_{00} & A_{01} & A_{02} \\ A_{10} & A_{11} & A_{12} \\ \hline A_{20} & A_{21} & A_{22} \end{array} \right), \left(\begin{array}{c} B_T \\ \hline B_B \end{array} \right) \leftarrow \left(\begin{array}{c} B_0 \\ \hline B_1 \\ B_2 \end{array} \right), \left(\begin{array}{c} C_T \\ \hline C_B \end{array} \right) \leftarrow \left(\begin{array}{c} C_0 \\ \hline C_1 \\ C_2 \end{array} \right)$$

endwhile