

Topological spaces

2016-01-25 9:00 -0500

X a set

X a set

$P(X)$ the **power set** of X , that is, the set of all subsets of X .

X a set

$P(X)$ the **power set** of X , that is, the set of all subsets of X .

\mathbb{R} the set of real numbers.

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

(T2) If $U_\alpha \in \tau$ for all $\alpha \in I$, then $\cup_\alpha U_\alpha \in \tau$.

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

(T2) If $U_\alpha \in \tau$ for all $\alpha \in I$, then $\bigcup_\alpha U_\alpha \in \tau$.

(T3) If $U_1, U_2 \in \tau$, then $U_1 \cap U_2 \in \tau$.

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

(T2) If $U_\alpha \in \tau$ for all $\alpha \in I$, then $\bigcup_\alpha U_\alpha \in \tau$.

(T3) If $U_1, U_2 \in \tau$, then $U_1 \cap U_2 \in \tau$.

Definition (Topological spaces)

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

(T2) If $U_\alpha \in \tau$ for all $\alpha \in I$, then $\bigcup_\alpha U_\alpha \in \tau$.

(T3) If $U_1, U_2 \in \tau$, then $U_1 \cap U_2 \in \tau$.

Definition (Topological spaces)

1. If τ is a topology on X , the pair (X, τ) is called a *topological space*.

Definitions

Definition (Topology)

A *topology* on X is a subset $\tau \subseteq P(X)$, such that:

(T1) $\emptyset, X \in \tau$.

(T2) If $U_\alpha \in \tau$ for all $\alpha \in I$, then $\bigcup_\alpha U_\alpha \in \tau$.

(T3) If $U_1, U_2 \in \tau$, then $U_1 \cap U_2 \in \tau$.

Definition (Topological spaces)

1. If τ is a topology on X , the pair (X, τ) is called a *topological space*.
2. The elements of τ are called *open sets*.

Examples

- Let X be any set. Then $\tau_d = P(X)$ is a topology, called the *discrete topology*.

Examples

- Let X be any set. Then $\tau_d = P(X)$ is a topology, called the *discrete topology*.
- Let X be any set. Then $\tau_i = \{\emptyset, X\}$ is a topology, called the *indiscrete topology*.

Examples

- Let X be any set. Then $\tau_d = P(X)$ is a topology, called the *discrete topology*.
- Let X be any set. Then $\tau_i = \{\emptyset, X\}$ is a topology, called the *indiscrete topology*.
- Let $X = \{1, 2\}$. Then $\tau = \{\emptyset, X, \{1\}\}$ is a topology, and (X, τ) is called the *Sierpiński space*.

More examples

- Let

More examples

- Let
 - $X = \mathbb{R}$,

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}.$

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}$.
 - Then τ is called the *usual topology* on \mathbb{R} .

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}$.
 - Then τ is called the *usual topology* on \mathbb{R} .
- Let

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}$.
 - Then τ is called the *usual topology* on \mathbb{R} .
- Let
 - (X, d) be any metric space. For $x \in X$, let $B_\epsilon(x) = \{y \in X \mid d(x, y) < \epsilon\}$.

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}$.
 - Then τ is called the *usual topology* on \mathbb{R} .
- Let
 - (X, d) be any metric space. For $x \in X$, let $B_\epsilon(x) = \{y \in X \mid d(x, y) < \epsilon\}$.
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } B_\epsilon(x) \subseteq U\}$.

More examples

- Let
 - $X = \mathbb{R}$,
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } (x-\epsilon, x+\epsilon) \subseteq U\}$.
 - Then τ is called the *usual topology* on \mathbb{R} .
- Let
 - (X, d) be any metric space. For $x \in X$, let $B_\epsilon(x) = \{y \in X \mid d(x, y) < \epsilon\}$.
 - $\tau = \{U \subseteq X \mid \forall x \in U \exists \epsilon > 0, \text{ such that } B_\epsilon(x) \subseteq U\}$.
 - Then τ is called the *metric topology* on X .

Cofinite topology

Example (Cofinite topology)

Let $\tau = \{U \subseteq X \mid X - U \text{ is finite}\} \cup \{\emptyset\}$. Then τ is a topology on X .

Cofinite topology

Example (Cofinite topology)

Let $\tau = \{U \subseteq X \mid X - U \text{ is finite}\} \cup \{\emptyset\}$. Then τ is a topology on X .

Sketch of proof.



Cofinite topology

Example (Cofinite topology)

Let $\tau = \{U \subseteq X \mid X - U \text{ is finite}\} \cup \{\emptyset\}$. Then τ is a topology on X .

Sketch of proof.

- $\emptyset \in \tau$ immediately. $X \in \tau$ because $X - X = \emptyset$ is finite.



Cofinite topology

Example (Cofinite topology)

Let $\tau = \{U \subseteq X \mid X - U \text{ is finite}\} \cup \{\emptyset\}$. Then τ is a topology on X .

Sketch of proof.

- $\emptyset \in \tau$ immediately. $X \in \tau$ because $X - X = \emptyset$ is finite.
- Let $U_\alpha \in \tau$ for all $\alpha \in I$. We need to show that either $\cup_\alpha U_\alpha$ is empty or $X - \cup_\alpha U_\alpha$ is finite.



Cofinite topology

Example (Cofinite topology)

Let $\tau = \{U \subseteq X \mid X - U \text{ is finite}\} \cup \{\emptyset\}$. Then τ is a topology on X .

Sketch of proof.

- $\emptyset \in \tau$ immediately. $X \in \tau$ because $X - X = \emptyset$ is finite.
- Let $U_\alpha \in \tau$ for all $\alpha \in I$. We need to show that either $\cup_\alpha U_\alpha$ is empty or $X - \cup_\alpha U_\alpha$ is finite.
- Let $U_1, U_2 \in \tau$. We need to show that either $U_1 \cap U_2$ is empty or $X - (U_1 \cap U_2)$ is finite.



Exercises

1. Show that if X is any set, there is a metric on X such that the metric topology is the same as the discrete topology.

Exercises

1. Show that if X is any set, there is a metric on X such that the metric topology is the same as the discrete topology.
2. Show that in general there is no metric on X such that the metric topology is the same as the indiscrete topology.

Exercises

1. Show that if X is any set, there is a metric on X such that the metric topology is the same as the discrete topology.
2. Show that in general there is no metric on X such that the metric topology is the same as the indiscrete topology.
3. Show that if τ_α is a topology on X for each $\alpha \in I$, then $\tau = \cap \tau_\alpha$ is a topology on X .

Links

- Topological space - Wikipedia, the free encyclopedia

Links

- Topological space - Wikipedia, the free encyclopedia
- How to get intuition in topology concerning the definitions? - Mathematics Stack Exchange

Links

- Topological space - Wikipedia, the free encyclopedia
- How to get intuition in topology concerning the definitions? - Mathematics Stack Exchange
- Why is a topology made up of 'open' sets? - MathOverflow

Links

- Topological space - Wikipedia, the free encyclopedia
- How to get intuition in topology concerning the definitions? - Mathematics Stack Exchange
- Why is a topology made up of 'open' sets? - MathOverflow
- Why does topology rarely come up outside of topology? - Mathematics Stack Exchange