

Hausdorff spaces

2016-02-11 9:00 -0500

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.
- Any totally ordered set with the order topology is Hausdorff.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.
- Any totally ordered set with the order topology is Hausdorff.
- The product of two Hausdorff spaces is Hausdorff.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.
- Any totally ordered set with the order topology is Hausdorff.
- The product of two Hausdorff spaces is Hausdorff.
- Any subspace of a Hausdorff space is Hausdorff.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.
- Any totally ordered set with the order topology is Hausdorff.
- The product of two Hausdorff spaces is Hausdorff.
- Any subspace of a Hausdorff space is Hausdorff.
- The Sierpiński space is not Hausdorff.

Definition

Definition (Hausdorff space)

We say that the space X is **Hausdorff** if for any $x, y \in X$ such that $x \neq y$, there are disjoint neighborhoods U, V of x, y , respectively.

Examples

- If d is a metric on X , (X, τ_d) is Hausdorff.
- Any totally ordered set with the order topology is Hausdorff.
- The product of two Hausdorff spaces is Hausdorff.
- Any subspace of a Hausdorff space is Hausdorff.
- The Sierpiński space is not Hausdorff.
- Is the cofinite topology on \mathbb{N} Hausdorff?

Properties of Hausdorff spaces

Theorem (Points in Hausdorff)

Every singleton in a Hausdorff space X is a closed set.

Properties of Hausdorff spaces

Theorem (Points in Hausdorff)

Every singleton in a Hausdorff space X is a closed set.

Proof

For $x \in X$, let $y \notin \{x\}$. If U, V are disjoint neighborhoods of x, y respectively, then $V \subseteq X - \{x\}$. Hence $X - \{x\}$ is open.

Properties of Hausdorff spaces

Theorem (Points in Hausdorff)

Every singleton in a Hausdorff space X is a closed set.

Proof

For $x \in X$, let $y \notin \{x\}$. If U, V are disjoint neighborhoods of x, y respectively, then $V \subseteq X - \{x\}$. Hence $X - \{x\}$ is open.

Theorem (Limit points in Hausdorff)

Let X be a Hausdorff space and $A \subseteq X$. Then $x \in X$ is a limit point of A if and only if every neighborhood of x contains infinitely many points of A .

Properties of Hausdorff spaces

Theorem (Points in Hausdorff)

Every singleton in a Hausdorff space X is a closed set.

Proof

For $x \in X$, let $y \notin \{x\}$. If U, V are disjoint neighborhoods of x, y respectively, then $V \subseteq X - \{x\}$. Hence $X - \{x\}$ is open.

Theorem (Limit points in Hausdorff)

Let X be a Hausdorff space and $A \subseteq X$. Then $x \in X$ is a limit point of A if and only if every neighborhood of x contains infinitely many points of A .

Proof.



Exercises

- We say that a space is a T_1 space if for any two different points x, y , each has a neighborhood that does not contain the other point. Show that all singletons are closed in X if and only if X is T_1 .

Exercises

- We say that a space is a T_1 space if for any two different points x, y , each has a neighborhood that does not contain the other point. Show that all singletons are closed in X if and only if X is T_1 .
- Show that If X is Hausdorff, then X is T_1 . Give an example of a T_1 space that is not Hausdorff.

Exercises

- We say that a space is a T_1 space if for any two different points x, y , each has a neighborhood that does not contain the other point. Show that all singletons are closed in X if and only if X is T_1 .
- Show that If X is Hausdorff, then X is T_1 . Give an example of a T_1 space that is not Hausdorff.
- We say that a space is a T_0 space if for any two different points x, y , one of them has a neighborhood that does not contain the other point. Show that a T_1 space is T_0 . Give an example of a T_0 space that is not T_1 . Give an example of a space that is not T_0 .

Links

- Hausdorff space - Wikipedia, the free encyclopedia