

# Connectedness

2015-04-26 11:00

# Separations

## Definition (Separation)

Let  $X$  be a topological space. A **separation** of  $X$  is a pair of open sets  $U, V$  such that:

# Separations

## Definition (Separation)

Let  $X$  be a topological space. A **separation** of  $X$  is a pair of open sets  $U, V$  such that:

- $U \cap V = \emptyset$ ,

# Separations

## Definition (Separation)

Let  $X$  be a topological space. A **separation** of  $X$  is a pair of open sets  $U, V$  such that:

- $U \cap V = \emptyset$ ,
- $U \cup V = X$ ,

# Separations

## Definition (Separation)

Let  $X$  be a topological space. A **separation** of  $X$  is a pair of open sets  $U, V$  such that:

- $U \cap V = \emptyset$ ,
- $U \cup V = X$ ,
- $U \neq \emptyset, V \neq \emptyset$

# Separations

## Definition (Separation)

Let  $X$  be a topological space. A **separation** of  $X$  is a pair of open sets  $U, V$  such that:

- $U \cap V = \emptyset$ ,
- $U \cup V = X$ ,
- $U \neq \emptyset, V \neq \emptyset$

## Definition (Connected space)

A topological space is **connected** if it has no separation.

## Remark

A space is connected if and only if the only subsets that are both open and closed are  $\emptyset$  and  $X$ .

## Remark

A space is connected if and only if the only subsets that are both open and closed are  $\emptyset$  and  $X$ .

This is because if  $U, V$  form a separation, then both are open and closed. Conversely, if  $A \subseteq X$  is not the empty set and not  $X$ , and is open and closed, then  $A, X - A$  form a separation.



# Lemmas

## Lemma

*Let  $X$  be a space and  $Y$  a subspace of  $X$ . Then a separation of  $Y$  is a pair of disjoint nonempty sets  $A, B$  with union  $Y$ , and none containing a limit point of the other.*

# Lemmas

## Lemma

*Let  $X$  be a space and  $Y$  a subspace of  $X$ . Then a separation of  $Y$  is a pair of disjoint nonempty sets  $A, B$  with union  $Y$ , and none containing a limit point of the other.*

# Lemmas

## Lemma

*Let  $X$  be a space and  $Y$  a subspace of  $X$ . Then a separation of  $Y$  is a pair of disjoint nonempty sets  $A, B$  with union  $Y$ , and none containing a limit point of the other.*

- Let  $A, B$  be a separation of  $Y$ . Then both  $A, B$  are open and closed in  $Y$ . We have that the closure of  $A$  in  $Y$  is  $\overline{A} \cap Y$ , and since  $A$  is closed, we have  $A = \overline{A} \cap Y$ . Since  $A, B$  are disjoint, we have  $\overline{A} \cap B = \emptyset$ .

# Lemmas

## Lemma

*Let  $X$  be a space and  $Y$  a subspace of  $X$ . Then a separation of  $Y$  is a pair of disjoint nonempty sets  $A, B$  with union  $Y$ , and none containing a limit point of the other.*

- Let  $A, B$  be a separation of  $Y$ . Then both  $A, B$  are open and closed in  $Y$ . We have that the closure of  $A$  in  $Y$  is  $\overline{A} \cap Y$ , and since  $A$  is closed, we have  $A = \overline{A} \cap Y$ . Since  $A, B$  are disjoint, we have  $\overline{A} \cap B = \emptyset$ .
- Now, suppose  $A, B \subseteq Y$  are disjoint, nonempty, and none containing a limit point of the other. Then  $\overline{A} \cap B = \emptyset$ , hence  $\overline{A} \cap Y = A$ . It follows that  $A$  is closed in  $Y$ , hence  $B$  is open in  $Y$ .

# Examples

- If  $X$  has the indiscrete topology, then  $X$  is connected.

# Examples

- If  $X$  has the indiscrete topology, then  $X$  is connected.
- If  $X$  is a space with only one point, then it is connected.

# Examples

- If  $X$  has the indiscrete topology, then  $X$  is connected.
- If  $X$  is a space with only one point, then it is connected.
- The subspace  $Y = [-1, 0) \cup (0, 1]$  of  $\mathbb{R}$  is not connected.

# Examples

- If  $X$  has the indiscrete topology, then  $X$  is connected.
- If  $X$  is a space with only one point, then it is connected.
- The subspace  $Y = [-1, 0) \cup (0, 1]$  of  $\mathbb{R}$  is not connected.
- $\mathbb{Q}$  as a subspace of  $\mathbb{R}$  is not connected.



# Properties of connected spaces

## Lemma

*If  $A, B$  form a separation of  $X$ , and  $Y \subseteq X$  is connected, then either  $Y \subseteq A$  or  $Y \subseteq B$ .*

# Properties of connected spaces

## Lemma

*If  $A, B$  form a separation of  $X$ , and  $Y \subseteq X$  is connected, then either  $Y \subseteq A$  or  $Y \subseteq B$ .*

We have that  $Y \cap A, Y \cap B$  are both open in  $Y$ , they are disjoint and its union is  $Y$ . Since  $Y$  is connected they cannot be both nonempty, and if  $Y \cap A = \emptyset$ , then  $Y \subseteq B$ .

# Properties of connected spaces

## Lemma

*If  $A, B$  form a separation of  $X$ , and  $Y \subseteq X$  is connected, then either  $Y \subseteq A$  or  $Y \subseteq B$ .*

We have that  $Y \cap A, Y \cap B$  are both open in  $Y$ , they are disjoint and its union is  $Y$ . Since  $Y$  is connected they cannot be both nonempty, and if  $Y \cap A = \emptyset$ , then  $Y \subseteq B$ .

## Lemma

*The union of connected subspaces with a point in common is connected.*

# Properties of connected spaces

## Lemma

*If  $A, B$  form a separation of  $X$ , and  $Y \subseteq X$  is connected, then either  $Y \subseteq A$  or  $Y \subseteq B$ .*

We have that  $Y \cap A, Y \cap B$  are both open in  $Y$ , they are disjoint and its union is  $Y$ . Since  $Y$  is connected they cannot be both nonempty, and if  $Y \cap A = \emptyset$ , then  $Y \subseteq B$ .

## Lemma

*The union of connected subspaces with a point in common is connected.*

Let  $X = \bigcup_{\alpha \in I} A_{\alpha}$  with  $A_{\alpha}$  connected, and  $x_0 \in \bigcap A_{\alpha}$ . Let  $U, V$  be a separation of  $X$ . Suppose  $x_0 \in U$ . By the previous lemma, we must have  $A_{\alpha} \subseteq U$  for all  $\alpha \in I$ . But then  $V = \emptyset$ , which is a contradiction.

# More theorems

## Theorem

*Let  $A \subseteq X$  be connected. Then if  $B \subseteq X$  is such that  $A \subseteq B \subseteq \overline{A}$ , then  $B$  is also connected.*

# More theorems

## Theorem

*Let  $A \subseteq X$  be connected. Then if  $B \subseteq X$  is such that  $A \subseteq B \subseteq \overline{A}$ , then  $B$  is also connected.*

## Proof

Suppose that  $C, D$  is a separation of  $B$ . By a previous lemma, we may assume without loss that  $A \subseteq C$ . But then  $\overline{A} \subseteq \overline{C}$ , and, since  $\overline{C}$  and  $D$  are disjoint and  $B \subseteq \overline{A}$ , we must have  $B \cap D = \emptyset$ , which is a contradiction.  $\square$

## Theorem

*Let  $f: X \rightarrow Y$  be a continuous map, with  $X$  connected. Then  $f(X)$  is connected.*

## Theorem

*Let  $f: X \rightarrow Y$  be a continuous map, with  $X$  connected. Then  $f(X)$  is connected.*

## Proof

Let  $Z = f(X)$ , and consider the surjective map  $f: X \rightarrow Z$ , which is also continuous. Let  $A, B$  be a separation of  $Z$ . Then  $f^{-1}(A), f^{-1}(B)$  would form a separation of  $X$ , which is impossible. Therefore, there is no separation of  $Z$ , hence  $f(X)$  is connected.  $\square$



## Theorem

*The arbitrary product of connected spaces is connected.*

## Theorem

*The arbitrary product of connected spaces is connected.*

## Theorem

*The arbitrary product of connected spaces is connected.*

- For the proof, we first prove that if  $X, Y$  are connected, then  $X \times Y$  is connected.

## Theorem

*The arbitrary product of connected spaces is connected.*

- For the proof, we first prove that if  $X, Y$  are connected, then  $X \times Y$  is connected.
- Let  $(a, b) \in X \times Y$ . Then  $X \times \{b\}$  and  $\{x\} \times Y$  are connected, for all  $y \in Y$ .

## Theorem

*The arbitrary product of connected spaces is connected.*

- For the proof, we first prove that if  $X, Y$  are connected, then  $X \times Y$  is connected.
- Let  $(a, b) \in X \times Y$ . Then  $X \times \{b\}$  and  $\{x\} \times Y$  are connected, for all  $y \in Y$ .
- Hence  $T_x = (X \times \{b\}) \cup (\{x\} \times Y)$  is connected, since it is the union of two connected spaces with a point in common.

## Theorem

*The arbitrary product of connected spaces is connected.*

- For the proof, we first prove that if  $X, Y$  are connected, then  $X \times Y$  is connected.
- Let  $(a, b) \in X \times Y$ . Then  $X \times \{b\}$  and  $\{x\} \times Y$  are connected, for all  $y \in Y$ .
- Hence  $T_x = (X \times \{b\}) \cup (\{x\} \times Y)$  is connected, since it is the union of two connected spaces with a point in common.
- Then,  $X \times Y = \bigcup_{x \in X} T_x$  is the union of connected spaces with the point  $(a, b)$  in common.

## Theorem

*The arbitrary product of connected spaces is connected.*

- For the proof, we first prove that if  $X, Y$  are connected, then  $X \times Y$  is connected.
- Let  $(a, b) \in X \times Y$ . Then  $X \times \{b\}$  and  $\{x\} \times Y$  are connected, for all  $y \in Y$ .
- Hence  $T_x = (X \times \{b\}) \cup (\{x\} \times Y)$  is connected, since it is the union of two connected spaces with a point in common.
- Then,  $X \times Y = \bigcup_{x \in X} T_x$  is the union of connected spaces with the point  $(a, b)$  in common.
- By induction, we obtain that the product of any finite number of connected spaces is connected.

- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .



- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .
- Given  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ , we define  $X(\alpha_1, \dots, \alpha_n)$  as the subspace of  $X$  with points such that  $x_\alpha = b_\alpha$  for  $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$ .

- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .
- Given  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ , we define  $X(\alpha_1, \dots, \alpha_n)$  as the subspace of  $X$  with points such that  $x_\alpha = b_\alpha$  for  $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$ .
- We have that  $X(\alpha_1, \dots, \alpha_n)$  is homeomorphic to  $X_{\alpha_1} \times \dots \times X_{\alpha_n}$ , and so it is connected.

- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .
- Given  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ , we define  $X(\alpha_1, \dots, \alpha_n)$  as the subspace of  $X$  with points such that  $x_\alpha = b_\alpha$  for  $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$ .
- We have that  $X(\alpha_1, \dots, \alpha_n)$  is homeomorphic to  $X_{\alpha_1} \times \dots \times X_{\alpha_n}$ , and so it is connected.
- Let  $Y$  be the subspace of  $X$  that is the union of all  $X(\alpha_1, \dots, \alpha_n)$ , for  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$  finite. Then  $Y$  is connected, as is the union of connected spaces with the point  $b$  in common.

- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .
- Given  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ , we define  $X(\alpha_1, \dots, \alpha_n)$  as the subspace of  $X$  with points such that  $x_\alpha = b_\alpha$  for  $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$ .
- We have that  $X(\alpha_1, \dots, \alpha_n)$  is homeomorphic to  $X_{\alpha_1} \times \dots \times X_{\alpha_n}$ , and so it is connected.
- Let  $Y$  be the subspace of  $X$  that is the union of all  $X(\alpha_1, \dots, \alpha_n)$ , for  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$  finite. Then  $Y$  is connected, as is the union of connected spaces with the point  $b$  in common.
- We finally prove that  $\overline{Y} = X$ . Let  $x = (x_\alpha) \in X$ , and  $U = \prod U_\alpha$  be a basis element of the product topology. We have that  $U = X_\alpha$  for all  $\alpha \in I - \{\alpha_1, \dots, \alpha_n\}$ .

- Now, let  $X_\alpha$  be a connected space, for  $\alpha \in I$ . We want to prove that  $\prod_\alpha X_\alpha$  is connected. Choose  $b = (b_\alpha) \in X$ .
- Given  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ , we define  $X(\alpha_1, \dots, \alpha_n)$  as the subspace of  $X$  with points such that  $x_\alpha = b_\alpha$  for  $\alpha \notin \{\alpha_1, \dots, \alpha_n\}$ .
- We have that  $X(\alpha_1, \dots, \alpha_n)$  is homeomorphic to  $X_{\alpha_1} \times \dots \times X_{\alpha_n}$ , and so it is connected.
- Let  $Y$  be the subspace of  $X$  that is the union of all  $X(\alpha_1, \dots, \alpha_n)$ , for  $\{\alpha_1, \dots, \alpha_n\} \subseteq I$  finite. Then  $Y$  is connected, as is the union of connected spaces with the point  $b$  in common.
- We finally prove that  $\overline{Y} = X$ . Let  $x = (x_\alpha) \in X$ , and  $U = \prod U_\alpha$  be a basis element of the product topology. We have that  $U = X_\alpha$  for all  $\alpha \in I - \{\alpha_1, \dots, \alpha_n\}$ .
- Let  $y = (y_\alpha)$  defined as  $y_\alpha = x_\alpha$  for  $\alpha \in \{\alpha_1, \dots, \alpha_n\}$ , and  $y_\alpha = b_\alpha$  otherwise. Then  $y \in X(\alpha_1, \dots, \alpha_n)$ , so  $Y \cap U \neq \emptyset$ .  
□

# Linear continuum

## Definition

A totally ordered set  $L$  is called a **linear continuum** if it satisfies the following:

# Linear continuum

## Definition

A totally ordered set  $L$  is called a **linear continuum** if it satisfies the following:

- $L$  has the least upper bound property.

# Linear continuum

## Definition

A totally ordered set  $L$  is called a **linear continuum** if it satisfies the following:

- $L$  has the least upper bound property.
- If  $x < y$ , there is  $z$  such that  $x < z < y$ .



# Linear continuum

## Definition

A totally ordered set  $L$  is called a **linear continuum** if it satisfies the following:

- $L$  has the least upper bound property.
- If  $x < y$ , there is  $z$  such that  $x < z < y$ .

## Theorem

*If  $L$  is a linear continuum, then  $L$  is connected in the order topology. (Hence, every interval and every ray in  $L$  is connected.)*

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.
- Suppose that  $Y = A \cup B$ , with  $A, B$  open in  $Y$ , disjoint and nonempty. Let  $a \in A$ ,  $b \in B$ , and assume that  $a < b$ .

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.
- Suppose that  $Y = A \cup B$ , with  $A, B$  open in  $Y$ , disjoint and nonempty. Let  $a \in A$ ,  $b \in B$ , and assume that  $a < b$ .
- Since  $Y$  is convex, we have  $[a, b] \subseteq Y$ . Then  $[a, b]$  is the union of the disjoint nonempty sets  $A_0, B_0$  given by:

$$A_0 = A \cap [a, b] \quad B_0 = B \cap [a, b]$$

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.
- Suppose that  $Y = A \cup B$ , with  $A, B$  open in  $Y$ , disjoint and nonempty. Let  $a \in A$ ,  $b \in B$ , and assume that  $a < b$ .
- Since  $Y$  is convex, we have  $[a, b] \subseteq Y$ . Then  $[a, b]$  is the union of the disjoint nonempty sets  $A_0, B_0$  given by:

$$A_0 = A \cap [a, b] \quad B_0 = B \cap [a, b]$$

- Note that  $A_0, B_0$  are open in  $[a, b]$ , and thus form a separation of  $[a, b]$ .

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.
- Suppose that  $Y = A \cup B$ , with  $A, B$  open in  $Y$ , disjoint and nonempty. Let  $a \in A$ ,  $b \in B$ , and assume that  $a < b$ .
- Since  $Y$  is convex, we have  $[a, b] \subseteq Y$ . Then  $[a, b]$  is the union of the disjoint nonempty sets  $A_0, B_0$  given by:

$$A_0 = A \cap [a, b] \quad B_0 = B \cap [a, b]$$

- Note that  $A_0, B_0$  are open in  $[a, b]$ , and thus form a separation of  $[a, b]$ .
- The interval  $[a, b]$  is a subspace of  $L$ . Therefore the topology on  $[a, b]$  is the order topology.

# Proof

- We say that a subset  $Y \subseteq L$  is **convex** if  $a, b \in Y$  with  $a < b$  implies  $[a, b] \subseteq Y$ . We will prove that any convex subset  $Y$  of  $L$  is connected.
- Suppose that  $Y = A \cup B$ , with  $A, B$  open in  $Y$ , disjoint and nonempty. Let  $a \in A$ ,  $b \in B$ , and assume that  $a < b$ .
- Since  $Y$  is convex, we have  $[a, b] \subseteq Y$ . Then  $[a, b]$  is the union of the disjoint nonempty sets  $A_0, B_0$  given by:

$$A_0 = A \cap [a, b] \quad B_0 = B \cap [a, b]$$

- Note that  $A_0, B_0$  are open in  $[a, b]$ , and thus form a separation of  $[a, b]$ .
- The interval  $[a, b]$  is a subspace of  $L$ . Therefore the topology on  $[a, b]$  is the order topology.
- Let  $c = \sup A_0$ . Then  $c \in [a, b]$ .

## Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .



## Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d \in [a, b]$  such that  $(d, c] \subseteq B_0$ .

## Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d \in [a, b]$  such that  $(d, c] \subseteq B_0$ .
- Hence  $d$  is an upper bound of  $A_0$ , which contradicts the choice of  $c$ .

## Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d \in [a, b]$  such that  $(d, c] \subseteq B_0$ .
- Hence  $d$  is an upper bound of  $A_0$ , which contradicts the choice of  $c$ .
- Then  $c \in A_0$ . Given that  $c \neq b$ , we have that either  $c = a$  or  $c \in (a, b)$ .

# Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d \in [a, b]$  such that  $(d, c] \subseteq B_0$ .
- Hence  $d$  is an upper bound of  $A_0$ , which contradicts the choice of  $c$ .
- Then  $c \in A_0$ . Given that  $c \neq b$ , we have that either  $c = a$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d$  such that  $[c, d) \subseteq A_0$ .

## Proof (continuation)

- Suppose  $c \in B_0$ . Given that  $c \neq a$ , we have that either  $c = b$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d \in [a, b]$  such that  $(d, c] \subseteq B_0$ .
- Hence  $d$  is an upper bound of  $A_0$ , which contradicts the choice of  $c$ .
- Then  $c \in A_0$ . Given that  $c \neq b$ , we have that either  $c = a$  or  $c \in (a, b)$ .
- In any case, we have that there is  $d$  such that  $[c, d) \subseteq A_0$ .
- By Property 2 of linear continuum, there is  $z$  such that  $c < z < d$ . But this contradicts that  $c$  is upper bound of  $A_0$ . □

# Connected subsets of $\mathbb{R}$

## Corollary

$\mathbb{R}$  is connected, and so is every interval and ray in  $\mathbb{R}$ .

# Connected subsets of $\mathbb{R}$

## Corollary

$\mathbb{R}$  is connected, and so is every interval and ray in  $\mathbb{R}$ .

## Theorem

*If  $X$  and  $Y$  are linear continuum, and  $Y$  is bounded above and below, then  $X \times Y$  is a linear continuum with dictionary order.*

# Intermediate value theorem

## Theorem

*Let  $f: X \rightarrow Y$  be a continuous map, where  $X$  is connected and  $Y$  is ordered and has the order topology. Suppose that  $a, b \in X$  and  $y \in Y$  are such that  $f(a) < y < f(b)$ . Then there is  $c \in X$  such that  $f(c) = y$ .*



## Definition

## Definition

- A **path** on a space  $X$  from  $x \in X$  to  $y \in X$  is a continuous function  $f: [a, b] \rightarrow X$  from some interval  $[a, b] \subseteq \mathbb{R}$ , such that  $f(a) = x, f(b) = y$ .

## Definition

- A **path** on a space  $X$  from  $x \in X$  to  $y \in X$  is a continuous function  $f: [a, b] \rightarrow X$  from some interval  $[a, b] \subseteq \mathbb{R}$ , such that  $f(a) = x$ ,  $f(b) = y$ .
- A space  $X$  is **path-connected** if for every  $x, y \in X$  there is a path from  $x$  to  $y$ .

## Definition

- A **path** on a space  $X$  from  $x \in X$  to  $y \in X$  is a continuous function  $f: [a, b] \rightarrow X$  from some interval  $[a, b] \subseteq \mathbb{R}$ , such that  $f(a) = x$ ,  $f(b) = y$ .
- A space  $X$  is **path-connected** if for every  $x, y \in X$  there is a path from  $x$  to  $y$ .

## Theorem

*If  $X$  is path-connected, then it is connected.*

## Definition

- A **path** on a space  $X$  from  $x \in X$  to  $y \in X$  is a continuous function  $f: [a, b] \rightarrow X$  from some interval  $[a, b] \subseteq \mathbb{R}$ , such that  $f(a) = x$ ,  $f(b) = y$ .
- A space  $X$  is **path-connected** if for every  $x, y \in X$  there is a path from  $x$  to  $y$ .

## Theorem

*If  $X$  is path-connected, then it is connected.*

## Examples

The converse is not true, see  $[0, 1] \times [0, 1]$  with dictionary order and the *deleted comb space*.