

# Components and local connectedness

2016-05-05 10:00

# Definition

## Definition (Equivalence relation)

Let  $X$  be a topological space. We define a relation  $\sim$  on  $X$  by declaring  $x \sim y$  if there is  $Y \subseteq X$  connected, such that  $x, y \in Y$ .

# Definition

## Definition (Equivalence relation)

Let  $X$  be a topological space. We define a relation  $\sim$  on  $X$  by declaring  $x \sim y$  if there is  $Y \subseteq X$  connected, such that  $x, y \in Y$ .

## Definition (Connected components)

One can prove that  $\sim$  is an equivalence relation. The equivalence classes are called **connected components** of  $X$ .

# Properties

## Theorem

*The connected components of  $X$  have the following properties:*

# Properties

## Theorem

*The connected components of  $X$  have the following properties:*

- *They are connected, disjoint subsets of  $X$  with union  $X$ .*

# Properties

## Theorem

*The connected components of  $X$  have the following properties:*

- *They are connected, disjoint subsets of  $X$  with union  $X$ .*
- *Each connected nonempty subset of  $X$  intersects exactly one of them.*

# Properties

## Theorem

*The connected components of  $X$  have the following properties:*

- *They are connected, disjoint subsets of  $X$  with union  $X$ .*
- *Each connected nonempty subset of  $X$  intersects exactly one of them.*
- *They are closed, and if there are finitely many components, they are also open.*

# Definition

## A new equivalence relation

We define a relation  $\sim$  on the topological space  $X$ , by setting  $x \sim y$  if there is a path from  $x$  to  $y$ .



# Definition

## A new equivalence relation

We define a relation  $\sim$  on the topological space  $X$ , by setting  $x \sim y$  if there is a path from  $x$  to  $y$ .

## Definition (Path components)

One can prove that  $\sim$  is an equivalence relation. The equivalence classes are called **path components** of  $X$ .

# Properties

## Theorem

*The path components of  $X$  have the following properties:*

# Properties

## Theorem

*The path components of  $X$  have the following properties:*

- *They are path-connected, disjoint subsets of  $X$  with union  $X$ .*


# Properties

## Theorem

*The path components of  $X$  have the following properties:*

- *They are path-connected, disjoint subsets of  $X$  with union  $X$ .*
- *Each nonempty path-connected subset of  $X$  intersects exactly one of them.*

# The topologist's sine curve



$$X = \{(x, \sin \frac{1}{x}) \mid x > 0\} \cup \{(0, x) \mid x \in [-1, 1]\}$$

# Definition

Locally connected space

# Definition

## Locally connected space

- $X$  is **locally connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a connected neighborhood of  $x$  contained in  $U$ .

# Definition

## Locally connected space

- $X$  is **locally connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a connected neighborhood of  $x$  contained in  $U$ .
- If  $X$  is locally connected at every point, then we say that  $X$  is **locally connected**.



# Definition

## Locally connected space

- $X$  is **locally connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a connected neighborhood of  $x$  contained in  $U$ .
- If  $X$  is locally connected at every point, then we say that  $X$  is **locally connected**.

## Locally path connected space

# Definition

## Locally connected space

- $X$  is **locally connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a connected neighborhood of  $x$  contained in  $U$ .
- If  $X$  is locally connected at every point, then we say that  $X$  is **locally connected**.

## Locally path connected space

- $X$  is **locally path connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a path connected neighborhood of  $x$  contained in  $U$ .

# Definition

## Locally connected space

- $X$  is **locally connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a connected neighborhood of  $x$  contained in  $U$ .
- If  $X$  is locally connected at every point, then we say that  $X$  is **locally connected**.

## Locally path connected space

- $X$  is **locally path connected at  $x \in X$**  if for any neighborhood  $U$  of  $x$ , there is a path connected neighborhood of  $x$  contained in  $U$ .
- If  $X$  is locally path connected at every point, then we say that  $X$  is **locally path connected**.

# Examples

:Bexampleblock:

# Examples

:Bexampleblock:

- Any interval in  $\mathbb{R}$  is connected and locally connected.

# Examples

:Bexampleblock:

- Any interval in  $\mathbb{R}$  is connected and locally connected.
- The subspace  $[0, 1] \cup [2, 3] \subseteq \mathbb{R}$  is locally connected and not connected.

# Examples

:Bexampleblock:

- Any interval in  $\mathbb{R}$  is connected and locally connected.
- The subspace  $[0, 1] \cup [2, 3] \subseteq \mathbb{R}$  is locally connected and not connected.
- The topologist's sine curve is connected but not locally connected.

# Examples

:Bexampleblock:

- Any interval in  $\mathbb{R}$  is connected and locally connected.
- The subspace  $[0, 1] \cup [2, 3] \subseteq \mathbb{R}$  is locally connected and not connected.
- The topologist's sine curve is connected but not locally connected.
- $\mathbb{Q}$  as a subspace of  $\mathbb{R}$  is neither connected nor locally connected.



## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

- Suppose that  $X$  is locally connected, and let  $C$  be a component of the open set  $U \subseteq X$ .

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

- Suppose that  $X$  is locally connected, and let  $C$  be a component of the open set  $U \subseteq X$ .
- Let  $x \in C$ , and choose  $V$  a connected neighborhood of  $x$  with  $V \subseteq U$ .

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

- Suppose that  $X$  is locally connected, and let  $C$  be a component of the open set  $U \subseteq X$ .
- Let  $x \in C$ , and choose  $V$  a connected neighborhood of  $x$  with  $V \subseteq U$ .
- Since  $V$  is connected, it must be contained in  $C$ .

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

- Suppose that  $X$  is locally connected, and let  $C$  be a component of the open set  $U \subseteq X$ .
- Let  $x \in C$ , and choose  $V$  a connected neighborhood of  $x$  with  $V \subseteq U$ .
- Since  $V$  is connected, it must be contained in  $C$ .
- Now suppose the components of open sets in  $X$  are open.

## Theorem

*A space  $X$  is locally connected if and only if for every open set  $U$  of  $X$ , each component of  $U$  is open in  $X$ .*

## Proof

- Suppose that  $X$  is locally connected, and let  $C$  be a component of the open set  $U \subseteq X$ .
- Let  $x \in C$ , and choose  $V$  a connected neighborhood of  $x$  with  $V \subseteq U$ .
- Since  $V$  is connected, it must be contained in  $C$ .
- Now suppose the components of open sets in  $X$  are open.
- Let  $x \in X$  and a neighborhood  $U$  of  $x$ . If  $C$  is the component of  $U$  that contains  $x$ , then  $C$  is a connected neighborhood of  $x$ .



The proof of the following theorem is similar and left as an exercise.

### Theorem

*A space  $X$  is locally path connected if and only if for every open set  $U$  of  $X$ , each path component of  $U$  is open in  $X$ .*



# Components and path components

Theorem

# Components and path components

## Theorem

- *For any space  $X$ , each path component of  $X$  is contained in a component of  $X$ .*

# Components and path components

## Theorem

- *For any space  $X$ , each path component of  $X$  is contained in a component of  $X$ .*
- *If  $X$  is locally path connected, the components and the path components are the same.*

# Components and path components

## Theorem

- *For any space  $X$ , each path component of  $X$  is contained in a component of  $X$ .*
- *If  $X$  is locally path connected, the components and the path components are the same.*

## Proof

# Components and path components

## Theorem

- *For any space  $X$ , each path component of  $X$  is contained in a component of  $X$ .*
- *If  $X$  is locally path connected, the components and the path components are the same.*

## Proof

- Let  $C$  be a component of  $X$ , let  $x \in C$ , and let  $P$  be the path component of  $X$  containing  $x$ . Since  $P$  is connected,  $P \subseteq C$ .

# Components and path components

## Theorem

- *For any space  $X$ , each path component of  $X$  is contained in a component of  $X$ .*
- *If  $X$  is locally path connected, the components and the path components are the same.*

## Proof

- Let  $C$  be a component of  $X$ , let  $x \in C$ , and let  $P$  be the path component of  $X$  containing  $x$ . Since  $P$  is connected,  $P \subseteq C$ .
- Suppose now that  $X$  is locally path connected, and we wish to prove  $P = C$ . Assume that  $P \subsetneq C$ .

## Proof (continuation)

## Proof (continuation)

- Let  $S$  be the union of all the path components of  $X$  different from  $P$  that intersect  $C$ . Since such components must lie in  $C$ , we have  $C = P \cup S$ .



## Proof (continuation)

- Let  $S$  be the union of all the path components of  $X$  different from  $P$  that intersect  $C$ . Since such components must lie in  $C$ , we have  $C = P \cup S$ .
- Since  $X$  is locally path connected, each path component of  $X$  is open in  $X$ . Hence  $P$  and  $S$  are open, disjoint and nonempty sets with union  $C$ . This contradicts that  $C$  is connected. □