

The order topology

2016-02-01 9:00 -0500

Total order

Definition (Total order)

A relation $<$ on a set X is called a **total (or linear) order** if the following is satisfied:

Total order

Definition (Total order)

A relation $<$ on a set X is called a **total (or linear) order** if the following is satisfied:

(O1) For every $x, y \in X$ with $x \neq y$, we have that $x < y$ or $y < x$ (**comparability**).

Total order

Definition (Total order)

A relation $<$ on a set X is called a **total (or linear) order** if the following is satisfied:

- (O1)** For every $x, y \in X$ with $x \neq y$, we have that $x < y$ or $y < x$ (**comparability**).
- (O2)** For every $x \in X$, we have that $x \not< x$ (**nonreflexivity**).

Total order

Definition (Total order)

A relation $<$ on a set X is called a **total (or linear) order** if the following is satisfied:

- (O1)** For every $x, y \in X$ with $x \neq y$, we have that $x < y$ or $y < x$ (**comparability**).
- (O2)** For every $x \in X$, we have that $x \not< x$ (**nonreflexivity**).
- (O3)** if $x < y$ and $y < z$, then $x < z$ (**transitivity**).

Total order

Definition (Total order)

A relation $<$ on a set X is called a **total (or linear) order** if the following is satisfied:

- (O1)** For every $x, y \in X$ with $x \neq y$, we have that $x < y$ or $y < x$ (**comparability**).
- (O2)** For every $x \in X$, we have that $x \not< x$ (**nonreflexivity**).
- (O3)** if $x < y$ and $y < z$, then $x < z$ (**transitivity**).

Definition (Totally ordered set)

In this case, the pair $(X, <)$ is called a **totally ordered set**. If $x < y$ or $x = y$, we write $x \leq y$. An example of a total order is given by $X = \mathbb{R}$ with usual order.

Intervals

Definition (Intervals)

If X is a totally ordered set, and $a, b \in X$, we will use the following notation for **intervals**:

$$[a, b] = \{x \in X \mid a \leq x \leq b\}$$

$$(a, b) = \{x \in X \mid a < x < b\}$$

$$(a, b] = \{x \in X \mid a < x \leq b\}$$

$$[a, b) = \{x \in X \mid a \leq x < b\}$$

$$(a, \infty) = \{x \in X \mid x > a\}$$

$$(-\infty, b) = \{x \in X \mid x < b\}$$

The order topology

Definition (Order topology)

Let X be a totally ordered set with more than one element.
Let \mathcal{S} be the collection:

$$\mathcal{S} = \{(-\infty, a) \mid a \in X\} \cup \{(b, \infty) \mid b \in X\}.$$

Then \mathcal{S} is subbase for a topology on X , called **order topology**.

Base for the order topology

Lemma (Base for the order topology)

Let X be a totally ordered set with more than one element. Let \mathcal{B} be the collection of all intervals of the form (a, b) , together with those of the form $[m, b)$, in case X has a minimum m , and those of the form $(a, M]$, in case X has a maximum M . Then \mathcal{B} is a base for the order topology on X .

Base for the order topology

Lemma (Base for the order topology)

Let X be a totally ordered set with more than one element. Let \mathcal{B} be the collection of all intervals of the form (a, b) , together with those of the form $[m, b)$, in case X has a minimum m , and those of the form $(a, M]$, in case X has a maximum M . Then \mathcal{B} is a base for the order topology on X .

Proof

The set of finite intersections of the intervals in the subbasis can be obtained as union of elements of the intervals described. \square

Dictionary order

Definition (Dictionary order)

Let X and Y be totally ordered sets. We define an order relation on the cartesian product $X \times Y$ by:

$$(x, y) < (x', y'),$$

if either:

Dictionary order

Definition (Dictionary order)

Let X and Y be totally ordered sets. We define an order relation on the cartesian product $X \times Y$ by:

$$(x, y) < (x', y'),$$

if either:

- $x < x'$, or

Dictionary order

Definition (Dictionary order)

Let X and Y be totally ordered sets. We define an order relation on the cartesian product $X \times Y$ by:

$$(x, y) < (x', y'),$$

if either:

- $x < x'$, or
- $x = x'$ and $y < y'$.

Dictionary order

Definition (Dictionary order)

Let X and Y be totally ordered sets. We define an order relation on the cartesian product $X \times Y$ by:

$$(x, y) < (x', y'),$$

if either:

- $x < x'$, or
- $x = x'$ and $y < y'$.

Lemma (The dictionary order is a total order)

*The order on $X \times Y$ just described is a total order, called the **dictionary order**.*

Proof that the dictionary order is a total order

- Let $(x, y) \neq (x', y')$ with $(x, y), (x', y') \in X \times Y$. If $x \neq x'$, then one of the pairs is less than the other. Otherwise $x = x'$, and since the pairs are distinct, we must have $y \neq y'$, and so (O1) follows.

Proof that the dictionary order is a total order

- Let $(x, y) \neq (x', y')$ with $(x, y), (x', y') \in X \times Y$. If $x \neq x'$, then one of the pairs is less than the other. Otherwise $x = x'$, and since the pairs are distinct, we must have $y \neq y'$, and so (O1) follows.
- If we had $(x, y) < (x, y)$, then it would follow that either $x < x$ or $x = x, y < y$. Since none of these is possible, we conclude that (O2) is true.

Proof that the dictionary order is a total order

- Let $(x, y) \neq (x', y')$ with $(x, y), (x', y') \in X \times Y$. If $x \neq x'$, then one of the pairs is less than the other. Otherwise $x = x'$, and since the pairs are distinct, we must have $y \neq y'$, and so (O1) follows.
- If we had $(x, y) < (x, y)$, then it would follow that either $x < x$ or $x = x, y < y$. Since none of these is possible, we conclude that (O2) is true.
- The proof of (O3) is left as an exercise.

Examples

- The order topology on \mathbb{R} is the same as the standard topology.

Examples

- The order topology on \mathbb{R} is the same as the standard topology.
- The order topology on $\mathbb{R} \times \mathbb{R}$ with the dictionary order is different from the usual metric topology on \mathbb{R}^2 .

Examples

- The order topology on \mathbb{R} is the same as the standard topology.
- The order topology on $\mathbb{R} \times \mathbb{R}$ with the dictionary order is different from the usual metric topology on \mathbb{R}^2 .
- The order topology on $\{0, 1\} \times \mathbb{N}$ has almost all singletons as open sets.

Exercises

1. Let $(X, <)$ be a totally ordered set, and let $Y \subseteq X$. If, for $a, b \in Y$ we define $a <' b$ whenever $a < b$ in X , show that $(Y, <')$ is a totally ordered set.

Exercises

1. Let $(X, <)$ be a totally ordered set, and let $Y \subseteq X$. If, for $a, b \in Y$ we define $a <' b$ whenever $a < b$ in X , show that $(Y, <')$ is a totally ordered set.
2. If X is a totally ordered set, and $Y \subseteq X$, show that Y can have at most one smallest element.

Exercises

1. Let $(X, <)$ be a totally ordered set, and let $Y \subseteq X$. If, for $a, b \in Y$ we define $a <' b$ whenever $a < b$ in X , show that $(Y, <')$ is a totally ordered set.
2. If X is a totally ordered set, and $Y \subseteq X$, show that Y can have at most one smallest element.
3. If X is a totally ordered set, and we have that $x < y$ and there are no $z \in X$ such that $x < z < y$, we say that y is an **immediate succesor** of x . Show that any $x \in X$ has at most one immediate succesor.

Exercises

1. Let $(X, <)$ be a totally ordered set, and let $Y \subseteq X$. If, for $a, b \in Y$ we define $a <' b$ whenever $a < b$ in X , show that $(Y, <')$ is a totally ordered set.
2. If X is a totally ordered set, and $Y \subseteq X$, show that Y can have at most one smallest element.
3. If X is a totally ordered set, and we have that $x < y$ and there are no $z \in X$ such that $x < z < y$, we say that y is an **immediate succesor** of x . Show that any $x \in X$ has at most one immediate succesor.
4. Let $(X, <)$ be a totally ordered set with no minimum element, and let $\mathcal{C} = \{(a, \infty) \mid a \in X\}$. Show that \mathcal{C} is a basis for a topology on X . Is $\tau_{\mathcal{C}}$ in general, the same as the order topology?

Links

- Total order - Wikipedia, the free encyclopedia

Links

- Total order - Wikipedia, the free encyclopedia
- Order topology - Wikipedia, the free encyclopedia