

# The subspace topology

2016-02-02 11:00 -0500

# Definition

## Definition (Subspace topology)

Let  $(X, \tau)$  be a topological space and  $Y \subseteq X$ . Then

$$\tau_Y = \{Y \cap U \mid U \in \tau\}$$

is a topology on  $Y$ , called **subspace topology**.

# Definition

## Definition (Subspace topology)

Let  $(X, \tau)$  be a topological space and  $Y \subseteq X$ . Then

$$\tau_Y = \{Y \cap U \mid U \in \tau\}$$

is a topology on  $Y$ , called **subspace topology**.

## Proof

# Definition

## Definition (Subspace topology)

Let  $(X, \tau)$  be a topological space and  $Y \subseteq X$ . Then

$$\tau_Y = \{Y \cap U \mid U \in \tau\}$$

is a topology on  $Y$ , called **subspace topology**.

## Proof

- Since  $\emptyset = Y \cap \emptyset$ , and  $Y = Y \cap X$ , we have (T1).

# Definition

## Definition (Subspace topology)

Let  $(X, \tau)$  be a topological space and  $Y \subseteq X$ . Then

$$\tau_Y = \{Y \cap U \mid U \in \tau\}$$

is a topology on  $Y$ , called **subspace topology**.

## Proof

- Since  $\emptyset = Y \cap \emptyset$ , and  $Y = Y \cap X$ , we have (T1).
- Let  $V_\alpha \in \tau_Y$  for  $\alpha \in I$ . For each  $\alpha \in I$ , there is  $U_\alpha \in \tau$  such that  $V_\alpha = Y \cap U_\alpha$ .

# Definition

## Definition (Subspace topology)

Let  $(X, \tau)$  be a topological space and  $Y \subseteq X$ . Then

$$\tau_Y = \{Y \cap U \mid U \in \tau\}$$

is a topology on  $Y$ , called **subspace topology**.

## Proof

- Since  $\emptyset = Y \cap \emptyset$ , and  $Y = Y \cap X$ , we have (T1).
- Let  $V_\alpha \in \tau_Y$  for  $\alpha \in I$ . For each  $\alpha \in I$ , there is  $U_\alpha \in \tau$  such that  $V_\alpha = Y \cap U_\alpha$ .
- Then:

$$Y \cap (\cup U_\alpha) = \cup (Y \cap U_\alpha) = \cup V_\alpha,$$

and we obtain (T2). The proof of (T3) is an exercise.

## Open sets relative to $Y$

If  $Y$  is a subspace of  $X$ , and  $U \subseteq Y$ , it could be that  $U$  is open in  $Y$  but not open in  $X$ .

# Base of the subspace topology

## Lemma

Let  $\mathcal{B}$  be a base for the topology of  $X$ . Then

$$\mathcal{B}_Y = \{Y \cap B \mid B \in \mathcal{B}\}$$

is a base for the subspace topology on  $Y$ .



# Base of the subspace topology

## Lemma

Let  $\mathcal{B}$  be a base for the topology of  $X$ . Then

$$\mathcal{B}_Y = \{Y \cap B \mid B \in \mathcal{B}\}$$

is a base for the subspace topology on  $Y$ .

## Proof

Let  $U$  be open in  $X$  and  $y \in Y \cap U$ . Let  $B \in \mathcal{B}$  such that  $y \in B \subseteq U$ . Then  $Y \cap B \in \mathcal{B}_Y$ , and  $y \in Y \cap B \subseteq Y \cap U$ .  $\square$

# Example

Subspace of the real line

# Example

## Subspace of the real line

- Consider  $Y = [0, 1] \subseteq X = \mathbb{R}$ , where  $X$  has the standard topology.

# Example

## Subspace of the real line

- Consider  $Y = [0, 1] \subseteq X = \mathbb{R}$ , where  $X$  has the standard topology.
- A base for the subspace topology on  $Y$  is the collection of all sets of the form  $(a, b) \cap Y$ .

# Example

## Subspace of the real line

- Consider  $Y = [0, 1] \subseteq X = \mathbb{R}$ , where  $X$  has the standard topology.
- A base for the subspace topology on  $Y$  is the collection of all sets of the form  $(a, b) \cap Y$ .
- They all are of one of the following forms:  $(a, b)$ ,  $[0, b)$ ,  $(a, 1]$ ,  $Y$ ,  $\emptyset$ .

# Example

## Subspace of the real line

- Consider  $Y = [0, 1] \subseteq X = \mathbb{R}$ , where  $X$  has the standard topology.
- A base for the subspace topology on  $Y$  is the collection of all sets of the form  $(a, b) \cap Y$ .
- They all are of one of the following forms:  $(a, b)$ ,  $[0, b)$ ,  $(a, 1]$ ,  $Y$ ,  $\emptyset$ .
- We obtain that these elements generate the order topology on  $[0, 1]$ .

# Example

Subspace and order topology not always the same

# Example

Subspace and order topology not always the same

- Consider now  $Y = [0, 1) \cup \{2\} \subseteq X = \mathbb{R}$ .



# Example

Subspace and order topology not always the same

- Consider now  $Y = [0, 1) \cup \{2\} \subseteq X = \mathbb{R}$ .
- In the subspace topology, the set  $\{2\}$  is open.

# Example

Subspace and order topology not always the same

- Consider now  $Y = [0, 1) \cup \{2\} \subseteq X = \mathbb{R}$ .
- In the subspace topology, the set  $\{2\}$  is open.
- But in the order topology on  $Y$ , considering that 2 is  $\max Y$ , any open set that contains 2 has to contain other points.

# Order topology in intervals

## Lemma (Order topology in intervals)

*If  $X$  is an ordered set that has the order topology, and  $Y \subseteq X$  is an interval, then the order topology and the subspace topology on  $Y$  are the same.*

# Order topology in intervals

## Lemma (Order topology in intervals)

*If  $X$  is an ordered set that has the order topology, and  $Y \subseteq X$  is an interval, then the order topology and the subspace topology on  $Y$  are the same.*

Proof.

Exercise



# Exercises

- Show that if we consider in  $\mathbb{R}$  the usual topology, then the topology on  $\mathbb{N}$  as a subspace of  $\mathbb{R}$  is discrete.

# Exercises

- Show that if we consider in  $\mathbb{R}$  the usual topology, then the topology on  $\mathbb{N}$  as a subspace of  $\mathbb{R}$  is discrete.
- Let  $Y \subseteq X$  be open. Show that  $A \subseteq Y$  is open in  $Y$  if and only if it is open in  $X$ .

# Exercises

- Show that if we consider in  $\mathbb{R}$  the usual topology, then the topology on  $\mathbb{N}$  as a subspace of  $\mathbb{R}$  is discrete.
- Let  $Y \subseteq X$  be open. Show that  $A \subseteq Y$  is open in  $Y$  if and only if it is open in  $X$ .
- Let  $A \subseteq Y \subseteq X$ , where  $X$  is a topological space. Show that the topology of  $A$  as a subspace of  $X$  is the same as  $(\tau_Y)_A$ .

# Exercises

- Show that if we consider in  $\mathbb{R}$  the usual topology, then the topology on  $\mathbb{N}$  as a subspace of  $\mathbb{R}$  is discrete.
- Let  $Y \subseteq X$  be open. Show that  $A \subseteq Y$  is open in  $Y$  if and only if it is open in  $X$ .
- Let  $A \subseteq Y \subseteq X$ , where  $X$  is a topological space. Show that the topology of  $A$  as a subspace of  $X$  is the same as  $(\tau_Y)_A$ .
- Show that if  $X$  is a topological space, and  $A \subseteq Y \subseteq X$ , then the topology of  $A$  as a subspace of  $Y$  is the same as a subspace of  $X$  ( $Y$  is a subspace of  $X$ ).



# Links

- Subspace topology - Wikipedia, the free encyclopedia