

Continuous functions


2016-02-15 9:00 -0500

Definition (Continuous functions)

Let X and Y be topological spaces. The function $f: X \rightarrow Y$ is **continuous** if for any open set V in Y , we have that $f^{-1}(V)$ is open in X .


Definition (Continuous functions)

Let X and Y be topological spaces. The function $f: X \rightarrow Y$ is **continuous** if for any open set V in Y , we have that $f^{-1}(V)$ is open in X .




Definition (Continuous functions)

Let X and Y be topological spaces. The function $f: X \rightarrow Y$ is **continuous** if for any open set V in Y , we have that $f^{-1}(V)$ is open in X .




Definition (Continuous functions)

Let X and Y be topological spaces. The function $f: X \rightarrow Y$ is **continuous** if for any open set V in Y , we have that $f^{-1}(V)$ is open in X .



Definition (Continuous functions)

Let X and Y be topological spaces. The function $f: X \rightarrow Y$ is **continuous** if for any open set V in Y , we have that $f^{-1}(V)$ is open in X .



Properties

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.
- If the topology of Y is given by a subbasis \mathcal{S} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{S}$.

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.
- If the topology of Y is given by a subbasis \mathcal{S} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{S}$.
- A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous if and only if for every $x_0 \in \mathbb{R}$ and $\epsilon > 0$, there is a $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$.

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.
- If the topology of Y is given by a subbasis \mathcal{S} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{S}$.
- A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous if and only if for every $x_0 \in \mathbb{R}$ and $\epsilon > 0$, there is a $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$.

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.
- If the topology of Y is given by a subbasis \mathcal{S} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{S}$.
- A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous if and only if for every $x_0 \in \mathbb{R}$ and $\epsilon > 0$, there is a $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$.
- Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be continuous, $x_0 \in \mathbb{R}$ and $\epsilon > 0$.

Properties

- If the topology of Y is given by a basis \mathcal{B} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{B}$.
 - If the topology of Y is given by a subbasis \mathcal{S} , then $f: X \rightarrow Y$ is continuous if and only if $f^{-1}(V)$ is open in X for every $V \in \mathcal{S}$.
 - A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous if and only if for every $x_0 \in \mathbb{R}$ and $\epsilon > 0$, there is a $\delta > 0$ such that $|x - x_0| < \delta$ implies $|f(x) - f(x_0)| < \epsilon$.
-
- Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be continuous, $x_0 \in \mathbb{R}$ and $\epsilon > 0$.
 - We have that $f^{-1}((f(x_0) - \epsilon, f(x_0) + \epsilon))$ is open in \mathbb{R} and contains x_0 . Hence there is $\delta > 0$ such that $(x_0 - \delta, x_0 + \delta) \subseteq f^{-1}(f(x_0) - \epsilon, f(x_0) + \epsilon)$.

Examples

Example (Continuous functions)

Examples

Example (Continuous functions)

- Any continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$ from calculus courses is continuous in this sense.

Examples

Example (Continuous functions)

- Any continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$ from calculus courses is continuous in this sense.
- Let \mathbb{R}_l be the set of real numbers with the Sorgenfrey topology. Then the identity function $1_{\mathbb{R}}: \mathbb{R} \rightarrow \mathbb{R}_l$ is not continuous.

Examples

Example (Continuous functions)

- Any continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$ from calculus courses is continuous in this sense.
- Let \mathbb{R}_l be the set of real numbers with the Sorgenfrey topology. Then the identity function $1_{\mathbb{R}}: \mathbb{R} \rightarrow \mathbb{R}_l$ is not continuous.
- On the other hand, the identity $1_{\mathbb{R}}: \mathbb{R}_l \rightarrow \mathbb{R}$ is continuous.

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

- *f is continuous.*

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

- *f is continuous.*
- *For every $A \subseteq X$, one has $f(\overline{A}) \subseteq \overline{f(A)}$.*

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

- *f is continuous.*
- *For every $A \subseteq X$, one has $f(\overline{A}) \subseteq \overline{f(A)}$.*
- *For every closed set C in Y , one has $f^{-1}(C)$ is closed in X .*

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

- *f is continuous.*
- *For every $A \subseteq X$, one has $f(\overline{A}) \subseteq \overline{f(A)}$.*
- *For every closed set C in Y , one has $f^{-1}(C)$ is closed in X .*
- *For each $x \in X$ and each neighborhood V of $f(x)$, there is a neighborhood U of x such that $f(U) \subseteq V$.*

Equivalences of continuity

Theorem

Let X, Y be topological spaces and $f: X \rightarrow Y$. Then the following are equivalent:

- *f is continuous.*
- *For every $A \subseteq X$, one has $f(\overline{A}) \subseteq \overline{f(A)}$.*
- *For every closed set C in Y , one has $f^{-1}(C)$ is closed in X .*
- *For each $x \in X$ and each neighborhood V of $f(x)$, there is a neighborhood U of x such that $f(U) \subseteq V$.*

When the last condition is satisfied at $x_0 \in X$, we say that f is **continuous at x_0** .

Exercises

1. Let $f: X \rightarrow Y$ continuous. If $A \subseteq X$ and x is a limit point of A , is $f(x)$ a limit point of $f(A)$?

Exercises

1. Let $f: X \rightarrow Y$ continuous. If $A \subseteq X$ and x is a limit point of A , is $f(x)$ a limit point of $f(A)$?
2. If the singleton $\{x_0\}$ is open in X , prove that every function $f: X \rightarrow Y$ is continuous at x_0 . Is the converse true?

Exercises

1. Let $f: X \rightarrow Y$ continuous. If $A \subseteq X$ and x is a limit point of A , is $f(x)$ a limit point of $f(A)$?
2. If the singleton $\{x_0\}$ is open in X , prove that every function $f: X \rightarrow Y$ is continuous at x_0 . Is the converse true?
3. Prove that $f: X \rightarrow Y$ is continuous if and only if for every $A \subseteq Y$ we have that $f^{-1}(A^\circ) \subseteq (f^{-1}(A))^\circ$.