

Higher genus theory and unramified quadratic extensions of biquadratic fields

Arithmetic Statistics, Paderborn

Robin Visser

joint work with Siu Hang Man and Pavlo Yatsyna

Faculty of Mathematics and Physics
Charles University

27 August 2026

Unit signature ranks

- Let K be a totally real number field of degree n , discriminant Δ_K , maximal order $\mathcal{O}_K$, unit group $\mathcal{O}_K^\times$, and real embeddings $\sigma_1, \dots, \sigma_n : K \rightarrow \mathbb{R}$.
- The **signature** of $\alpha \in K^\times$ is the vector $\text{sgn}(\alpha) := (\text{sgn } \sigma_1(\alpha), \dots, \text{sgn } \sigma_n(\alpha))$.
(E.g. For $K = \mathbb{Q}(\sqrt{2})$, $1 + \sqrt{2}$ has signature $(+, -)$, and $3 + 2\sqrt{2}$ has signature $(+, +)$)
- The collection of all signatures of units $\alpha \in \mathcal{O}_K^\times$ can be viewed as a subspace of $\mathbb{F}_2^n$.
The dimension of this subspace (over $\mathbb{F}_2$) is the **unit signature rank** of K .
- Since $\text{sgn}(-1) = (-, \dots, -)$, we always have $1 \leq \text{sgnrk}(K) \leq n$.

Question: How is $\text{sgnrk}(K)$ distributed amongst some family of totally real fields K ?

Real quadratic fields

Let $K = \mathbb{Q}(\sqrt{D})$ be real quadratic, $D > 0$ squarefree. Then $\text{sgnrk}(K) \in \{1, 2\}$, and

$$\text{sgnrk}(K) = 2 \iff N_{K/\mathbb{Q}}(\varepsilon_K) = -1 \iff x^2 - Dy^2 = -1 \text{ solvable} \quad (1)$$

- If D divisible by some prime p which is 3 mod 4, then (1) is not solvable mod p , and so $\text{sgnrk}(K) = 1$.
- If D is not divisible by any prime $p \equiv 3 \pmod{4}$, then $\text{sgnrk}(K) \in \{1, 2\}$. But we still find many examples in both cases:

$$\text{sgnrk}(K) = 1 : K = \mathbb{Q}(\sqrt{34}), \mathbb{Q}(\sqrt{146}), \mathbb{Q}(\sqrt{178}), \mathbb{Q}(\sqrt{194}), \mathbb{Q}(\sqrt{205}), \dots$$

$$\text{sgnrk}(K) = 2 : K = \mathbb{Q}(\sqrt{2}), \mathbb{Q}(\sqrt{5}), \mathbb{Q}(\sqrt{10}), \mathbb{Q}(\sqrt{13}), \mathbb{Q}(\sqrt{17}), \mathbb{Q}(\sqrt{26}), \dots$$

Which case occurs more often?

Real quadratic fields: a brief history

Define the sets:

$$\mathcal{D} := \{D > 0 \text{ squarefree} : p \mid D \Rightarrow p \not\equiv 3 \pmod{4}\},$$

$$\mathcal{D}^- := \{D > 0 \text{ squarefree} : \text{sgnrk } \mathbb{Q}(\sqrt{D}) = 2\}.$$

How big is $\mathcal{D}^-$ inside $\mathcal{D}$?

Let $\mathcal{D}(X) := \mathcal{D} \cap [1, X]$ and $\mathcal{D}^-(X) := \mathcal{D}^- \cap [1, X]$.

- (Dirichlet 1834) Proved that $p \in \mathcal{D}^-$ for all primes $p \equiv 1 \pmod{4}$.
- (Nagell 1932) Conjectured that the limit

$$\lim_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)}$$

exists and lies in the open interval $(0, 1)$.

Real quadratic fields: a brief history

Let $\alpha := \prod_{j \text{ odd}} (1 - 2^{-j}) \approx 0.41942\dots$

- (Stevenhagen 1993) Developed a heuristic model to conjecture that

$$\lim_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} = 1 - \alpha \approx 0.5805\dots$$

- (Fouvry–Klüners 2010) Proved the lower and upper bounds

$$\alpha \leq \liminf_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \leq \limsup_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \leq \frac{2}{3}$$

- (Fouvry–Klüners 2010) Improved their earlier lower bound to

$$\liminf_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \geq \frac{5}{4}\alpha \approx 0.52475\dots$$

Real quadratic fields: a brief history

- (Chan–Koymans–Milovic–Pagano 2019) Used the results of Smith to further improve the lower bound

$$\liminf_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \geq \alpha \cdot \sum_{n=0}^{\infty} 2^{-n(n+3)/2} \approx 0.538 \dots$$

- (Koymans–Pagano 2020) Improved the lower and upper bounds:

$$0.54302 \dots \leq \liminf_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \leq \limsup_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} \leq 0.59944 \dots$$

- (Koymans–Pagano 2022) Proved Stevenhagen's conjecture holds, i.e. the limit exists and is equal to

$$\lim_{X \rightarrow \infty} \frac{\#\mathcal{D}^-(X)}{\#\mathcal{D}(X)} = 1 - \alpha \approx 0.5805 \dots \quad \text{☺}$$

From units to class groups

To get a handle on when K has full unit signature rank, we briefly recall the ideas from the above papers:

- For a finite abelian group G , and integer $m \geq 1$, the 2^m -rank of G is

$$\mathrm{rk}_{2^m}(G) := \dim_{\mathbb{F}_2} 2^{m-1}G/2^mG$$

- Note that $\mathrm{rk}_2(G) \geq \mathrm{rk}_4(G) \geq \mathrm{rk}_8(G) \geq \dots$, and $\mathrm{rk}_{2^m}(G) = 0$ for all $m \gg 0$. The sequence $\{\mathrm{rk}_{2^m}(G)\}_{m \geq 1}$ uniquely determines the 2-primary part of G .
- For any K totally real of degree n :

$$\begin{aligned} \mathrm{sgnrk}(K) = n &\iff \mathrm{Cl}(K) \cong \mathrm{Cl}^+(K) \\ &\iff \mathrm{rk}_{2^m} \mathrm{Cl}(K) = \mathrm{rk}_{2^m} \mathrm{Cl}^+(K) \text{ for all } m \geq 1 \end{aligned}$$

since the odd parts of $\mathrm{Cl}(K)$ and $\mathrm{Cl}^+(K)$ agree.

Real quadratic case

Let $K = \mathbb{Q}(\sqrt{D})$ be real quadratic with $D \in \mathcal{D}$.

- By a theorem of Gauss, $\mathrm{rk}_2 \mathrm{Cl}(K) = \mathrm{rk}_2 \mathrm{Cl}^+(K) = \omega(D) - 1$.
- Thus, if for example either $\mathrm{rk}_4 \mathrm{Cl}^+(K) = 0$, or $\mathrm{rk}_8 \mathrm{Cl}^+(K) = 0$ and $\mathrm{rk}_4 \mathrm{Cl}^+(K) = \mathrm{rk}_4 \mathrm{Cl}(K)$, then $\mathrm{sgnrk}(K) = 2$.

To therefore obtain bounds for the ratio $\#\mathcal{D}^-(X)/\#\mathcal{D}(X)$, we must count fields K with prescribed $\mathrm{rk}_4 \mathrm{Cl}^+(K)$, $\mathrm{rk}_8 \mathrm{Cl}^+(K)$, etc.

Rédei matrices

Let $D \in \mathcal{D}$ with prime divisors $p_1, p_2, \dots, p_t$. The **Rédei matrix** $R(D)$ is a $t \times t$ matrix over $\mathbb{F}_2$ where

1. For $i \neq j$, we define $R(D)_{ij} := r_{ij}$ where $\left(\frac{p_i}{p_j}\right) = (-1)^{r_{ij}}$.
2. The diagonal entries $R(D)_{ii}$ are chosen such that $\sum_{j=1}^t R(D)_{ij} = 0 \in \mathbb{F}_2$ for every i .

Note $R(D)$ is a symmetric matrix (by quadratic reciprocity).

Theorem (Rédei 1934)

Let $K = \mathbb{Q}(\sqrt{D})$ with $D \in \mathcal{D}$. Then

$$\mathrm{rk}_4 \mathrm{Cl}^+(K) = \omega(D) - 1 - \mathrm{rank}_{\mathbb{F}_2} R(D) \quad (= \mathrm{corank}_{\mathbb{F}_2} R(D) - 1)$$

Rédei matrices: an example

Example: Let $K = \mathbb{Q}(\sqrt{D})$ where $D = 5 \cdot 13 \cdot 37 = 2405$. The Legendre symbols are

$$\left(\frac{5}{13}\right) = \left(\frac{5}{37}\right) = \left(\frac{13}{37}\right) = -1,$$

so all off-diagonal entries equal 1, and all diagonal entries equal 0:

$$R(2405) = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

We compute $\text{rank } R(D) = 2$ and $\text{corank } R(D) = 1$. Hence

$$\text{rk}_2 \text{Cl}^+ = 2, \quad \text{rk}_4 \text{Cl}^+ = 0, \quad \text{Cl}^+(K) \cong C_2 \times C_2,$$

and in particular $x^2 - 2405y^2 = -1$ is soluble.

Counting Rédei Matrices

Let $\mathcal{R}_t$ be the set of all Rédei matrices of size $t \times t$. We have a rank-preserving bijection from $\mathcal{R}_t$ to the set of symmetric $(t-1) \times (t-1)$ matrices over $\mathbb{F}_2$

Theorem (MacWilliams 1969)

Let $N(t, r)$ be the number of symmetric $t \times t$ matrices over $\mathbb{F}_2$ of rank r . Then:

$$N(t, 2s) = \prod_{i=1}^s \frac{2^{2i}}{2^{2i} - 1} \cdot \prod_{i=0}^{2s-1} (2^{t-i} - 1), \quad N(t, 2s+1) = \prod_{i=1}^s \frac{2^{2i}}{2^{2i} - 1} \cdot \prod_{i=0}^{2s} (2^{t-i} - 1).$$

Corollary

Fix some non-negative integer $m \geq 0$. Then

$$\lim_{t \rightarrow \infty} \frac{\#\{M \in \mathcal{R}_t \mid \text{corank}_{\mathbb{F}_2} M = m+1\}}{\#\mathcal{R}_t} = \alpha \prod_{i=1}^m (2^i - 1)^{-1}$$

Biquadratic fields

Let $K = \mathbb{Q}(\sqrt{d_1}, \sqrt{d_2})$ be a real biquadratic field, with quadratic subfields $K_i = \mathbb{Q}(\sqrt{d_i})$, where $d_3 = d_1 d_2 / \gcd(d_1, d_2)^2$. Here $n = 4$.

- By a theorem of Edgar–Mollin–Peterson (1986), we have

$$\begin{aligned} \text{sgnrk}(K) = 4 &\implies \text{sgnrk}(K_i) = 2 && \text{for all } i = 1, 2, 3 \\ &\implies x^2 - D_i y^2 = -1 \text{ solvable} && \text{for all } i = 1, 2, 3. \end{aligned}$$

- (Kubota 1956) If $\text{sgnrk}(K_i) = 2$ for each $i = 1, 2, 3$, then $\text{sgnrk}(K) \in \{3, 4\}$.

We again find many examples in both cases:

$$\text{sgnrk}(K) = 3 : K = \mathbb{Q}(\sqrt{2}, \sqrt{85}), \mathbb{Q}(\sqrt{10}, \sqrt{17}), \mathbb{Q}(\sqrt{13}, \sqrt{85}), \mathbb{Q}(\sqrt{17}, \sqrt{65}), \dots$$

$$\text{sgnrk}(K) = 4 : K = \mathbb{Q}(\sqrt{2}, \sqrt{5}), \mathbb{Q}(\sqrt{5}, \sqrt{13}), \mathbb{Q}(\sqrt{5}, \sqrt{17}), \mathbb{Q}(\sqrt{2}, \sqrt{13}), \dots$$

Which case occurs more often?

Rédei matrices for biquadratic fields

Write $\Delta_K = D^2$ with $D \in \mathcal{D}$, and let $p_1, \dots, p_t$ be the primes dividing D . They partition into three blocks

$$\gamma_1 = \gcd(d_1, d_3), \quad \gamma_2 = \gcd(d_2, d_3), \quad \gamma_3 = \gcd(d_1, d_2),$$

of sizes t_1, t_2, t_3 (so $d_1 = \gamma_1\gamma_3$, $d_2 = \gamma_2\gamma_3$, $d_3 = \gamma_1\gamma_2$). Let $\underline{R}(D)$ be the Rédei *prematrix* of D consisting of Jacobi symbols (p_i/p_j) . We break $\underline{R}(D)$ into blocks,

$$\underline{R}(D) = \begin{pmatrix} \underline{A}_{11} & A_{12} & A_{13} \\ A_{21} & \underline{A}_{22} & A_{23} \\ A_{31} & A_{32} & \underline{A}_{33} \end{pmatrix},$$

where the Rédei prematrices of D_i are given by

$$\underline{R}(D_1) = \begin{pmatrix} \underline{A}_{11} & A_{13} \\ A_{31} & \underline{A}_{33} \end{pmatrix}, \quad \underline{R}(D_2) = \begin{pmatrix} \underline{A}_{22} & A_{23} \\ A_{32} & \underline{A}_{33} \end{pmatrix}, \quad \underline{R}(D_3) = \begin{pmatrix} \underline{A}_{11} & A_{12} \\ A_{21} & \underline{A}_{22} \end{pmatrix}.$$

The corresponding Rédei matrices $R(D_i)$ are obtained by filling in the diagonals of $\underline{R}(D_i)$.

Rédei matrices for biquadratic fields

Theorem (Man-V-Yatsyna)

Let $m_1, m_2, m_3 \in \mathbb{N}_0$ be fixed integers. Then, as $t \rightarrow \infty$ with $t_1 \geq t_2 \geq t_3 \geq \frac{t}{10}$, we have

$$\lim_{t \rightarrow \infty} \frac{\#\{A \in \mathcal{R}_t \mid \text{corank}_{\mathbb{F}_2} A_{[i]} = m_i + 1, i = 1, 2, 3\}}{\#\mathcal{R}_t} = \alpha^3 \prod_{i=1}^3 \prod_{j=1}^{m_i} (2^j - 1)^{-1},$$

Construction of $A_{[i]}$:

1. Break down the prematrix $\underline{A}$ associated to A into blocks of size t_1, t_2, t_3 respectively
2. Take $A_{[i]}$ to be Rédei matrices:

$$\underline{A} = \begin{pmatrix} \underline{A}_{11} & A_{12} & A_{13} \\ A_{21} & \underline{A}_{22} & A_{23} \\ A_{31} & A_{32} & \underline{A}_{33} \end{pmatrix},$$

$$A_{[1]} = R \begin{pmatrix} \underline{A}_{11} & A_{13} \\ A_{31} & \underline{A}_{33} \end{pmatrix}, \quad A_{[2]} = R \begin{pmatrix} \underline{A}_{22} & A_{23} \\ A_{32} & \underline{A}_{33} \end{pmatrix},$$
$$A_{[3]} = R \begin{pmatrix} \underline{A}_{11} & A_{12} \\ A_{21} & \underline{A}_{22} \end{pmatrix}.$$

Proof strategy


Sketch proof:

1. **Reverse the construction:** First fill in the diagonal entries of the three Rédei prematrices, independently of each other, and then fill in the non-diagonal blocks
2. **Grow one row and column at a time.** Add a row and column of an off-diagonal block one at a time, keeping track of the rank. Show that from a symmetric matrix corank m , ones moves to corank

$$m - 1 \text{ w.p. } 1 - 2^{-m}, \quad m \text{ w.p. } 2^{-m-1}, \quad m + 1 \text{ w.p. } 2^{-m-1}.$$

3. **Model this process as a Markov chain**, i.e. the corank performs a random walk on the sequence of nonnegative integers.
4. **Independence:** Do the above to track the coranks of each of $A_{[1]}$, $A_{[2]}$, and $A_{[3]}$, and show these distributions are independent.

Proof strategy



The probability of landing on corank m approaches $\alpha \prod_{j=1}^m (2^j - 1)$.

Back to fields

Let $\mathcal{K}(X) := \{K \text{ real biquadratic} \mid \Delta_K = D^2, D \in \mathcal{D}(X)\}$

Theorem (Man-V-Yatsyna)

$$\liminf_{X \rightarrow \infty} \frac{\#\{K \in \mathcal{K}(X) \mid N_{K_i/\mathbb{Q}}(\varepsilon_i) = -1, \forall 1 \leq i \leq 3\}}{\#\mathcal{K}(X)} \geq \alpha^3 \approx 0.0738 \dots$$

- This is only a lower bound, as we have contented ourselves with the case $\text{rk}_4 \text{Cl}^+(K) = 0$ (though the method extends to positive rk_4 , as in Fouvry-Klüners and CKMP)

Sketch proof

We apply the toolbox from Chan–Koymans–Milovic–Pagano (2022):

1. By Erdős–Kac, almost all $D \in \mathcal{D}(X)$ satisfy $|\omega(D) - \mu| \leq \mu^{2/3}$, with $\mu := \frac{1}{2} \log \log X$. The exceptions have relative size $O(\exp(-\frac{1}{3}\mu^{1/3}))$. So it suffices to work with the subsets

$$\mathcal{D}_t(X) := \{D \in \mathcal{D}(X) \mid \omega(D) = t\}, \quad |t - \mu| \leq \mu^{2/3}$$

2. For each t , we can find some “acceptable” boxes $W_i \subseteq \mathcal{D}_t(X)$ such that
 - (i) almost all $D \in \mathcal{D}_t(X)$ is contained in some W_i ; the exceptions have relative size $\ll (\log \log X)^{-c}$ for some small $c > 0$.
 - (ii) within each W_i , the Rédei matrices are equidistributed, up to some permutation of rows and columns, with relative error $\ll (\log \log X)^{-c}$ for some small $c > 0$.
3. For fixed $D \in \mathcal{D}_t(X)$, almost all biquadratic K with $\Delta_K = D^2$ satisfy $t_1, t_2, t_3 \geq \frac{t}{10}$. The exceptions have relative size $\ll e^{-c \log \log X}$ for some small $c > 0$.

Kuroda's formula

Theorem (Kuroda's formula)

Let K be real biquadratic with quadratic subfields K_1, K_2, K_3 . Then

$$h(K) = \frac{Q}{4} \cdot h(K_1)h(K_2)h(K_3), \quad \text{where } Q := [\mathcal{O}_K^\times : \mathcal{O}_{K_1}^\times \mathcal{O}_{K_2}^\times \mathcal{O}_{K_3}^\times].$$

Proposition

Suppose $h^+(K_i) = h(K_i)$ for $i = 1, 2, 3$. Then

$$h^+(K) = \frac{1}{2} h(K_1)h(K_2)h(K_3),$$

and $h^+(K) = h(K) \iff Q = 2 \iff \text{sgnrk}(K) = 4$.

$$\text{sgnrk}(K) = 4 \iff h^+(K) = h(K) \iff H_2^+(K) \text{ totally real.}$$

From class numbers to fields

To obtain bounds on the number of fields with $\text{sgnrk}(K) = 4$, we make the following simplifying assumption:

$$\text{rk}_4 \text{Cl}^+(K_1) = \text{rk}_4 \text{Cl}^+(K_2) = \text{rk}_4 \text{Cl}^+(K_3) = 0. \quad (*)$$

Consequences:

- All three negative Pell equations are solvable (Gauss).
- We have $\text{rk}_8 \text{Cl}^+(K) = 0$, (Chan–Milovic 2022)
- We have the order: $h_2(K)^+ = \frac{1}{2}h_2(K_1)h_2(K_2)h_2(K_3)$.

To determine if $\text{sgnrk}(K) = 4$, we need to know if $h_2(K) = h_2^+(K)$, i.e. if the two narrow class field $H_2^+(K)$ is *totally real*. We do this by classifying quadratic extensions of K which are unramified at all finite places (UAFP) and determining which are real.

Quadratic UAFP extensions

Recall $\text{rk}_2 \text{Cl}^+(K) = \#\{\text{independent UAFP quadratic extensions of } K\}$.

Two known sources of quadratic UAFP extensions:

- **Genus theory:** For each prime $p \mid \Delta_K$, the extension $K(\sqrt{p})/K$ is UAFP.
- **Fouvry–Klüners (2010):** Decompositions of the second type $D = D_1 D_2$: solve $x^2 - D_1 y^2 - D_2 z^2 = 0$ and adjoin $\sqrt{x + y\sqrt{D_1}}$.

But there are more!

Example

For $K = \mathbb{Q}(\sqrt{130}, \sqrt{493})$, the extension $K(\sqrt{\beta})/K$ is UAFP, where

$$\beta = 77 + 4\sqrt{130} + \sqrt{493}.$$

Can we count how many there are, and can we explicitly classify them ?

Higher genus theory

To do this, we use the **higher genus theory** of Koymans and Pagano.

Let $K = \mathbb{Q}(\sqrt{a_1}, \dots, \sqrt{a_m})$ be a multiquadratic field of degree 2^m , where $a_1, \dots, a_m$ are pairwise coprime, squarefree, and have all prime factors $\equiv 1 \pmod{4}$. Let H be the maximal elementary exponent 2 extension of K unramified at all finite places. We have $\text{Gal}(H/K) \cong \text{Cl}^+(K)[2]$.

Koymans–Pagano (2022) study the group $G = \text{Gal}(H/\mathbb{Q})$ as an *expansion group*. The descending central series of G can be equipped with a graded Lie algebra structure. They construct a universal expansion group $\mathcal{G}$, of which any such G is a quotient.

The theory gives us a way to handle $\text{Cl}^+(K)[2]$. Unramified $\mathbb{F}_2$ -extensions of K are built from 2-cocycles assembled out of data on the multiquadratic subfields of index 2 (the “corners” of K), and whether such a cocycle is trivial in $H^2(G_{\mathbb{Q}}, \mathbb{F}_2)$ is a condition on local Hilbert symbols.

Classifying quadratic UAFP extensions

Let $\gamma_1, \gamma_2, \gamma_3 \in \mathcal{D}$ be pairwise coprime, and let $d_1 = \gamma_1\gamma_3$, $d_2 = \gamma_2\gamma_3$, $d_3 = \gamma_1\gamma_2$. Let $K = \mathbb{Q}(\sqrt{d_1}, \sqrt{d_2})$. From higher genus theory, the unramified non-genus $\mathbb{F}_2$ -extensions of K are characterised by 2-cocycles $\theta : G_{\mathbb{Q}} \rightarrow \mathbb{F}_2$ of the form

$$\theta = \chi_{d_1} \cup \chi_{b_2 b_3} + \chi_{d_2} \cup \chi_{b_1 b_3}, \quad b_i \mid \gamma_i,$$

such that θ is trivial in $H^2(G_{\mathbb{Q}}, \mathbb{F}_2)$, i.e. θ is a *coboundary*. Galois cohomology turns this into a condition on local Hilbert symbols:

$$\begin{aligned} \theta \text{ trivial in } H^2(G_{\mathbb{Q}}, \mathbb{F}_2) &\iff (d_1, b_2 b_3)_v + (d_2, b_1 b_3)_v = 0 \quad \forall v \\ &\iff (d_1, b_2)_v + (d_2, b_1)_v + (d_3, b_3)_v = 0 \quad \forall v. \end{aligned}$$

- The Fouvry–Klüners extensions are exactly the *pure* cocycles, where only one of b_1, b_2, b_3 is nontrivial (hence why these miss some UAFP extensions).

Counting UAFP extensions

Define the auxiliary Rédei matrix R' as the symmetric $\mathbb{F}_2$ -matrix with entries:

$$w_{p,q} = \begin{cases} \ell, & q \mid \gamma_j, j \neq i, \left(\frac{q}{p}\right) = (-1)^\ell, \\ 0, & q \mid \gamma_i, q \neq p, \\ \tau, & q = p, \left(\frac{\prod_{j \neq i} \gamma_j}{p}\right) = (-1)^\tau. \end{cases}$$

i.e. R' is the Rédei matrix of D with the intra-block off-diagonal entries deleted and diagonals recomputed such that rows sum to 0.

Theorem

The number of independent non-genus UAFP C_2 -extensions of K is $\text{corank } R' - 1$, and

$$\text{rk}_2 \text{Cl}^+(K) = \omega(K) + \text{corank}(R') - 3.$$

Example

Let $d_1 = p_1 p_2$, $d_2 = p_3 p_4$. Under our assumption $\text{rk}_4 \text{Cl}^+(K_i) = 0$, up to permuting the primes, the Rédei prematrix $A_K = \left(\left(\frac{p_i}{p_j} \right) \right)_{ij}$ is either

$$\begin{pmatrix} * & 1 & 1 & 0 \\ 1 & * & 0 & 0 \\ 1 & 0 & * & 1 \\ 0 & 0 & 1 & * \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} * & 1 & 1 & 1 \\ 1 & * & 0 & 0 \\ 1 & 0 & * & 1 \\ 1 & 0 & 1 & * \end{pmatrix}.$$

Deleting the intra-block entries ($p_1 p_2$ and $p_3 p_4$) and refilling the diagonals gives

$$R' = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \text{ corank} = 3, \quad \text{resp.} \quad R' = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \text{ corank} = 2.$$

With $\omega(K) = 4$ the theorem gives $\text{Cl}_2^+(K) \cong C_2^4$, resp. $C_4 \times C_2^2$.

But this is not yet enough

Two fields with the same Legendre data

Take $\gamma_1 = 5$, $\gamma_2 = 13$ and $\gamma_3 = 37$ or 97 . In both cases all three Legendre symbols equal -1 , so A_K is the same, and indeed $\text{Cl}_2^+(K) \cong C_4$ in both cases. But:

K	$\text{Cl}(K)$	$\text{Cl}^+(K)$	$\text{sgnrk}(K)$
$\mathbb{Q}(\sqrt{185}, \sqrt{481})$	C_4	C_4	4
$\mathbb{Q}(\sqrt{485}, \sqrt{1261})$	C_2	C_4	3

- $h^+(K)$ is determined by from our assumption $\text{rk}_4 \text{Cl}^+(K_i) = 0$ and the Legendre symbols, but $h(K)$ is not!
- We need to determine whether the narrow 2-Hilbert class field $H_2^+(K)$ is totally real.
- Instead of only Legendre symbols, we must use **Rédei symbols**.

Detecting real UAFP extensions

Let $\theta = \chi_{d_1} \cup \chi_{b_2 b_3} + \chi_{d_2} \cup \chi_{b_1 b_3}$ be a 2-cocycle as above, and let L_θ/K be the corresponding UAFP extension. From higher genus theory, we have that $\theta = d\psi$ is the coboundary of a 1-cochain, whose field of definition $L(\psi)$ is an unramified $\mathbb{F}_2$ -extension of the multiquadratic field $M_\theta = \mathbb{Q}(\sqrt{d_1}, \sqrt{d_2}, \sqrt{b_1 b_3}, \sqrt{b_2 b_3})$.

We obtain the condition

$$L_\theta \text{ totally real} \iff \text{Inv}_\infty(\text{res}_{G_\infty} \psi \cup \chi_{-1}) = \psi(\text{Frob}_\infty) = 0.$$

In order to make this more tractable, we use *Rédei symbols*.

- For Fourvy–Klüner extensions, coming from a decomposition of second type $D = D_1 D_2$, realness is determined by the single Rédei symbol $[D_1, D_2, -1]$.
- For non-genus non-FK extensions, this unfortunately isn't determined by a single Rédei symbol. We must in general take a sum of Rédei symbols.

The criterion

Theorem (Man–V–Yatsyna)

Let R be the Rédei matrix of $\mathbb{Q}(\sqrt{\gamma_1\gamma_2\gamma_3})$ and R' the auxiliary matrix above, and let $t_k = \omega(\gamma_k)$. Let $\emptyset \subsetneq B \subsetneq [t]$ be a subset such that $\sum_{i \in B} e_i \in \ker R'$, and let $b_i = \gcd(\prod_{j \in B} p_j, \gamma_i)$.

Then $\theta = \chi_{d_1} \cup \chi_{b_2 b_3} + \chi_{d_2} \cup \chi_{b_1 b_3}$ is trivial in $H^2(G_{\mathbb{Q}}, \mathbb{F}_2)$, and there exist subsets $C_i, D_i \subseteq [t]$ depending only on t_1, t_2, t_3, R and B (and not on the primes) such that, writing $u_i = \prod_{j \in C_i} p_j$ and $v_i = \prod_{j \in D_i} p_j$, the field L_θ is totally real if and only if

$$\sum_i [u_i, v_i, -1] = 0 \in \mathbb{F}_2.$$

What we expect to prove

Theorem? (Man–V–Yatsyna 2026+)

A positive density of real biquadratic fields, ordered by discriminant, whose quadratic subfields all have unit signature rank 2, have unit signature rank 3.

- Chan–Milovic (2022) computed numerically that the quadratic extensions attached to the non-genus part of the 2-rank, and those attached to the 4-rank, are totally real about half the time (and independently).

Conjecture

100% of real biquadratic fields, ordered by discriminant, whose quadratic subfields all have unit signature rank 2, have unit signature rank 3.

What remains

- Can we model the distribution of the corank(R') for the auxiliary Rédei matrix R' , i.e. the distribution of $\text{rk}_2 \text{Cl}^+(K)$? (this is almost done!)
- Can we remove the simplifying hypothesis that $\text{rk}_4 \text{Cl}^+(K_i) = 0$ for every $i = 1, 2, 3$, i.e. control $\text{rk}_8 \text{Cl}^+(K)$ and characterise UAFP extensions in this more general setting.
- Can we count and classify the cyclic quartic UAFP extensions? (this will be needed to attack the strong conjecture).

Thank you!