

Solvability of Equations in Elementary Functions

Rodion Zaytsev

Moscow Institute of Physics and Technology, Department of Innovation and High Technology

Introduction

The most general problem of solvability can be formulated as follows: we have a set of "known quantities" K and a set of "allowed" operations \mathcal{F} . The question is, can a given element be obtained from K by repeatedly applying operations from \mathcal{F} ? The main approach to these questions is Galois theory (algebraic, differential and topological), which connects the problem of solvability to the properties of an associated group and a method developed by Liouville, where the idea is that either the solution is "simple" or doesn't exist [1].

Historical background

In 1770 Lagrange unified various methods for solving polynomial equations in radicals [2]. The method relied on Lagrange resolvents, which allowed to reduce an equation to one of a lower degree. This method, however, does not work for higher degrees, as the new equation has a bigger degree than the original polynomial. Lagrange's work inspired Ruffini, who made 6 publications in an attempt to prove insolubility of equations of degree higher than 4, the first publication made in 1799 and the last in 1813 [3]. The first accepted proof is due to Abel [4], published in 1824. Later in that decade, Galois developed a general theory which could determine whether a given equation is solvable by radicals [5]. Galois method connects the solvability of an equation with the solvability of its Galois group. In 1833 Liouville extended the works of Abel in 1823 about expression of integrals in elementary functions [6] (Abel's work "a general representation of the possibility to integrate all differential formulas" was not published). Around 1883-1904 Picard and Vessiot associated a Galois group with differential equation, to describe when it can be solved in quadratures and this theory was further refined by Kolchin around 1950 [7]. In 1963 V.I. Arnold discovered a topological proof of the Abel-Ruffini theorem [8], which was the starting point of topological Galois theory, to which this work is devoted.

Notions from topological Galois theory

- "Known quantities" $K = \mathbb{C}(a)$, "allowed" operations $\mathcal{F} = \{\text{algebraic operations and elementary functions}\}$
- **Def.** *Monodromy group is the group acting on the roots of the equation when the parameter a makes a loop.*
- Monodromy group acting on an elementary function is cyclic.
- When the equation is solvable the monodromy group must be solvable.

My contribution

$$\tan x - x = a$$

- Calculated singular points
- Proved a lemma about the location of roots of $\tan(x) - x = 0$
- Calculated the action of the group for paths around singular points
- Proved that the monodromy group contained the alternating group

$$x^x = a$$

- Transforms to equation $x + e^x = a$
- Singular points were calculated
- Action of paths around these point was calculated
- Using transitivity it was shown that the group of monodromy is the symmetric group, hence unsolvable

Main result 1

The equation $\tan x - x = a$ is not solvable in elementary functions [9]

Main result 2

The equation $x^x = a$ is not solvable in elementary functions




Figure 1: $\tan x - x = a$ monodromy




Figure 2: $x^x = a$ monodromy

References

- [1] AG Khovansky. Topological galois theory. *Solvability and non-solvability of equations in finite terms (in Russian)*, MCCME, Moscow, 2008.
- [2] Joseph-Louis Lagrange. Reflexions sur la resolution algebrique des equations, nouveaux memoires de l'acade. *Royale des sciences et belles-lettres, avec l'histire pour la meme annee*, 1:134–215, 1770.
- [3] Paolo Ruffini. Teoria generale delle equazioni, in cui si dimostra impossibile la soluzione algebrica delle equazioni generali di grado superiore al quarto. *Stamperia di S. Tommaso d'Aquino*, 1799.
- [4] Niels Henrik Abel. *œuvres complètes de Niels Henrik Abel*, volume 1. Grøndahl & søn, 1881.
- [5] E Galois. Sur les conditions de résolubilité des équations par radicaux. *Journal de mathématiques pures et appliquées*, 11:417–444, 1846.
- [6] Joseph Liouville. *Premier mémoire sur la détermination des intégrales dont la valeur est algébrique*. Impr. Royale, 1833.
- [7] Ellis Robert Kolchin. The picard-vessiot theory of homogeneous linear ordinary differential equations. *Proceedings of the National Academy of Sciences*, 32(12):308–311, 1946.
- [8] Henryk Żołądek. The topological proof of abel-ruffini theorem. *Topological Methods in Nonlinear Analysis*, 16(2):253–265, 2000.
- [9] Alexey Belov-Kanel, Alexei Malistov, and Rodion Zaytsev.

Solvability of equations in elementary functions. *Journal of Knot Theory and Its Ramifications*, 29(02):2040005, feb 2020.