

# Magnetic Flux

## 1 Introduction

---

The definition of magnetic flux is

$$\Phi_B = \int \mathbf{B} \cdot d\mathbf{A}$$

When the magnitude and direction of  $\vec{\mathbf{B}}$  is the same at all points on the surface, the integral simplifies to

$$\Phi_B = \mathbf{B} \cdot \mathbf{A}$$

or, equivalently,

$$\Phi_B = BA \cos \phi$$


where  $\theta$  is the angle between the  $\vec{\mathbf{B}}$  and  $\vec{\mathbf{A}}$  vectors.

Note that previously electric flux was covered. The same techniques that apply to computing electric flux apply to magnetic flux. See the Electric Flux activity for additional discussion.

## 2 Computing $\Phi_B$

---

In the following figure, a square loop of area  $A$  that can be rotated about the  $z$  axis is shown. Assume that the normal direction of the loop is as shown in the diagram. A uniform external magnetic field points in the  $\hat{j}$  direction.




1. At what angles in the range of  $\phi = [0, 360^\circ]$  is the magnetic flux zero?
2. Draw the loop,  $\hat{\mathbf{n}}$ , and  $\vec{\mathbf{B}}_{\text{ext}}$  as they would appear when viewed from large  $z$  when  $\phi = 0^\circ$ ,  $\phi = 45^\circ$ ,  $\phi = 135^\circ$ , and  $\phi = 180^\circ$ .
3. When  $\phi = 45^\circ$ , is the magnetic flux positive or negative? What is its value in terms of  $B_o A$ ?
4. When  $\phi = 135^\circ$ , is the magnetic flux positive or negative? What is its value in terms of  $B_o A$ ?
5. Sketch a plot of the magnetic flux,  $\Phi$ , as a function of  $\phi$ .

### 3 Computing $d\Phi_B/dt$

---

The time rate of change of magnetic flux through a closed loop is a quantity that will be used when Faraday's law is covered.

Suppose the loop in the previous problem rotates at a constant rate.



1. At what angles is  $d\Phi_B/dt = 0$ ?
2. For what angle range is  $d\Phi_B/dt$  increasing?
3. For what angle range is  $d\Phi_B/dt$  decreasing?
4. What is the formula for  $d\Phi_B/dt$ ?