

Probability: The Foundation of Uncertainty


PSC4375: Week 9

Prof. Weldzius


Villanova University

Slides Updated: 2025-03-19


Learning about populations




Learning about populations



Learning about populations




Learning about populations



Probability: formalize the uncertainty about how our data came to be

Learning about populations



Probability: formalize the uncertainty about how our data came to be

Inference: learning about the population from a sample of data

Why probability?

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?
 - Weather changes \rightsquigarrow slightly different estimated effect.

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?
 - Weather changes \rightsquigarrow slightly different estimated effect.
- Statistical inference is a **thought experiment** about uncertainty.

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?
 - Weather changes \rightsquigarrow slightly different estimated effect.
- Statistical inference is a **thought experiment** about uncertainty.
 - Imagine a world where the treatment effect were 0 in the population.

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?
 - Weather changes \rightsquigarrow slightly different estimated effect.
- Statistical inference is a **thought experiment** about uncertainty.
 - Imagine a world where the treatment effect were 0 in the population.
 - What types of estimated effects would we see in this world by chance?

Why probability?

- Probability quantifies chance variation or uncertainty in outcomes.
 - It might rain or be sunny today, we don't know which.
- We estimated a treatment effect of 7.2, but what if we reran history?
 - Weather changes \rightsquigarrow slightly different estimated effect.
- Statistical inference is a **thought experiment** about uncertainty.
 - Imagine a world where the treatment effect were 0 in the population.
 - What types of estimated effects would we see in this world by chance?
- Probability to the rescue!

Sample spaces & events

Sample spaces & events

- To formalize chance, we need to define the set of possible outcomes.

Sample spaces & events

- To formalize chance, we need to define the set of possible outcomes.
- **Sample space:** Ω the set of possible outcomes.

Sample spaces & events

- To formalize chance, we need to define the set of possible outcomes.
- **Sample space:** Ω the set of possible outcomes.
- **Event:** any subset of outcomes in the sample space

Example: gambling





Example: gambling

- A standard deck of playing cards has 52 cards:





Example: gambling

- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)





Example: gambling

- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)
 - in each of 4 suits: (, , , )






Example: gambling

- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)
 - in each of 4 suits: (, , , )
 - Hypothetical trial: pick a card, any card.






Example: gambling


























- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)
 - in each of 4 suits: (, , , )
 - Hypothetical trial: pick a card, any card.
 - Uncertainty: we don't know which card we're going to get.





Example: gambling

- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)
 - in each of 4 suits: (, , , )
 - Hypothetical trial: pick a card, any card.
 - Uncertainty: we don't know which card we're going to get.
 - One possible outcome: picking a 4

Example: gambling

- A standard deck of playing cards has 52 cards:
 - 13 rank cards: (2,3,4,5,6,7,8,9,10,J,Q,K,A)
 - in each of 4 suits: (, , , )
 - Hypothetical trial: pick a card, any card.
 - Uncertainty: we don't know which card we're going to get.
 - One possible outcome: picking a 4
 - Sample space:

2 	3 	4 	5 	6 	7 	8 	9 	10 	J 	Q 	K 	A 
2 	3 	4 	5 	6 	7 	8 	9 	10 	J 	Q 	K 	A 
2 	3 	4 	5 	6 	7 	8 	9 	10 	J 	Q 	K 	A 
2 	3 	4 	5 	6 	7 	8 	9 	10 	J 	Q 	K 	A 

- An event: picking a Queen, $\{Q, Q, Q, Q\}$

What is probability?

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing $4\clubsuit$:

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing $4\clubsuit$: $\frac{1}{52}$

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣:

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣: $\frac{13}{52}$

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣: $\frac{13}{52}$
- Same math, but different interpretations:

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣: $\frac{13}{52}$
- Same math, but different interpretations:
 - **Frequentist**: $\mathbb{P}()$ reflects relative frequency in a large number of trials.

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣: $\frac{13}{52}$
- Same math, but different interpretations:
 - **Frequentist:** $\mathbb{P}()$ reflects relative frequency in a large number of trials.
 - **Bayesian:** $\mathbb{P}()$ are subjective beliefs about outcomes.

What is probability?

- The probability $\mathbb{P}(A)$ represents how likely event A occurs.
- If **all outcomes equally likely**, then:

$$\mathbb{P}(A) = \frac{\text{number of elements in } A}{\text{number of elements in } \Omega}$$

- Example: randomly draw 1 card:
 - probability of drawing 4♣: $\frac{1}{52}$
 - probability of drawing any ♣: $\frac{13}{52}$
- Same math, but different interpretations:
 - **Frequentist**: $\mathbb{P}()$ reflects relative frequency in a large number of trials.
 - **Bayesian**: $\mathbb{P}()$ are subjective beliefs about outcomes.
- Not our fight \rightsquigarrow stick to frequentism in this class

Probability axioms

- Probability quantifies how likely or unlikely events are.

Probability axioms

- Probability quantifies how likely or unlikely events are.
- We'll define the probability $\mathbb{P}(A)$ with three axioms:

Probability axioms

- Probability quantifies how likely or unlikely events are.
 - We'll define the probability $\mathbb{P}(A)$ with three axioms:
- 1 (Nonnegativity) $\mathbb{P}(A) \geq 0$ for every event A

Probability axioms

- Probability quantifies how likely or unlikely events are.
 - We'll define the probability $\mathbb{P}(A)$ with three axioms:
- 1 (Nonnegativity) $\mathbb{P}(A) \geq 0$ for every event A
 - 2 (Normalization) $\mathbb{P}(\Omega) = 1$

Probability axioms

- Probability quantifies how likely or unlikely events are.
 - We'll define the probability $\mathbb{P}(A)$ with three axioms:
- 1 (Nonnegativity) $\mathbb{P}(A) \geq 0$ for every event A
 - 2 (Normalization) $\mathbb{P}(\Omega) = 1$
 - 3 (Addition Rule) If two events A and B are mutually exclusive

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

Gambling 102

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{\text{Q}\clubsuit\}$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{\text{Q}\clubsuit \text{ or } \text{Q}\spadesuit\}$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{\text{Q}\clubsuit \text{ or } \text{Q}\spadesuit \text{ or } \text{Q}\heartsuit\}$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{Q\clubsuit \text{ or } Q\spadesuit \text{ or } Q\heartsuit \text{ or } Q\diamondsuit\}$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{Q\clubsuit \text{ or } Q\spadesuit \text{ or } Q\heartsuit \text{ or } Q\diamondsuit\}$
- Union of mutually exclusive events \rightsquigarrow use addition rule

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{\text{Q}\clubsuit \text{ or } \text{Q}\spadesuit \text{ or } \text{Q}\heartsuit \text{ or } \text{Q}\diamondsuit\}$
- Union of mutually exclusive events \rightsquigarrow use addition rule
 - $\rightsquigarrow \mathbb{P}(\text{Q card}) = \mathbb{P}(\text{Q}\clubsuit) + \mathbb{P}(\text{Q}\spadesuit) + \mathbb{P}(\text{Q}\heartsuit) + \mathbb{P}(\text{Q}\diamondsuit)$

Gambling 102

- What is $\mathbb{P}(\text{Q card})$ if a single card is randomly selected from a deck?
 - “randomly selected” \rightsquigarrow all cards have prob. $1/52$
- “4 card” event = $\{\text{Q}\clubsuit \text{ or } \text{Q}\spadesuit \text{ or } \text{Q}\heartsuit \text{ or } \text{Q}\diamondsuit\}$
- Union of mutually exclusive events \rightsquigarrow use addition rule
 - $\rightsquigarrow \mathbb{P}(\text{Q card}) = \mathbb{P}(\text{Q}\clubsuit) + \mathbb{P}(\text{Q}\spadesuit) + \mathbb{P}(\text{Q}\heartsuit) + \mathbb{P}(\text{Q}\diamondsuit) = \frac{4}{52}$

Useful probability facts

- Probability of the complement:

Useful probability facts

- Probability of the complement: $\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A)$

Useful probability facts

- Probability of the complement: $\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A)$
 - Probability of **not** drawing a Queen is $1 - \frac{4}{52} = \frac{48}{52}$

Useful probability facts

- Probability of the complement: $\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A)$
 - Probability of **not** drawing a Queen is $1 - \frac{4}{52} = \frac{48}{52}$
- **General addition rule** for any events, A and B :

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \text{ and } B)$$

Useful probability facts

- Probability of the complement: $\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A)$
 - Probability of **not** drawing a Queen is $1 - \frac{4}{52} = \frac{48}{52}$
- **General addition rule** for any events, A and B :

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \text{ and } B)$$

- Probability of drawing Queen or ♠

Useful probability facts

- Probability of the complement: $\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A)$
 - Probability of **not** drawing a Queen is $1 - \frac{4}{52} = \frac{48}{52}$
- **General addition rule** for any events, A and B :

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \text{ and } B)$$

- Probability of drawing Queen or ♠
- $\frac{4}{52} + \frac{13}{52} - \frac{1}{52} = \frac{16}{52}$

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - 1 Linda is a bank teller?
 - 2 Linda is a bank teller and is active in the feminist movement?

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - 1 Linda is a bank teller?
 - 2 Linda is a bank teller and is active in the feminist movement?
- Famous example of the **conjunction fallacy** called the Linda problem

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - ① Linda is a bank teller?
 - ② Linda is a bank teller and is active in the feminist movement?
- Famous example of the **conjunction fallacy** called the Linda problem
 - Majority of respondents chose 2, but this is impossible!

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - 1 Linda is a bank teller?
 - 2 Linda is a bank teller and is active in the feminist movement?
- Famous example of the **conjunction fallacy** called the Linda problem
 - Majority of respondents chose 2, but this is impossible!
- **Law of total probability** for any events A and B :

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - 1 Linda is a bank teller?
 - 2 Linda is a bank teller and is active in the feminist movement?
- Famous example of the **conjunction fallacy** called the Linda problem
 - Majority of respondents chose 2, but this is impossible!
- **Law of total probability** for any events A and B :

$$\mathbb{P}(A) = \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B)$$

Conjunction fallacy

Linda is 31 years old, single, outspoken, and very bright. She majored in philosophy. As a student, she was deeply concerned with issues of discrimination and social justice, and also participated in anti-nuclear demonstrations.

- What is more probable?
 - 1 Linda is a bank teller?
 - 2 Linda is a bank teller and is active in the feminist movement?
- Famous example of the **conjunction fallacy** called the Linda problem
 - Majority of respondents chose 2, but this is impossible!
- **Law of total probability** for any events A and B :

$$\mathbb{P}(A) = \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B)$$

$$\mathbb{P}(\text{teller and feminist}) = \mathbb{P}(\text{teller}) - \mathbb{P}(\text{teller and not feminist})$$

Law of total probability

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

Law of total probability

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- What's the probability of randomly selecting a woman senator?

Law of total probability

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- What's the probability of randomly selecting a woman senator?

$$\mathbb{P}(\text{woman}) = \mathbb{P}(\text{woman and a Democrat}) + \mathbb{P}(\text{woman and not a Democrat})$$

Law of total probability

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- What's the probability of randomly selecting a woman senator?

$$\mathbb{P}(\text{woman}) = \mathbb{P}(\text{woman and a Democrat}) + \mathbb{P}(\text{woman and not a Democrat})$$

$$= \frac{16}{100} + \frac{10}{100} = \frac{26}{100}$$

Break time!

Conditional probability

- If we know that B has occurred, what is the probability of A ?
 - Conditioning our analysis on B having occurred.
- Examples:
 - Probability of two states going to war if they are both democracies?
 - Probability of a judge issuing a pro-choice ruling if they have daughters?
 - Probability of a coup in a country if it has a presidential system?
- Conditional probability extremely useful for data analysis.

Conditional probability definition

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

- How often A and B occur divided by how often B occurs.

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

- How often A and B occur divided by how often B occurs.
- **WARNING!** $\mathbb{P}(A|B)$ does **not**, in general, equal $\mathbb{P}(B|A)$

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

- How often A and B occur divided by how often B occurs.
- **WARNING!** $\mathbb{P}(A|B)$ does **not**, in general, equal $\mathbb{P}(B|A)$
 - $\mathbb{P}(\text{smart}|\text{in QSS})$ is high

Conditional probability definition

- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

- How often A and B occur divided by how often B occurs.
- **WARNING!** $\mathbb{P}(A|B)$ does **not**, in general, equal $\mathbb{P}(B|A)$
 - $\mathbb{P}(\text{smart}|\text{in QSS})$ is high
 - $\mathbb{P}(\text{in QSS}|\text{smart})$ is low

Conditional probability definition


- Definition: if $\mathbb{P}(B) > 0$ then the **conditional probability** of A given B is


$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \text{ and } B)}{\mathbb{P}(B)}$$

- How often A and B occur divided by how often B occurs.
- **WARNING!** $\mathbb{P}(A|B)$ does **not**, in general, equal $\mathbb{P}(B|A)$
 - $\mathbb{P}(\text{smart}|\text{in QSS})$ is high
 - $\mathbb{P}(\text{in QSS}|\text{smart})$ is low
 - There are many many smart people who are not in this class (tell your friends)
- If all outcomes equally likely:

$$\mathbb{P}(A|B) = \frac{\text{number of outcomes in both } A \text{ and } B}{\text{number of outcomes in just } B}$$

Conditional probability





US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Choose one senator at random from this population

US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Choose one senator at random from this population
- What is the probability of choosing a woman?

US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Choose one senator at random from this population
- What is the probability of choosing a woman?
 - $\mathbb{P}(\text{Woman}) = \frac{26}{100} = 0.26$

US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Choose one senator at random from this population
- What is the probability of choosing a woman?
 - $\mathbb{P}(\text{Woman}) = \frac{26}{100} = 0.26$
- What is the probability of choosing a Republican who is a woman?

US Senate example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Choose one senator at random from this population
- What is the probability of choosing a woman?
 - $\mathbb{P}(\text{Woman}) = \frac{26}{100} = 0.26$
- What is the probability of choosing a Republican who is a woman?
 - $\mathbb{P}(\text{Woman} \mid \text{Rep.}) = \frac{10/100}{53/100} \approx 0.19$

Conditional probability rules

- Multiplication rule:

$$\mathbb{P}(A \text{ and } B) = \mathbb{P}(A|B)\mathbb{P}(B) = \mathbb{P}(B|A)\mathbb{P}(A)$$

Multiplication rule example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Draw the names of two senators from a hat
- What's the probability that we draw two women?

Multiplication rule example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Draw the names of two senators from a hat
- What's the probability that we draw two women?
 - Let W_1 and W_2 be the events that 1st and 2nd draws are women

Multiplication rule example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100

- Draw the names of two senators from a hat
- What's the probability that we draw two women?
 - Let W_1 and W_2 be the events that 1st and 2nd draws are women
 - We could make a list of all possible pairs to draw and count them...


Multiplication rule example

	<i>Democrats</i>	<i>Republicans</i>	<i>Independents</i>	<i>Total</i>
<i>Men</i>	29	43	2	74
<i>Women</i>	16	10	0	26
<i>Total</i>	45	53	2	100


- Draw the names of two senators from a hat
- What's the probability that we draw two women?
 - Let W_1 and W_2 be the events that 1st and 2nd draws are women
 - We could make a list of all possible pairs to draw and count them...
 - Or we could just use the multiplication rule:

$$\mathbb{P}(W_1 \text{ and } W_2) = \mathbb{P}(W_1)\mathbb{P}(W_2|W_1)$$

Law of total probability




Law of total probability




- Conditional probability lets us restate the law of total probability

Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**


Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B) \\ &= \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|\text{ not } B)\mathbb{P}(\text{not } B)\end{aligned}$$


Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B) \\ &= \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|\text{ not } B)\mathbb{P}(\text{not } B)\end{aligned}$$


Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B) \\ &= \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|\text{ not } B)\mathbb{P}(\text{not } B)\end{aligned}$$


Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B) \\ &= \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|\text{ not } B)\mathbb{P}(\text{not } B)\end{aligned}$$

Law of total probability



- Conditional probability lets us restate the law of total probability
- **Law of total probability:**

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \text{ and } B) + \mathbb{P}(A \text{ and not } B) \\ &= \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|\text{ not } B)\mathbb{P}(\text{not } B)\end{aligned}$$

Sampling and independence

Sampling and independence

- Sampling > 1 with replacement: **independent draws**

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.
 - First draw doesn't affect second \leadsto independence

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.
 - First draw doesn't affect second \rightsquigarrow independence
- Sampling > 1 without replacement: **dependent draws**

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.
 - First draw doesn't affect second \rightsquigarrow independence
- Sampling > 1 without replacement: **dependent draws**
 - Randomly draw 1 senator, note the name, leave it out.

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.
 - First draw doesn't affect second \rightsquigarrow independence
- Sampling > 1 without replacement: **dependent draws**
 - Randomly draw 1 senator, note the name, leave it out.
 - Randomly draw 2nd senator from the remaining 99 senators.

Sampling and independence

- Sampling > 1 with replacement: **independent draws**
 - Randomly draw 1 senator, note the name, then put it back in the hat.
 - Shuffle, randomly draw 2nd senator, note the senator.
 - First draw doesn't affect second \rightsquigarrow independence
- Sampling > 1 without replacement: **dependent draws**
 - Randomly draw 1 senator, note the name, leave it out.
 - Randomly draw 2nd senator from the remaining 99 senators.
 - First draw affects the probability of the second.