#### **PSC4375: Randomized Experiments**

Week 1: Lecture 2

Prof. Weldzius

Villanova University

Slides Updated: 2025-01-22

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.
- Psychological studies show it isn't easy.

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.
- Psychological studies show it isn't easy.
- **Contact Hypothesis**: outgroup hostility diminished when people from different groups interact with one another.

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.
- Psychological studies show it isn't easy.
- **Contact Hypothesis**: outgroup hostility diminished when people from different groups interact with one another.
- Today we'll explore this question the context of support for gay marriage and contact with a member of the LGBT community.

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.
- Psychological studies show it isn't easy.
- **Contact Hypothesis**: outgroup hostility diminished when people from different groups interact with one another.
- Today we'll explore this question the context of support for gay marriage and contact with a member of the LGBT community.
  - $Y_i = \text{support for gay marriage } (1) \text{ or not } (0)$

- Question: can we effectively persuade people to change their minds?
- Hugely important question for political campaigns, companies, etc.
- Psychological studies show it isn't easy.
- Contact Hypothesis: outgroup hostility diminished when people from different groups interact with one another.
- Today we'll explore this question the context of support for gay marriage and contact with a member of the LGBT community.
  - $Y_i = \text{support for gay marriage } (1) \text{ or not } (0)$
  - $T_i = \text{contact with member of the LGBT community } (1) \text{ or not } (0)$

• What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen *i* have supported gay marriage if they had contact with a member of the LGBT community?

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen *i* have supported gay marriage if they had contact with a member of the LGBT community?
- Two potential outcomes:

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen *i* have supported gay marriage if they had contact with a member of the LGBT community?
- Two potential outcomes:
  - $Y_i(1)$ : would i have supported gay marriage if they **had** contact with a member of the LGBT community?

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen i have supported gay marriage if they had contact with a member of the LGBT community?
- Two potential outcomes:
  - $Y_i(1)$ : would i have supported gay marriage if they **had** contact with a member of the LGBT community?
  - $Y_i(0)$ : would i have supported gay marriage if they **didn't have** contact with a member of the LGBT community?

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen i have supported gay marriage if they had contact with a member of the LGBT community?
- Two potential outcomes:
  - $Y_i(1)$ : would i have supported gay marriage if they **had** contact with a member of the LGBT community?
  - $Y_i(0)$ : would *i* have supported gay marriage if they **didn't** have contact with a member of the LGBT community?
- Causal effect for citizen i:  $Y_i(1) Y_i(0)$

3 / 10

- What does " $T_i$  causes  $Y_i$ " mean?  $\rightsquigarrow$  counterfactuals, "what if"
- Would citizen *i* have supported gay marriage if they had contact with a member of the LGBT community?
- Two potential outcomes:
  - $Y_i(1)$ : would i have supported gay marriage if they **had** contact with a member of the LGBT community?
  - $Y_i(0)$ : would i have supported gay marriage if they **didn't have** contact with a member of the LGBT community?
- Causal effect for citizen i:  $Y_i(1) Y_i(0)$
- Fundamental problem of causal inference: only one of the two potential outcomes is observable.

3 / 10

• We will often refer to the **sample size** (number of units) as *n*.

- We will often refer to the **sample size** (number of units) as *n*.
- We often have *n* measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$

- We will often refer to the **sample size** (number of units) as *n*.
- We often have *n* measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$
- We often want sums: how many in our sample support gay marriage?

- We will often refer to the **sample size** (number of units) as *n*.
- We often have n measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$
- We often want sums: how many in our sample support gay marriage?

$$Y_1 + Y_2 + ... + Y_n$$

- We will often refer to the **sample size** (number of units) as *n*.
- We often have n measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$
- We often want sums: how many in our sample support gay marriage?

$$Y_1 + Y_2 + ... + Y_n$$

\* Notation is a bit clunky, so we often use the **Sigma notation**:

- We will often refer to the sample size (number of units) as n.
- We often have *n* measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$
- We often want sums: how many in our sample support gay marriage?

$$Y_1 + Y_2 + ... + Y_n$$

\* Notation is a bit clunky, so we often use the **Sigma notation**:

$$\sum_{i=1}^{n} Y_i = Y_1 + Y_2 + \dots + Y_n$$

- We will often refer to the **sample size** (number of units) as *n*.
- We often have n measurements of some variable:  $(Y_1, Y_2, ..., Y_n)$
- We often want sums: how many in our sample support gay marriage?

$$Y_1 + Y_2 + ... + Y_n$$

\* Notation is a bit clunky, so we often use the **Sigma notation**:

$$\sum_{i=1}^{n} Y_i = Y_1 + Y_2 + \dots + Y_n$$

\*  $\sum_{i=1}^{n}$  means sum each value from  $Y_1$  to  $Y_n$ 

• The **sample average** or **sample mean** is simply the sum of all values divided by the number of values.

- The **sample average** or **sample mean** is simply the sum of all values divided by the number of values.
- Sigma notation allows us to write this in a compact way:

- The sample average or sample mean is simply the sum of all values divided by the number of values.
- Sigma notation allows us to write this in a compact way:

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^{n} Y_i$$

- The sample average or sample mean is simply the sum of all values divided by the number of values.
- Sigma notation allows us to write this in a compact way:

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^{n} Y_i$$

\* Suppose we surveyed 6 people and 3 supported gay marriage:

5/10

- The sample average or sample mean is simply the sum of all values divided by the number of values.
- Sigma notation allows us to write this in a compact way:

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^{n} Y_i$$

\* Suppose we surveyed 6 people and 3 supported gay marriage:

$$\bar{Y} = \frac{1}{6}(1+1+1+0+0+0) = 0.5$$

Slides Updated: 2025-01-22

• We want to estimate the average causal effects over all units:

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}\left(Y_{i}(1)-Y_{i}(0)\right)$$

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}(Y_i(1)-Y_i(0))$$

\* Why can't we just calculate this quantity directly?

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}(Y_i(1)-Y_i(0))$$

\* Why can't we just calculate this quantity directly? \* What we can estimate instead:

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}(Y_i(1)-Y_i(0))$$

\* Why can't we just calculate this quantity directly? \* What we can estimate instead:

Difference in means  $= \bar{Y}_{treated} - \bar{Y}_{control}$ 

### Quantity of interest

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}\left(Y_{i}(1)-Y_{i}(0)\right)$$

\* Why can't we just calculate this quantity directly? \* What we can estimate instead:

Difference in means 
$$= \bar{Y}_{treated} - \bar{Y}_{control}$$

-  $\bar{Y}_{treated}$ : observed average outcome for treated group -  $\bar{Y}_{control}$ : observed average outcome for control group

# **Quantity of interest**

### **Quantity of interest**

• We want to estimate the average causal effects over all units:

Sample Average Treatment Effect (SATE) 
$$=\frac{1}{n}\sum_{i=1}^{n}\left(Y_{i}(1)-Y_{i}(0)\right)$$

\* Why can't we just calculate this quantity directly? \* What we can estimate instead:

Difference in means 
$$= \bar{Y}_{treated} - \bar{Y}_{control}$$

-  $\bar{Y}_{treated}$ : observed average outcome for treated group -  $\bar{Y}_{control}$ : observed average outcome for control group \* When will the difference-in-means be a good estimate of the SATE?

6/10

 Randomize control trial: each unit's treatment assignment is determined by chance

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.
  - Treatment and control group are identical on average

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.
  - Treatment and control group are identical on average
  - Similar on both observable and unobservable characteristics.

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.
  - Treatment and control group are identical on average
  - Similar on both observable and unobservable characteristics.
- $\bullet$  Control group  $\approx$  what would have happened to treatment group if they had taken control

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.
  - Treatment and control group are identical on average
  - Similar on both observable and unobservable characteristics.
- ullet Control group pprox what would have happened to treatment group if they had taken control
  - $\bar{Y}_{control} \approx \frac{1}{n} \sum_{i=1}^{n} Y_i(0)$

- Randomize control trial: each unit's treatment assignment is determined by chance
  - e.g., flip a coin; draw read and blue chips from a hat; etc.
- Randomization ensures balance between treatment and control group.
  - Treatment and control group are identical on average
  - Similar on both observable and unobservable characteristics.
- ullet Control group pprox what would have happened to treatment group if they had taken control
  - $\bar{Y}_{control} \approx \frac{1}{\underline{n}} \sum_{i=1}^{n} Y_i(0)$
  - $\bar{Y}_{treated} \bar{Y}_{control} \approx \mathsf{SATE}$

Placebo effects:

#### Placebo effects:

 Respondents will be affected by any intervention, even if they shouldn't have any effect

- Placebo effects:
  - Respondents will be affected by any intervention, even if they shouldn't have any effect
- Hawthorne effects:

#### Placeho effects:

 Respondents will be affected by any intervention, even if they shouldn't have any effect

#### • Hawthorne effects:

Respondents act differently just knowing that they are under study.

• Can we determine if randomization "worked"?

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on **pretreatment variable**

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on pretreatment variable
  - Pretreatment variable are those that are unaffected by treatment

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on pretreatment variable
  - Pretreatment variable are those that are unaffected by treatment
- We can check in the actual data for some pretreatment variable X

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on pretreatment variable
  - Pretreatment variable are those that are unaffected by treatment
- We can check in the actual data for some pretreatment variable X
  - $\bar{X}_{treated}$ : average value of variable for treated group

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on **pretreatment variable** 
  - Pretreatment variable are those that are unaffected by treatment
- We can check in the actual data for some pretreatment variable X
  - $X_{treated}$ : average value of variable for treated group
  - X<sub>control</sub>: average value of variable for control group

- Can we determine if randomization "worked"?
- If it did, we shouldn't see large differences between treatment and control group on pretreatment variable
  - Pretreatment variable are those that are unaffected by treatment
- ullet We can check in the actual data for some pretreatment variable X
  - $\bar{X}_{treated}$ : average value of variable for treated group
  - $\bar{X}_{control}$ : average value of variable for control group
  - ullet Under randomization,  $ar{X}_{treated} ar{X}_{control} pprox 0$

• Instead of 1 treatment, we might have multiple **treatment arms**:

10 / 10

- Instead of 1 treatment, we might have multiple **treatment arms**:
  - Control condition
  - Treatment A
  - Treatment B
  - Treatment C, etc.
- In this case, we will look at multiple comparisons:

- Instead of 1 treatment, we might have multiple **treatment arms**:
  - Control condition
  - Treatment A
  - Treatment B
  - Treatment C, etc.
- In this case, we will look at multiple comparisons:
  - $\bar{Y}_{treated,A} \bar{Y}_{control}$

- Instead of 1 treatment, we might have multiple **treatment arms**:
  - Control condition
  - Treatment A
  - Treatment B
  - Treatment C, etc.
- In this case, we will look at multiple comparisons:
  - $\bar{Y}_{treated,A} \bar{Y}_{control}$
  - $\bar{Y}_{treated B} \bar{Y}_{control}$

- Instead of 1 treatment, we might have multiple **treatment arms**:
  - Control condition
  - Treatment A
  - Treatment B
  - Treatment C, etc.
- In this case, we will look at multiple comparisons:
  - $\bar{Y}_{treated,A} \bar{Y}_{control}$
  - $\bar{Y}_{treated,B} \bar{Y}_{control}$
  - $\bar{Y}_{treated,A} \bar{Y}_{treated,B}$

- Instead of 1 treatment, we might have multiple **treatment arms**:
  - Control condition
  - Treatment A
  - Treatment B
  - Treatment C, etc.
- In this case, we will look at multiple comparisons:
  - $\bar{Y}_{treated,A} \bar{Y}_{control}$
  - $\bar{Y}_{treated.B} \bar{Y}_{control}$
  - $Y_{treated,A} Y_{treated,B}$
- If treatment arms are randomly assigned, these differences will be good estimators for each causal contrast.