Probability: Bayes Rule Slides from Prof. Blackwell (Harvard)

Prof. Weldzius

Villanova University

Slides Updated: 2025-03-19

QAnon



You meet a man named Steve and he tells you that he is a Republican. You have been interested in meeting someone who believes in the QAnon conspiracy theory. Given what you know about Steve, would you guess that he believes in OAnon or not?

QAnon



You meet a man named Steve and he tells you that he is a Republican. You have been interested in meeting someone who believes in the QAnon conspiracy theory. Given what you know about Steve, would you guess that he believes in OAnon or not?

 Common response: probably believes in QAnon since believers tend to be Republicans.


QAnon





You meet a man named Steve and he tells you that he is a Republican. You have been interested in meeting someone who believes in the QAnon conspiracy theory. Given what you know about Steve, would you guess that he believes in OAnon or not?


- Common response: probably believes in QAnon since believers tend to be Republicans.
- Base rate fallacy: ignores how uncommon QAnon believers are!















Chance a random Republican believes QAnon =





Chance a random Republican believes QAnon =




















Chance a random Republican believes QAnon =
$$\frac{\mathbb{P}(R|Q)\mathbb{P}(Q)}{\mathbb{P}(R|Q)\mathbb{P}(Q)} +$$



Chance a random Republican believes QAnon =
$$\frac{\mathbb{P}(R|Q)\mathbb{P}(Q)}{\mathbb{P}(R|Q)\mathbb{P}(Q)} +$$



Chance a random Republican believes QAnon =
$$\frac{\mathbb{P}(R|Q)\mathbb{P}(Q)}{\mathbb{P}(R|Q)\mathbb{P}(Q)} + \frac{\mathbb{P}(R|Q)\mathbb{P}(Q)}{\mathbb{P}(Q)}$$




Chance a random Republican believes QAnon =
$$\frac{\mathbb{P}(R|Q)\mathbb{P}(Q)}{\mathbb{P}(R|Q)\mathbb{P}(Q) + \mathbb{P}(R|\text{not }Q)\mathbb{P}(\text{not }Q)}$$

Bayes' rule



• Reverend Thomas Bayes (1701–61): English minister and statistician

Bayes' rule




- Reverend Thomas Bayes (1701–61): English minister and statistician
- Bayes' rule: if $\mathbb{P}(B) > 0$, then:

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(B \mid A)\mathbb{P}(A)}{\mathbb{P}(B)}$$

4/8

Bayes' rule



- Reverend Thomas Bayes (1701–61): English minister and statistician
- Bayes' rule: if $\mathbb{P}(B) > 0$, then:

$$\mathbb{P}(A\mid B) = \frac{\mathbb{P}(B\mid A)\mathbb{P}(A)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B\mid A)\mathbb{P}(A)}{\mathbb{P}(B\mid A)\mathbb{P}(A) + \mathbb{P}(B\mid \operatorname{not} A)\mathbb{P}(\operatorname{not} A)}$$

• What is the probability of some hypothesis given some evidence?

- What is the probability of some hypothesis given some evidence?
 - $\mathbb{P}(QAnon \mid Republican)$?

- What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.

- · What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - $\mathbb{P}(Republican \mid QAnon)$

- · What is the probability of some hypothesis given some evidence?
 - $\mathbb{P}(QAnon \mid Republican)$?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the **prior probability** of the hypothesis.

- · What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the prior probability of the hypothesis.
 - Prior: $\mathbb{P}(QAnon)$

- What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the prior probability of the hypothesis.
 - Prior: $\mathbb{P}(QAnon)$
 - **Posterior**: $\mathbb{P}(\mathsf{QAnon} \mid \mathsf{Republican})$

- What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the prior probability of the hypothesis.
 - Prior: ℙ(QAnon)
 - **Posterior**: $\mathbb{P}(\mathsf{QAnon} \mid \mathsf{Republican})$
- Applying Bayes' rule is often called updating the prior.

- What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the prior probability of the hypothesis.
 - Prior: ℙ(QAnon)
 - **Posterior**: $\mathbb{P}(\mathsf{QAnon} \mid \mathsf{Republican})$
- Applying Bayes' rule is often called updating the prior.
 - $\mathbb{P}(\mathsf{QAnon}) \leadsto \mathbb{P}(\mathsf{QAnon} \mid \mathsf{Republican})$

- What is the probability of some hypothesis given some evidence?
 - P(QAnon | Republican)?
- Often easier to know probability of evidence given hypothesis.
 - P(Republican | QAnon)
- Combine this with the prior probability of the hypothesis.
 - Prior: $\mathbb{P}(QAnon)$
 - **Posterior**: $\mathbb{P}(\mathsf{QAnon} \mid \mathsf{Republican})$
- Applying Bayes' rule is often called updating the prior.
 - $\mathbb{P}(QAnon) \rightsquigarrow \mathbb{P}(QAnon \mid Republican)$
 - How does the evidence change the chance of the hypothesis being true?

• Medical testing:

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$

- · Medical testing:
 - Want to know: $\mathbb{P}(Disease \mid Test Positive)$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$

- · Medical testing:
 - Want to know: $\mathbb{P}(Disease \mid Test Positive)$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$
- · Predicting traits from names:

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$
- · Predicting traits from names:
 - Want to know: $\mathbb{P}(African American | Last Name)$

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$
- · Predicting traits from names:
 - Want to know: $\mathbb{P}(African American | Last Name)$
 - Have: $\mathbb{P}(\text{Last Name} \mid \text{African American})$ and $\mathbb{P}(\text{African American})$

Uses of Bayes' rule

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$
- · Predicting traits from names:
 - Want to know: P(African American | Last Name)
 - Have: $\mathbb{P}(\text{Last Name} \mid \text{African American})$ and $\mathbb{P}(\text{African American})$
- Spam filtering:

Uses of Bayes' rule

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$
 - Have: $\mathbb{P}(\text{Test Positive} \mid \text{Disease})$ and $\mathbb{P}(\text{Disease})$
- · Predicting traits from names:
 - Want to know: P(African American | Last Name)
 - Have: $\mathbb{P}(\text{Last Name} \mid \text{African American})$ and $\mathbb{P}(\text{African American})$
- · Spam filtering:
 - Want to know: $\mathbb{P}(\operatorname{Spam} \mid \operatorname{Email} \operatorname{text})$

Uses of Bayes' rule

- · Medical testing:
 - Want to know: $\mathbb{P}(\text{Disease} \mid \text{Test Positive})$
 - Have: $\mathbb{P}(\mathsf{Test}\;\mathsf{Positive}\;|\;\mathsf{Disease})\;\mathsf{and}\;\mathbb{P}(\mathsf{Disease})$
- · Predicting traits from names:
 - Want to know: P(African American | Last Name)
 - Have: $\mathbb{P}(\text{Last Name} \mid \text{African American})$ and $\mathbb{P}(\text{African American})$
- · Spam filtering:
 - Want to know: $\mathbb{P}(\text{Spam} \mid \text{Email text})$
 - Have: $\mathbb{P}(\text{Email text} \mid \text{Spam})$ and $\mathbb{P}(\text{Spam})$

• Suppose you go and get a COVID-19 test and it comes back positive!

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be PT.

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.
 - Question: What is $\mathbb{P}(C \mid PT)$?

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.
 - Question: What is $\mathbb{P}(C \mid PT)$?
- · Components for calculating Bayes' rule:

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.
 - Question: What is $\mathbb{P}(C \mid PT)$?
- · Components for calculating Bayes' rule:
 - $\mathbb{P}(PT|C) = 0.7$: true positive rate

- · Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.
 - Question: What is $\mathbb{P}(C \mid PT)$?
- · Components for calculating Bayes' rule:
 - $\mathbb{P}(PT|C) = 0.7$: true positive rate
 - $\mathbb{P}(PT \mid \text{not } C) = 0.05$: false positive rate

- Suppose you go and get a COVID-19 test and it comes back positive!
 - Let a positive test be *PT*.
- What's the probability you actually have COVID-19?
 - Let having COVID be labeled *C*.
 - Question: What is $\mathbb{P}(C \mid PT)$?
- · Components for calculating Bayes' rule:
 - $\mathbb{P}(PT|C) = 0.7$: true positive rate
 - $\mathbb{P}(PT \mid \text{not } C) = 0.05$: false positive rate
 - $\mathbb{P}(C) = 0.001$ rough prevalance of active COVID cases.

• Use the law of total probability to get the denominator:

• Use the law of total probability to get the denominator:

$$\mathbb{P}(PT)$$

· Use the law of total probability to get the denominator:

$$\mathbb{P}(PT) = \mathbb{P}(PT \mid C)\mathbb{P}(C)$$

• Use the law of total probability to get the denominator:

$$\mathbb{P}(PT) = \mathbb{P}(PT \mid C)\mathbb{P}(C) + \mathbb{P}(PT \mid \text{not } C)\mathbb{P}(\text{not } C)$$

· Use the law of total probability to get the denominator:

$$\begin{split} \mathbb{P}(PT) &= \mathbb{P}(PT \mid C)\mathbb{P}(C) + \mathbb{P}(PT | \text{not } C)\mathbb{P}(\text{not } C) \\ &= (0.7 \times 0.001) + (0.05 \times 0.999) \end{split}$$

• Use the law of total probability to get the denominator:

$$\mathbb{P}(PT) = \mathbb{P}(PT \mid C)\mathbb{P}(C) + \mathbb{P}(PT \mid \text{not } C)\mathbb{P}(\text{not } C)$$

$$= (0.7 \times 0.001) + (0.05 \times 0.999)$$

$$= 0.051$$

• Use the law of total probability to get the denominator:

$$\mathbb{P}(PT) = \mathbb{P}(PT \mid C)\mathbb{P}(C) + \mathbb{P}(PT \mid \text{not } C)\mathbb{P}(\text{not } C)$$

$$= (0.7 \times 0.001) + (0.05 \times 0.999)$$

$$= 0.051$$

• Now plug in all values to Bayes' rule:

• Use the law of total probability to get the denominator:

$$\begin{split} \mathbb{P}(PT) &= \mathbb{P}(PT \mid C) \mathbb{P}(C) + \mathbb{P}(PT | \text{not } C) \mathbb{P}(\text{not } C) \\ &= (0.7 \times 0.001) + (0.05 \times 0.999) \\ &= 0.051 \end{split}$$

Now plug in all values to Bayes' rule:

$$\mathbb{P}(C \mid PT)$$

• Use the law of total probability to get the denominator:

$$\begin{split} \mathbb{P}(PT) &= \mathbb{P}(PT \mid C) \mathbb{P}(C) + \mathbb{P}(PT | \text{not } C) \mathbb{P}(\text{not } C) \\ &= (0.7 \times 0.001) + (0.05 \times 0.999) \\ &= 0.051 \end{split}$$

Now plug in all values to Bayes' rule:

$$\mathbb{P}(C \mid PT) = \frac{\mathbb{P}(PT \mid C)\mathbb{P}(C)}{\mathbb{P}(PT)}$$

• Use the law of total probability to get the denominator:

$$\begin{split} \mathbb{P}(PT) &= \mathbb{P}(PT \mid C) \mathbb{P}(C) + \mathbb{P}(PT | \text{not } C) \mathbb{P}(\text{not } C) \\ &= (0.7 \times 0.001) + (0.05 \times 0.999) \\ &= 0.051 \end{split}$$

• Now plug in all values to Bayes' rule:

$$\mathbb{P}(C \mid PT) = \frac{\mathbb{P}(PT \mid C)\mathbb{P}(C)}{\mathbb{P}(PT)} = \frac{0.7 \times 0.001}{0.051} \approx 0.014$$