

**Problem 1)**

Consider a potential problem in the half-space defined by  $z \geq 0$ , with Dirichlet boundary conditions on the plane  $z = 0$  (and at infinity).

- (a) Write down the appropriate Green function  $G(\vec{x}, \vec{x}')$ .
- (b) If the potential on the plane  $z = 0$  is specified to be  $\Phi = V$  inside a circle of radius  $a$  centered at the origin, and  $\Phi = 0$  outside that circle, find an integral expression for the potential at the point  $P$  specified in terms of cylindrical coordinates  $(\rho, \phi, z)$ .
- (c) Show that, along the axis of the circle ( $\rho = 0$ ), the potential is given by

$$\Phi = V \left( 1 - \frac{z}{\sqrt{a^2 + z^2}} \right). \quad (1)$$

- (d) Show that at large distances ( $\rho^2 + z^2 \gg a^2$ ) the potential can be expanded in a power series in  $(\rho^2 + z^2)^{-1}$ , and that the leading terms are

$$\Phi = \frac{Va^2}{2} \frac{z}{(\rho^2 + z^2)^{3/2}} \left[ 1 - \frac{3a^2}{4(\rho^2 + z^2)} + \frac{5(3\rho^2 a^2 + a^4)}{8(\rho^2 + z^2)^2} + \dots \right]. \quad (2)$$

Verify that the results of parts (c) and (d) are consistent with each other in their common range of validity.

- (a) A Green's function satisfying

$$\nabla^2 G(\vec{x}, \vec{x}') = -4\pi\delta(\vec{x} - \vec{x}') \quad (3)$$

is  $G(\vec{x}, \vec{x}') = |\vec{x} - \vec{x}'|^{-1}$ . Now, we must find a Green's function satisfying Dirichlet boundary conditions. That is, we take  $G \rightarrow G + F$ , where  $F$  solves Laplace's equation (in the half-space where  $z \geq 0$ ) and  $G + F$  satisfies the boundary condition  $\Phi = 0$  when  $z = 0$ . Such a choice is

$$G(\vec{x}, \vec{x}') = \frac{1}{|\vec{x}' - \vec{x}|} - \frac{1}{|\vec{x}' - \vec{y}|}, \quad (4)$$

where  $\vec{y}$  is  $\vec{x}$  translated over the  $xy$ -plane (i.e.  $z \rightarrow -z$  for  $\vec{y}$ ). It is clear then that the second term satisfies Laplace's equation for  $z \geq 0$  and that the sum of these two functions is identically zero on the  $xy$ -plane.

- (b) If we have the potential on the plane

$$\Phi(x, y, z = 0) = \begin{cases} V & x^2 + y^2 \leq a^2 \\ 0 & \text{otherwise,} \end{cases} \quad (5)$$

we can write the potential for  $z \geq 0$  as

$$\Phi = \frac{1}{4\pi\epsilon_0} \int_V d^3\vec{x}' G(\vec{x}, \vec{x}') \rho(\vec{x}') - \frac{1}{4\pi} \int_S dS' \Phi(\vec{x}') \frac{\partial G(\vec{x}, \vec{x}')}{\partial n'}. \quad (6)$$

Observe that the first term is zero since there is no charge. The second integral is over the circular surface of radius  $a$  in the  $xy$  plane such that

$$\begin{aligned}
 \Phi &= -\frac{V}{4\pi} \int_0^a \int_0^{2\pi} d\rho' d\phi' \rho' \left( -\frac{\partial G}{\partial z'} \right) \\
 &= \frac{V}{4\pi} \int_0^a \int_0^{2\pi} d\rho' d\phi' \rho' \frac{2z}{[(x' - x)^2 + (y' - y)^2 + z^2]^{3/2}} \\
 &= \boxed{\frac{Vz}{2\pi} \int_0^a \int_0^{2\pi} d\rho' d\phi' \frac{\rho'}{[(\rho' \cos \phi' - \rho \cos \phi)^2 + (\rho' \sin \phi' - \rho \sin \phi)^2 + z^2]^{3/2}}}.
 \end{aligned} \tag{7}$$

At this point, this is all we can do in full generality.

(c) We consider a special case along the  $z$  axis such that  $\rho = 0$ , which causes the previous expression to simply as

$$\begin{aligned}
 \Phi &= \frac{Vz}{2\pi} \int_0^a \int_0^{2\pi} d\rho' d\phi' \frac{\rho'}{(\rho'^2 + z^2)^{3/2}} = Vz \int_0^a \frac{\rho' d\rho'}{(\rho'^2 + z^2)^{3/2}} \\
 &= Vz \left[ \frac{1}{z} - \frac{1}{\sqrt{z^2 + a^2}} \right] = V \left[ 1 - \frac{z}{\sqrt{z^2 + a^2}} \right]
 \end{aligned} \tag{8}$$

as desired.

(d) There is another case in which we can do the integration, which is when we are far away from circular surface with radius  $a$  in the  $xy$ -plane (i.e.  $\rho^2 + z^2 \gg a^2$ ). In this case, we can write

$$\Phi = \frac{Vz}{2\pi} \frac{1}{(\rho^2 + z^2)^{3/2}} \int_0^a \int_0^{2\pi} d\rho' d\phi' \rho' \left[ 1 + \frac{\rho'^2 - 2\rho'\rho \cos(\phi' - \phi)}{\rho^2 + z^2} \right]^{-3/2}. \tag{9}$$

Notice that the expression in brackets is of the form  $(1+x)^n = 1+nx+[n(n-1)/2!]x^2+\dots$ , so the double integral becomes

$$\begin{aligned}
 &\int_0^a \int_0^{2\pi} d\rho' d\phi' \rho' \left[ 1 - \frac{3}{2} \left( \frac{\rho'^2 - 2\rho'\rho \cos(\phi' - \phi)}{\rho^2 + z^2} \right) + \frac{15}{8} \left( \frac{\rho'^2 - 2\rho'\rho \cos(\phi' - \phi)}{\rho^2 + z^2} \right)^2 + \dots \right] \\
 &= \pi a^2 - \frac{1}{\rho^2 + z^2} \frac{3}{2} \frac{2\pi}{4} a^4 + \frac{1}{(\rho^2 + z^2)^2} \frac{15}{8} \left[ \frac{2\pi}{6} a^6 + \frac{4\pi}{4} \rho^2 a^4 \right] + \dots \\
 &= \pi a^2 \left[ 1 - \frac{3a^2}{4(\rho^2 + z^2)} + \frac{5(3\rho^2 a^2 + a^4)}{8(\rho^2 + z^2)^2} + \dots \right].
 \end{aligned} \tag{10}$$

Plugging this back into Eq. (9), we find

$$\Phi = \boxed{\frac{Va^2}{2} \frac{z}{(\rho^2 + z^2)^{3/2}} \left[ 1 - \frac{3a^2}{4(\rho^2 + z^2)} + \frac{5(3\rho^2 a^2 + a^4)}{8(\rho^2 + z^2)^2} + \dots \right]}. \tag{11}$$

(d) One sanity check to perform is that the result of part (c) reduces to that of part (b) in the limit  $z \gg a$ , where  $\rho = 0$ . Observe that part (c) gives

$$\Phi = \frac{V}{2} \left( \frac{a}{z} \right)^2 \left[ 1 - \frac{3}{4} \left( \frac{a}{z} \right)^2 + \frac{5}{8} \left( \frac{a}{z} \right)^4 + \dots \right], \quad (12)$$

and expanding the result of part (b) in powers of  $\epsilon = a/z$ , we have

$$\begin{aligned} \Phi &= V \left[ 1 - (1 + \epsilon^2)^{-1/2} \right] = V \left( \frac{1}{2} \epsilon^2 - \frac{3}{8} \epsilon^4 + \frac{5}{16} \epsilon^6 + \dots \right) \\ &= \frac{V}{2} \epsilon^2 \left( 1 - \frac{3}{4} \epsilon^2 + \frac{5}{8} \epsilon^4 + \dots \right). \end{aligned} \quad (13)$$

One can see that Eq. (12) and Eq. (13) match as needed.

### Problem 2)

A two-dimensional potential problem is defined by two straight parallel line charges separated by a distance  $R$  with equal and opposite linear charge densities  $\lambda$  and  $-\lambda$

(a) Show by direct construction that the surface of constant potential  $V$  is a circular cylinder (circle in the transverse dimensions) and find the coordinates of the axis of the cylinder and its radius in terms of  $R$ ,  $\lambda$ , and  $V$ .

(b) Use the results of part (a) to show that the capacitance per unit length  $C$  of two right-circular cylindrical conductors, with radii  $a$  and  $b$  separated by a distance  $d > a + b$ , is

$$C = \frac{2\pi\epsilon_0}{\cosh^{-1} \left( \frac{d^2 - a^2 - b^2}{2ab} \right)}. \quad (14)$$

(c) Verify that the result for  $C$  agrees with the answer in Problem 1.7 of *Jackson textbook* in the appropriate limit and determine the next nonvanishing order correction in powers of  $a/d$  and  $b/d$ .

(d) Repeat the calculation of the capacitance per unit length for two cylinders inside each other ( $d < |b - a|$ ). Check the result for concentric cylinders ( $d = 0$ ).

Figure 1:

The potential a distance  $\rho$  from a line charge with linear charge density  $\lambda$  is

$$\Phi = \frac{\lambda}{2\pi\epsilon_0} \ln \left( \frac{\rho_0}{\rho} \right), \quad (15)$$

where  $\rho_0$  is some reference distance from the line charge where  $\Phi = 0$ . The potential of a configuration of two parallel, oppositely charged lines is just

$$\Phi = \Phi_+ + \Phi_- = \frac{\lambda}{2\pi\epsilon_0} \ln \left( \frac{\rho_-}{\rho_+} \right), \quad (16)$$

where  $\rho_{\pm}$  is just the distance between the point at which the potential is being evaluated and the line with charge per unit length  $\pm\lambda$ . Notice that the dependence on  $\rho_0$  cancels. If we set up our coordinate system as in Fig. 1, then we can express

$$\rho_{\pm} = \sqrt{\left(\frac{R}{2}\right)^2 + r^2 \pm 2\left(\frac{R}{2}\right)r \cos \phi}. \quad (17)$$

Hence, the potential in our coordinate system is

$$\Phi = \frac{\lambda}{4\pi\epsilon_0} \ln \left( \frac{R^2 + 4r^2 - 4Rr \cos \phi}{R^2 + 4r^2 + 4Rr \cos \phi} \right). \quad (18)$$

On the surface of constant potential  $\Phi = V$  we have

$$e^{4\pi\epsilon_0 V/\lambda} = \frac{R^2 + 4r^2 - 4Rr \cos \phi}{R^2 + 4r^2 + 4Rr \cos \phi} \quad (19)$$

$$a(R^2 + 4r^2 + 4Rr \cos \phi) = R^2 + 4r^2 - 4Rr \cos \phi \quad (20)$$

$$(a-1)R^2 + 4(a-1)r^2 + 4(a+1)Rr \cos \phi = 0 \quad (21)$$

where we have denoted  $a = \exp(4\pi\epsilon_0 V/\lambda)$ . It is difficult to see a resemblance to any familiar surface in this form, so let us transform back into Cartesian coordinates, where  $x = r \cos \phi$  and  $y = r \sin \phi$ :

$$(a-1)R^2 + 4(a-1)(x^2 + y^2) + 4(a+1)Rx = 0 \quad (22)$$

$$x^2 + \frac{a+1}{a-1}Rx + y^2 = -\frac{R^2}{4} \quad (23)$$

$$\left[ x + \frac{a+1}{a-1} \frac{R}{2} \right]^2 + y^2 = -\frac{R^2}{4} + \left( \frac{a+1}{a-1} \right)^2 \frac{R^2}{4} \quad (24)$$

$$\left[ x + \coth \left( \frac{2\pi\epsilon_0 V}{\lambda} \right) \frac{R}{2} \right]^2 + y^2 = \left[ \coth^2 \left( \frac{2\pi\epsilon_0 V}{\lambda} \right) - 1 \right] \left( \frac{R}{2} \right)^2 \quad (25)$$

$$\left[ x + \coth \left( \frac{2\pi\epsilon_0 V}{\lambda} \right) \frac{R}{2} \right]^2 + y^2 = \frac{(R/2)^2}{\sinh^2 (2\pi\epsilon_0 V/\lambda)}. \quad (26)$$

This is just the equation of a circle with center  $(-\coth(2\pi\epsilon_0 V/\lambda)[R/2], 0)$  and radius  $(R/2)/|\sinh(2\pi\epsilon_0 V/\lambda)|$ . Notice that if  $V > 0$  that the center of this circle is always Of course, the equipotential surface is a cylinder since the problem is translation invariant parallel to the lines.

(b) The capacitance of a setup with two conductors, one with charge  $Q$  and the other with charge  $-Q$ , is just  $C = Q/V$ , where  $V$  is the potential difference between the conductors. In this problem, we have two cylindrical conductors, with radii  $a$  and  $b$ , respectively. We know that conductors are equipotential surfaces. Hence, we can treat the potential on these surfaces as being set up by two line charges separated by some distance.

**Problem 3)**

An insulated, spherical, conducting shell of radius  $a$  is in a uniform electric field  $E_0$ . If the sphere is cut into two hemispheres by a plane perpendicular to the field, find the force required to prevent the hemispheres from separating

- (a) if the shell is uncharged;
- (b) if the total charge on the shell is  $Q$ .