

UCD Yang–Mills Existence and Mass-Gap Construction

Jaffe–Witten Specification: Compact-Group Regulator, Exact Schur
Renormalization, Reflection Positivity, and Spectral Transfer

Richard Wise

10 September 2026

Abstract

The Yang–Mills existence and mass-gap problem requires, for a compact simple gauge group G , a nontrivial quantum Yang–Mills theory on $\mathbb{R}^4$ with a physical Hamiltonian whose spectrum has a strictly positive interval above the vacuum containing no spectrum. This manuscript places the UCD construction directly into the Jaffe–Witten proof architecture.

The proof regulator is the standard compact-group lattice gauge system with link Hilbert space $L^2(G)$, gauge-invariant plaquette action, positive transfer structure, and independent ultraviolet and thermodynamic limits. For $SU(3)$ the Peter–Weyl decomposition

$$L^2(SU(3)) = \widehat{\bigoplus}_{p,q \geq 0} V_{p,q} \otimes V_{p,q}^*$$

gives a controlled finite-matrix verification hierarchy. The previously constructed 20-dimensional UCD quantum-link block is retained as an independent finite algebraic workbench; it is not substituted for $L^2(SU(3))$ in the continuum proof.

The manuscript proves an exact quadratic Schur-stability theorem. If the eliminated spectral sector has scale-uniform separation $\delta_* > 0$ and the local interface norm is η_k , then exact Feshbach reduction generates no term linear in the eliminated coupling and a local two-step path count A gives

$$\eta_{k+1} \leq \frac{A}{\delta_*} \eta_k^2.$$

For $x_0 = A\eta_0/\delta_* < 1$ this yields double-exponential contraction and the summable self-energy bound

$$\sum_{k \geq 0} \|\Sigma_k\| \leq \frac{\delta_*}{A^2} \frac{x_0^2}{1 - x_0^2}.$$

Consequently an initial spectral gap Δ_0 survives all subsequent exact Schur steps whenever

$$\Delta_0 > \frac{2\delta_*}{A^2} \frac{x_0^2}{1 - x_0^2}.$$

The 4-dimensional hypercubic plaquette overlap graph has degree at most 20 through shared links, giving the conservative ordered two-step path majorant $A \leq 400$. The executable UCD

$SU(3)$ block provides a numerical stress certificate for this theorem with $x_0 = 0.3294450611$ and a positive infinite-step gap floor 0.09985923249 in the specified strong-block test point.

The remaining Jaffe-Witten chain is formulated as explicit lemmas: all-field UV-to-block renormalization control, entrance of the standard Wilson/heat-kernel regulator into the Schur gap basin, convergence and nontriviality of the gauge-invariant Schwinger family, restoration of Euclidean symmetry, Osterwalder-Schrader reconstruction, and spectral-gap transfer. The package includes executable tests, machine-readable proof dependencies, Peter-Weyl truncation data, reflection-positive heat-kernel checks, Wilson continuum scaling, Schur certificates, and independent verification.

1. Exact Jaffe-Witten target

Let G be a compact simple Lie group. We seek a quantum gauge theory on $\mathbb{R}^4$ with Hilbert space $\mathcal{H}$, vacuum Ω , local gauge-invariant observable algebra $\mathcal{A}$, translation generators P_μ , positive Hamiltonian $H = P_0$, and

$$\boxed{\text{Spec}(H) \cap (0, \Delta) = \emptyset}$$

for some finite $\Delta > 0$.

The construction must satisfy axiomatic properties at least as strong as the Wightman/Osterwalder-Schrader formulations used in the Jaffe-Witten specification. In Euclidean form this means constructing gauge-invariant Schwinger distributions S_n with the positivity, covariance, locality/symmetry, regularity, and clustering properties required for reconstruction.

We organize the proof as

$$\boxed{A \implies B \implies C \implies D}$$

with:

- A.** a regulator- and volume-uniform positive physical spectral threshold;
- B.** existence of a non-Gaussian Euclidean continuum Yang-Mills state on $\mathbb{R}^4$;
- C.** Osterwalder-Schrader reconstruction of the physical relativistic theory;
- D.** transfer of the regulator-uniform spectral threshold to the reconstructed Hamiltonian.

2. Proof regulator: compact-group lattice Yang-Mills

Let $\Lambda_{a,L} \subset a\mathbb{Z}^4$ be a periodic hypercubic lattice with physical side L . Each oriented edge e carries a group element $U_e \in G$ with $U_{\bar{e}} = U_e^{-1}$. The Euclidean lattice configuration space is

$$\mathcal{C}_{a,L} = G^{E(\Lambda_{a,L})}.$$

For a vertex gauge transformation $g = (g_x)$,

$$U_{xy} \mapsto g_x U_{xy} g_y^{-1}.$$

For an oriented plaquette p ,

$$U_p = \prod_{e \in \partial p} U_e, \quad U_p \mapsto g_x U_p g_x^{-1},$$

so every class function of U_p is gauge invariant.

The Wilson action in a unitary representation R is

$$S_W(U) = \beta \sum_p \left(1 - \frac{1}{d_R} \operatorname{Re} \chi_R(U_p) \right).$$

The finite Euclidean measure is

$$d\mu_{a,L}(U) = Z_{a,L}^{-1} e^{-S_W(U)} \prod_e dU_e,$$

where dU is normalized Haar measure. Compactness of G makes every finite-volume integral well-defined.

For transfer-positivity arguments one may equivalently use the heat-kernel plaquette weight

$$K_t(g) = \sum_{\lambda \in \widehat{G}} d_\lambda e^{-tC_2(\lambda)} \chi_\lambda(g),$$

whose character coefficients are strictly positive and obey

$$K_t * K_s = K_{t+s}.$$

For any finite collection $g_i \in G$ and coefficients c_i ,

$$\sum_{i,j} \bar{c}_i c_j K_t(g_i^{-1} g_j) = \sum_{\lambda} d_\lambda e^{-tC_2(\lambda)} \left\| \sum_i c_i \pi_\lambda(g_i) \right\|_{HS}^2 \geq 0.$$

Thus K_t is of positive type for every $t > 0$. This is the algebraic core of the positive transfer kernel.

3. Hamiltonian regulator and Gauss law

On a spatial lattice, the link Hilbert space is

$$\mathcal{H}_e = L^2(G, dU),$$

and

$$\mathcal{H}_\Lambda = \bigotimes_{e \in E(\Lambda)} L^2(G).$$

Left and right regular actions implement local color rotations. The gauge-invariant physical space is

$$\mathcal{H}_\Lambda^{\text{phys}} = \{\psi : G_x \psi = \psi \ \forall x\}.$$

A Kogut-Susskind-type Hamiltonian is

$$H_{a,L} = \frac{g_0(a)^2}{2a} \sum_e (-\Delta_G)_e + \frac{1}{2g_0(a)^2 a} \sum_p (2d_R - \chi_R(U_p) - \overline{\chi_R(U_p)}).$$

Both terms are gauge invariant. The electric operator is positive. The magnetic bracket is nonnegative because $|\chi_R(U)| \leq d_R$.

4. Peter-Weyl finite verification hierarchy for SU(3)

The Peter-Weyl theorem gives

$$L^2(SU(3)) = \widehat{\bigoplus_{p,q \geq 0} V_{p,q} \otimes V_{p,q}^*}.$$

The irreducible representation (p, q) has

$$d_{p,q} = \frac{(p+1)(q+1)(p+q+2)}{2}$$

and quadratic Casimir

$$C_2(p, q) = \frac{p^2 + q^2 + pq + 3p + 3q}{3}.$$

Hence

$$C_F = C_2(1, 0) = \frac{4}{3}, \quad C_A = C_2(1, 1) = 3.$$

Define the finite Peter-Weyl truncation

$$\mathcal{H}_K = \bigoplus_{p+q \leq K} V_{p,q} \otimes V_{p,q}^*.$$

Then $P_K \rightarrow I$ strongly because the algebraic Peter-Weyl sum is dense. The executable dimensions are

$$1, 19, 155, 805, 3136, 9996, 27468, 67320, 150645, \dots$$

for $K = 0, 1, \dots, 8$.

Multiplication by a fundamental matrix coefficient obeys the exact tensor rule

$$(p, q) \otimes (1, 0) = (p+1, q) \oplus (p-1, q+1) \oplus (p, q-1),$$

with terms of negative Dynkin label omitted. Therefore, on every finite algebraic Peter-Weyl core vector, $P_K U P_K$ agrees with U for all sufficiently large K . This gives a finite-matrix verification route to the genuine $L^2(SU(3))$ link regulator.

5. UCD native refinement family

The UCD spatial refinement family is

$$\Gamma_{r,s} = \mathbb{Z}_3 2^{r+s} \times \mathbb{Z}_2 2^{r+s} \times \mathbb{Z}_2 2^{r+s},$$

with

$$a_r = L_* 2^{-r}.$$

The parameter r removes the short-distance regulator; s supplies independent thermodynamic enlargement. A fixed physical block of side L_B contains

$$n_r = \frac{L_B}{a_r}, \quad n_r a_r = L_B.$$

The executable refinement ledger verifies this identity exactly in dyadic arithmetic for the tested range.

The finite UCD carrier

$$\mathcal{H}_{504} = V_7 \otimes \mathbb{C}[\mathbb{Z}_3] \otimes \mathbb{C}[\text{Dic}_6]$$

has a separately proved ring translation

$$\Delta_{504} = 2\kappa e^{-1/72}.$$

Its exact Schur inheritance identity

$$H_{\text{eff}}(z) = H_{cc} - H_{cl}(H_{\ell\ell} - zI)^{-1}H_{\ell c}$$

supplies the finite-registry model for information-preserving elimination. This finite H504 gap is not inserted as the continuum Yang-Mills mass; it supplies algebraic Schur structure and a finite-registry sector.

6. Local Yang-Mills continuum asymptotics

For a smooth connection, a small plaquette satisfies

$$U_p = \exp(a^2 F_{\mu\nu} + O(a^3)).$$

After taking the real trace, the linear anti-Hermitian term cancels and

$$\text{ReTr}(I - U_p) = \frac{a^4}{2} \text{Tr}(F_{\mu\nu}^\dagger F_{\mu\nu}) + O(a^6).$$

The executable constant-color test uses

$$U_p = \exp(ia^2 X)$$

and verifies that the scaled density tends to $\frac{1}{2} \text{Tr}(X^2)$ with the expected $O(a^4)$ remainder in that commuting test.

7. Independent finite quantum-link workbench

The finite UCD $SU(3)$ workbench uses

$$\mathcal{H}_{\text{link}} = \Lambda^3 \mathbb{C}^6, \quad \dim \mathcal{H}_{\text{link}} = 20,$$

with

$$L^a = \sum_{ij} t_{ij}^a c_{L,i}^\dagger c_{L,j}, \quad R^a = \sum_{ij} t_{ij}^a c_{R,i}^\dagger c_{R,j}, \quad U_{ij} = c_{L,i}^\dagger c_{R,j}.$$

The machine verifies left/right $SU(3)$ covariance, $[L^a, R^b] = 0$, and

$$\sum_a t^a t^a = \frac{4}{3} I_3$$

to numerical roundoff. Exact Gauss reduction of one plaquette gives an 18-dimensional gauge-singlet sector.

The finite interacting diagnostic Hamiltonian is

$$H_B(g, \lambda; L_B) = \frac{g^2}{4L_B} Q_E - \frac{1}{2g^2 L_B} (B_p + B_p^\dagger) - \frac{\lambda}{L_B} D.$$

This block is used to test Schur, spectral-parent, transfer-matrix, and contraction machinery. The continuum proof regulator remains the compact-group $L^2(G)$ system of Sections 2-4.

8. Exact Feshbach-Schur theorem

Let a self-adjoint operator be decomposed by complementary projections P, Q :

$$H = \begin{pmatrix} A & B \\ B^\dagger & D \end{pmatrix}.$$

If $D - E$ is invertible, define

$$F_E(H) = A - E - B(D - E)^{-1} B^\dagger.$$

Then

$$0 \in \text{Spec } F_E(H) \iff E \in \text{Spec } H$$

away from $\text{Spec } D$. Moreover, if

$$D - E \geq \delta Q,$$

then

$$\|B(D - E)^{-1}B^\dagger\| \leq \frac{\|B\|^2}{\delta}.$$

The correction is exactly quadratic in the retained-eliminated coupling B . There is no first-order Schur self-energy term.

The finite $SU(3)$ workbench verifies the four lowest full eigenvalues against their energy-dependent Schur operators with maximum residual 1.33×10^{-14} .

9. Quadratic Schur contraction theorem

Consider a sequence of exact block eliminations. At scale k , absorb the complete PHP term into the next retained Hamiltonian and write the remaining retained-eliminated coupling as a sum of local interface operators whose local interaction norm is at most η_k .

Assume:

1. the eliminated sector in the spectral window is separated by

$$Q(H_k - E)Q \geq \delta_* Q,$$

with $\delta_* > 0$ independent of k ;

2. a retained local cell participates in at most A ordered two-step paths through one eliminated cell;
3. the next-scale residual interaction norm is defined by summing the norms of local Schur-generated paths touching a fixed retained cell.

Then each two-step term satisfies

$$\|B_Y(Q(H_k - E)Q)^{-1}B_Z^\dagger\| \leq \frac{\eta_k^2}{\delta_*},$$

and at most A such ordered paths contribute. Therefore

$$\eta_{k+1} \leq \frac{A}{\delta_*} \eta_k^2.$$

Set

$$x_k = \frac{A\eta_k}{\delta_*}.$$

Then

$$x_{k+1} \leq x_k^2.$$

If $0 < x_0 < 1$,

$$x_k \leq x_0^{2^k}$$

and therefore

$$\eta_k \leq \frac{\delta_*}{A} x_0^{2^k}.$$

The k th self-energy satisfies

$$\epsilon_k \leq \frac{\eta_k^2}{\delta_*} \leq \frac{\delta_*}{A^2} x_0^{2^{k+1}}.$$

Since $2^{k+1} \geq 2(k+1)$,

$$\sum_{k=0}^{\infty} x_0^{2^{k+1}} \leq \sum_{j=1}^{\infty} x_0^{2^j} = \frac{x_0^2}{1 - x_0^2}.$$

Hence

$$\boxed{\sum_{k \geq 0} \epsilon_k \leq \frac{\delta_*}{A^2} \frac{x_0^2}{1 - x_0^2}.}$$

By Weyl's inequality, a perturbation of operator norm ϵ moves each of the two lowest spectral edges by at most ϵ , so the gap decreases by at most 2ϵ . Thus an initial gap Δ_0 satisfies

$$\boxed{\Delta_{\infty} \geq \Delta_0 - \frac{2\delta_*}{A^2} \frac{x_0^2}{1 - x_0^2}.}$$

This is a regulator-step-independent sufficient criterion for gap survival through an exact Schur tower.

10. Four-dimensional local path constant

On a 4-dimensional hypercubic lattice, a link belongs to

$$2(d-1) = 6$$

plaquettes. Fix one plaquette. Each of its four boundary links is shared with five other plaquettes. Distinct edge-sharing neighboring plaquettes therefore number at most

$$D = 4(6-1) = 20.$$

An ordered Schur path retained $\rightarrow$ eliminated $\rightarrow$ retained is bounded by

$$\boxed{A \leq D^2 = 400.}$$

This is deliberately a path-count majorant; additional cancellations can only reduce it.

For the finite quantum-link stress point $g = 8$, $\lambda = 0.05$, $L_B = 1$, the machine obtains

$$\Delta_0 = 0.09998915391358655,$$

$$\eta_0 = 0.0703125,$$

$$\delta_0 = 170.74167028648398.$$

Taking the conservative invariant target

$$\delta_* = \frac{1}{2}\delta_0 = 85.37083514324199$$

and $A = 400$ gives

$$x_0 = 0.32944506110089744 < 1,$$

$$\sum_k \epsilon_k \leq 6.496071390629655 \times 10^{-5},$$

and

$$\boxed{\Delta_\infty \geq 0.09985923248577395 > 0.}$$

The corresponding separation floor is

$$170.74154036505618 > \delta_*;$$

hence the assumed half-separation is self-consistent for this abstract Schur tower.

11. Strong-coupling gap basin for the standard compact-group Hamiltonian

For the genuine $L^2(G)$ Hamiltonian, split

$$H = H_E + V_B.$$

The unperturbed electric operator is a sum of independent positive group Laplacians. By Peter-Weyl,

$$-\Delta_G|_{V_\lambda \otimes V_\lambda^*} = C_2(\lambda).$$

A rigorous Euclidean strong-coupling endpoint is also available for the standard lattice theory. In the t'Hooft-scaled normalization of Shen, Zhu, and Zhu, the $SU(N)$ lattice Yang-Mills measure in dimension $d > 1$ has a unique infinite-volume Gibbs state, satisfies Poincaré and logarithmic-Sobolev inequalities, and exhibits exponential decay of correlations when

$$|\beta| < \frac{1}{16(d-1)}.$$

Hence in four dimensions the explicit sufficient region is

$$|\beta| < \frac{1}{48}.$$

This supplies a quantitative infrared endpoint for the A→B chain: once the exact blocked Wilson/heat-kernel action is shown to enter the theorem's strong-coupling action class with controlled residual interactions, infinite-volume convergence and exponential clustering follow from an existing rigorous theorem. The ultraviolet-to-infrared burden is thereby localized to the matching estimate that carries the asymptotically-free regulator into that certified basin while preserving gauge invariance, locality, and reflection positivity.

For compact simple G , every nontrivial irreducible representation has

$$C_2(\lambda) > 0.$$

Thus the product electric vacuum is unique and has a local spectral gap. The magnetic plaquette term is a bounded finite-range perturbation because $|\chi_R(U)| \leq d_R$. Standard relatively-bounded perturbation theorems for gapped lattice systems therefore give a volume-independent gapped phase whenever the local magnetic interaction is sufficiently small relative to the electric gap. For $SU(3)$ the dimensionless local scale ratio behaves as

$$\frac{\|V_p\|}{\Delta_E} \lesssim \frac{9}{g^4},$$

so the large- g region lies in a rigorous gapped strong-coupling basin.

The purpose of Sections 8-10 is to give a UCD-native exact Schur mechanism for maintaining such a basin under subsequent coarse elimination, with explicit summable error control.

12. Ultraviolet asymptotic freedom and renormalized trajectory

For pure $SU(N)$,

$$\mu \frac{dg}{d\mu} = -b_0 g^3 + O(g^5), \quad b_0 = \frac{11N}{48\pi^2} > 0.$$

For $SU(3)$,

$$b_0 = \frac{11}{16\pi^2} = 0.0696583137541 \dots$$

At one loop, fixing a block coupling g_B at physical scale L_B gives

$$\frac{1}{g_0(a)^2} = \frac{1}{g_B^2} + 2b_0 \log \frac{L_B}{a}.$$

Thus

$$g_0(a) \rightarrow 0 \quad (a \rightarrow 0).$$

The executable trajectory verifies the algebraic round-trip relation to machine precision. Balaban's four-dimensional small-field renormalization construction provides rigorous localized effective actions and coupling-constant renormalization in the weak-coupling ultraviolet regime. The complete Jaffe–Witten bridge is the all-field continuation from that ultraviolet regime into a fixed-physical-scale gapped basin with constants uniform in a and L .

We isolate that step as the following lemma.

Bridge Lemma A (UV-to-gap-basin). There exist $L_B > 0$, $a_0 > 0$, and regulator couplings $g_0(a)$ for $0 < a < a_0$ such that the exact gauge-invariant Wilson/heat-kernel RG map to scale L_B produces an effective Hamiltonian satisfying the hypotheses of Theorem 9 with constants $A, \delta_*, \eta_0, \Delta_0$ independent of a and thermodynamic volume, with

$$\frac{A\eta_0}{\delta_*} < 1$$

and

$$\Delta_0 > \frac{2\delta_*}{A^2} \frac{(A\eta_0/\delta_*)^2}{1 - (A\eta_0/\delta_*)^2}.$$

Once Bridge Lemma A is established, Theorem 9 produces a regulator- and volume-uniform positive spectral threshold $m_* > 0$.

13. A: regulator-uniform mass gap

Let $P_{0,a,L}$ be the finite-volume vacuum projection. Under Bridge Lemma A and the exact Schur contraction theorem,

$$H_{a,L} - E_{0,a,L} \geq m_*(I - P_{0,a,L})$$

with one $m_* > 0$ independent of sufficiently small a and sufficiently large L . Therefore

$$\inf_{a < a_0, L > L_0} (E_1(a, L) - E_0(a, L)) \geq m_* > 0.$$

This is precisely the form of uniform finite-volume gap singled out in the Jaffe–Witten discussion as the spectral input needed for the thermodynamic construction.

14. B: infinite-volume local state

For a bounded gauge-invariant observable A supported in a fixed compact physical region K , define

$$\omega_{a,L}(A) = \langle \Omega_{a,L}, A \Omega_{a,L} \rangle.$$

A volume-uniform gap and finite-range locality give exponential decay of connected correlations in the Hamiltonian lattice theory. The required thermodynamic theorem is:

Bridge Lemma B1 (thermodynamic uniqueness and convergence). For each sufficiently small a , the local expectations $\omega_{a,L}(A)$ converge as $L \rightarrow \infty$, independently of boundary conditions compatible with the vacuum phase, and the limit obeys a uniform exponential clustering estimate

$$|\omega_a(AB) - \omega_a(A)\omega_a(B)| \leq C_{A,B} e^{-m_* d(A,B)}.$$

The quasi-local algebra is the norm closure

$$\mathcal{A}_a = \overline{\bigcup_{K \in \mathbb{R}^3} \mathcal{A}_a(K)}.$$

The thermodynamic state ω_a is positive and normalized by construction.

15. B: continuum Schwinger family

Let $\mathcal{O}_i^{(a)}(f_i)$ be renormalized gauge-invariant local composite observables smeared with test functions. Define

$$S_n^{(a)}(f_1, \dots, f_n) = \omega_a \left(\mathcal{O}_1^{(a)}(f_1) \cdots \mathcal{O}_n^{(a)}(f_n) \right).$$

The continuum construction requires scale-dependent field/operator renormalizations $Z_{\mathcal{O}}(a)$ and counterterms fixed by gauge-invariant renormalization conditions.

Bridge Lemma B2 (continuum convergence). For every finite collection of gauge-invariant local test observables, the renormalized distributions converge in $\mathcal{S}'(\mathbb{R}^{4n})$:

$$S_n = \lim_{a \rightarrow 0} S_n^{(a)}.$$

The bounds are uniform on bounded sets of Schwartz test functions and compatible with the exponential clustering constant m_* .

Bridge Lemma B3 (nontriviality). At least one connected gauge-invariant continuum distribution is nonzero, equivalently

$$S_n^T \neq 0$$

for some $n \geq 4$, or another equivalent non-Gaussian observable criterion holds.

The finite workbench already contains nonzero connected Wilson spectral weight; B3 is its continuum persistence statement.

15.1 Local curvature fields, stress tensor, and short-distance operator structure

The Jaffe–Witten specification requires the continuum theory to contain local gauge-invariant quantum fields corresponding to classical gauge-invariant differential polynomials in the curvature and its covariant derivatives. Let

$$P(F, DF, D^2F, \dots)$$

be such a polynomial of engineering dimension d_P . On a lattice, curvature is represented by oriented plaquette combinations. For $SU(N)$ one convenient Lie-algebra-valued representative is

$$F_{\mu\nu}^{(a)}(x) = \frac{1}{2ia^2g_0(a)} \left[U_{\mu\nu}(x) - U_{\mu\nu}(x)^\dagger - \frac{1}{N} \text{Tr}(U_{\mu\nu}(x) - U_{\mu\nu}(x)^\dagger)I \right],$$

and covariant finite differences give approximants to $D_\rho F_{\mu\nu}$. Gauge-invariant lattice composites $P_j^{(a)}$ are then mixed and renormalized by a finite matrix at each engineering dimension,

$$\mathcal{O}_P^{(a)} = \sum_{j: d_j \leq d_P} Z_{Pj}(a, \mu) P_j^{(a)}.$$

The required continuum statement is:

Bridge Lemma B4 (composite-field convergence). For every gauge-invariant differential polynomial $P(F, DF, \dots)$ there is a renormalized lattice approximant $\mathcal{O}_P^{(a)}$ such that all mixed Schwinger distributions containing finitely many such insertions converge in $\mathcal{S}'$ as $a \rightarrow 0$, and the limiting operators transform covariantly under the restored Euclidean group.

The stress tensor is obtained by differentiating an anisotropic/background-metric lattice action with respect to the metric and then renormalizing the finite set of operators allowed by symmetry. Its continuum Ward identity is

$$\partial^\mu T_{\mu\nu} = 0,$$

and after OS reconstruction

$$H = \int_{\mathbb{R}^3} T_{00}(0, \mathbf{x}) d^3x$$

in the standard quadratic-form sense on the energy domain.

At short distance, asymptotic freedom fixes the ultraviolet coefficients in products of local fields. The required operator-product statement is

$$\mathcal{O}_P(x) \mathcal{O}_Q(0) \sim \sum_R C_{PQ}^R(x, \mu, g(\mu)) \mathcal{O}_R(0), \quad |x| \downarrow 0,$$

with $g(\mu) \rightarrow 0$ as $\mu \rightarrow \infty$ and Wilson coefficients having the perturbative asymptotic expansion determined by Yang-Mills renormalization.

Bridge Lemma B5 (short-distance matching). The renormalized continuum composite fields of B4 obey the Yang-Mills Ward identities and their short-distance Schwinger distributions have the asymptotic-freedom/OPE singular structure fixed by perturbative renormalization. In particular a conserved renormalized stress tensor exists.

B4 and B5 are the exact operator-theoretic form of the local-field and ultraviolet requirements in the Jaffe-Witten problem statement; they prevent a merely gapped lattice theory from being substituted for four-dimensional quantum Yang-Mills.

16. Reflection positivity and passage to the limit

At finite regulator the Wilson approximation is reflection positive; the heat-kernel formulation exhibits positivity directly through nonnegative character coefficients.

Let F be a polynomial in positive-time gauge-invariant smeared observables. Then

$$\langle \Theta F, F \rangle_{a,L} \geq 0.$$

If Bridge Lemma B2 gives convergence of every correlator entering this finite quadratic form, then

$$\langle \Theta F, F \rangle = \lim_{a \rightarrow 0} \lim_{L \rightarrow \infty} \langle \Theta F, F \rangle_{a,L} \geq 0.$$

Therefore reflection positivity is closed under the continuum limit once the Schwinger distributions converge.

The same argument passes permutation symmetry. Translation invariance passes from the translation-invariant regulators. Full Euclidean rotation invariance requires restoration from the hypercubic subgroup:

Bridge Lemma C1 (Euclidean restoration). The limiting Schwinger distributions are invariant under $E(4)$ rather than only the hypercubic subgroup, with all Lorentz-breaking irrelevant operators vanishing under the renormalized trajectory.

17. C: Osterwalder-Schrader reconstruction

Assume B1-B5 and C1 together with the standard OS regularity bounds. Then the limiting Schwinger family satisfies the reconstruction hypotheses. The OS quotient is formed from positive-time test polynomials by

$$(F, G)_{OS} = \langle \Theta F, G \rangle,$$

modulo null vectors and completed to a Hilbert space $\mathcal{H}$. Euclidean time translations generate a positive contraction semigroup

$$T(t) = e^{-tH}, \quad H \geq 0.$$

Spatial translations and rotations continue to unitary operators, analytic continuation produces the Poincaré representation, and the reconstructed local observable fields satisfy the corresponding Wightman-strength axioms.

Thus

$$\boxed{\{S_n\}_{n \geq 0} \xrightarrow{OS} (\mathcal{H}, \Omega, H, P_\mu, \mathcal{A}).}$$

18. D: spectral-gap transfer

For a local gauge-invariant observable A with $\langle \Omega, A\Omega \rangle = 0$,

$$C_A(t) = \langle \Omega, A e^{-tH} A \Omega \rangle = \int_{[0, \infty)} e^{-Et} d\rho_A(E),$$

where ρ_A is a positive spectral measure.

If the uniform regulator gap gives

$$|C_A^{(a,L)}(t)| \leq C_A e^{-m_* t}$$

with C_A stable under the limits, then

$$|C_A(t)| \leq C_A e^{-m_* t}.$$

Suppose $\rho_A([E_1, E_2]) > 0$ for some

$$0 < E_1 \leq E_2 < m_*.$$

Then positivity implies

$$C_A(t) \geq e^{-E_2 t} \rho_A([E_1, E_2]),$$

which cannot be bounded by $C_A e^{-m_* t}$ as $t \rightarrow \infty$. Hence

$$\rho_A((0, m_*)) = 0.$$

If local gauge-invariant vectors $A\Omega$ are dense in $\Omega^\perp$, then the bounded spectral projection $1_{(0, m_*)}(H)$ vanishes on a dense set and therefore vanishes identically. Thus

$$\boxed{\text{Spec}(H) \subset \{0\} \cup [m_*, \infty)}.$$

19. Finite upper edge and nontrivial spectral support

The Jaffe–Witten definition also requires the supremal gap parameter to be finite. It is enough to construct one non-vacuum local state of finite energy. For a compactly supported gauge-invariant local operator A in the Hamiltonian form domain,

$$0 < \|(I - P_0)A\Omega\| < \infty,$$

and

$$\frac{\langle A\Omega, H A\Omega \rangle}{\|(I - P_0)A\Omega\|^2} < \infty$$

places spectrum at finite positive energy. The continuum nontriviality lemma B3 supplies the required non-vacuum local sector.

20. Arbitrary compact simple gauge group

The argument above is not special to $SU(3)$ at the structural level. For compact simple G :

1. Haar measure exists and is finite.
2. Peter-Weyl gives

$$L^2(G) = \widehat{\bigoplus_{\lambda \in \widehat{G}} V_\lambda \otimes V_\lambda^*}.$$

3. The Laplace-Casimir eigenvalue obeys $C_2(\lambda) > 0$ for every nontrivial irreducible representation.
4. Heat-kernel character coefficients are positive.
5. Wilson/character plaquette observables are local and gauge invariant.
6. The Schur theorem is representation independent.
7. The OS reconstruction and spectral-transfer arguments are group independent.

The group-dependent part is therefore concentrated in the uniform RG estimates of Bridge Lemmas A and B2. A complete “for any compact simple G ” proof supplies those estimates for each compact simple group, with no requirement that the numerical constant m_* be universal across different groups.

21. Positivity-bootstrap cross-check lane

Hermitian positivity, reflection positivity, and Schwinger-Dyson loop equations can be imposed simultaneously on finite sets of Wilson loops. The 2025 $SU(3)$ lattice bootstrap literature demonstrates rigorous expectation-value bounds from these constraints in 2D, 3D, and 4D. In this package that machinery is assigned as an independent finite-regulator cross-check lane:

Wilson loop algebra + reflection positivity + SD equations $\implies$ certified convex feasible region.

The natural next use is to bound the renormalized block parameters $(\eta_k, \delta_k, \Delta_k)$ entering Bridge Lemma A rather than merely bound the plaquette expectation.

22. Machine-verifiable results

The package runs the following independent checks:

- $SU(3)$ left/right covariance and fundamental Casimir;
- exact 18-dimensional finite Gauss-sector construction;
- positive interacting finite-workbench block gap;
- low-spectrum Feshbach-Schur isospectrality;
- exact spectral-parent inequality;
- positive self-adjoint transfer matrix and transfer-gap identity;
- positive-kernel preservation under coarse pullback;
- nonzero gauge-invariant finite spectral weight and exponential spectral bound;
- Wilson small-plaquette continuum scaling;
- one-loop asymptotically-free trajectory arithmetic;
- exact $SU(3)$ Peter-Weyl dimensions and Casimir values;
- fundamental tensor-product adjacency graph;
- heat-kernel positive coefficients and convolution semigroup coefficients;
- 4D hypercubic overlap/path-count constants;
- quadratic Schur contraction and infinite-step gap-floor certificate.

The current automated test suite returns **10/10 tests passed**. The prior independent finite-block verifier returns **11/11 checks passed**.

23. Jaffe-Witten proof ledger

The machine-readable file `results/jaffe_witten_obligation_ledger_v4.csv` maps every theorem to its dependencies. The shortest remaining $SU(3)$ proof chain is:

all-field UV RG control
↓
entry into the exact Schur gap basin
↓
 $\inf_{a,L} \Delta(a, L) = m_* > 0$
↓
thermodynamic + continuum Schwinger convergence
↓
Euclidean symmetry + OS axioms
↓
OS reconstruction
↓
 $\text{Spec}(H) \subset \{0\} \cup [m_*, \infty)$.

The ledger deliberately separates a theorem already proved in the package, a standard theorem invoked from the literature, and a bridge lemma that must be discharged by the UCD/Wilson construction. That separation is part of the proof object: no finite-matrix eigenvalue is silently relabeled as a continuum mass.

24. Reproducibility

Install and run:

```
python -m pip install -r requirements.txt
python run_all.py
python -m pytest -q
```

The GitHub Actions workflow runs the same verification on each push and pull request. Every release file is hashed in `MANIFEST.sha256`.

25. Conclusion

The Jaffe-Witten problem has been converted into an explicit finite dependency graph. The compact-group proof regulator, gauge symmetry, Gauss sector, local Yang-Mills continuum asymptotics, Peter-Weyl finite verification hierarchy, positive Euclidean transfer structure, exact Feshbach-Schur identity, and a quantitative quadratic Schur gap-stability theorem are all written in one common normalization.

The new quantitative result is the exact summable Schur criterion

$$\Delta_\infty \geq \Delta_0 - \frac{2\delta_*}{A^2} \frac{(A\eta_0/\delta_*)^2}{1 - (A\eta_0/\delta_*)^2}.$$

This converts the infrared part of the mass-gap problem into measured/derived block constants. The continuum proof then rests on an equally explicit ultraviolet-to-block bridge and continuum Schwinger convergence estimates. Once those bridge lemmas are discharged, the OS and spectral-transfer sections give the required theory on $\mathbb{R}^4$ and

$$\text{Spec}(H) \cap (0, \Delta) = \emptyset, \quad \Delta > 0.$$

References

1. A. Jaffe and E. Witten, *Quantum Yang–Mills Theory*, in *The Millennium Prize Problems*, Clay Mathematics Institute / American Mathematical Society.
2. K. Osterwalder and R. Schrader, *Axioms for Euclidean Green’s Functions I*, *Communications in Mathematical Physics* 31 (1973).
3. K. Osterwalder and R. Schrader, *Axioms for Euclidean Green’s Functions II*, *Communications in Mathematical Physics* 42 (1975).
4. K. Osterwalder and E. Seiler, *Gauge Field Theories on a Lattice*, *Annals of Physics* 110 (1978), 440–471.
5. T. Balaban, *Renormalization Group Approach to Lattice Gauge Field Theories I: Generation of Effective Actions in a Small Field Approximation and a Coupling Constant Renormalization in Four Dimensions*, *Communications in Mathematical Physics* 109 (1987), 249–301.
6. T. Balaban, *Renormalization Group Approach to Lattice Gauge Field Theories II: Cluster Expansions*, *Communications in Mathematical Physics* 116 (1988), 1–22.
7. T. Balaban, *Convergent Renormalization Expansions for Lattice Gauge Theories*, *Communications in Mathematical Physics* 119 (1988), 243–285.
8. T. Balaban, *Large Field Renormalization I: The Basic Step of the R Operation*, *Communications in Mathematical Physics* 122 (1989), 175–202.
9. T. Balaban, *Large Field Renormalization II: Localization, Exponentiation, and Bounds for the R Operation*, *Communications in Mathematical Physics* 122 (1989), 355–392.
10. D. A. Yarotsky, *Ground States in Relatively Bounded Quantum Perturbations of Classical Lattice Systems*, *Communications in Mathematical Physics* 261 (2006), 799–819.
11. M. B. Hastings and T. Koma, *Spectral Gap and Exponential Decay of Correlations*, *Communications in Mathematical Physics* 265 (2006), 781–804.
12. Y. Guo, Z. Li, G. Yang, and G. Zhu, *Bootstrapping SU(3) Lattice Yang–Mills Theory*, *JHEP* 12 (2025) 033; arXiv:2502.14421.
13. R. F. Streater and A. S. Wightman, *PCT, Spin and Statistics, and All That*, Princeton University Press.
14. R. Haag, *Local Quantum Physics*, Springer.
15. H. Shen, R. Zhu, and X. Zhu, *A Stochastic Analysis Approach to Lattice Yang–Mills at Strong Coupling*, arXiv:2204.12737 (2022; revised 2026).