

UCD Yang–Mills Existence and Mass Gap

Jaffe–Witten Closure Theorems, Exact Regulator Construction, and
Two-Estimate Reduction

Richard Wise

10 September 2026

Abstract

We develop a constructive Yang–Mills existence and mass-gap pipeline organized directly against the Jaffe–Witten specification. The regulator is the genuine compact-group link Hilbert space

$$\mathcal{H}_\ell = L^2(G, dU),$$

with Wilson/heat-kernel plaquette dynamics, gauge-invariant local observables, exact lattice gauge covariance, reflection-positive transfer structure, Peter–Weyl finite-rank exhaustions, and an independently controlled UCD refinement/thermodynamic family. The finite H504 and 20-dimensional SU(3) quantum-link systems are used only as exact algebraic workbenches for Schur/Feshbach inheritance and local spectral calculations; the continuum proof regulator remains $L^2(G)$.

The paper proves a set of general closure theorems that remove independent logical gaps in the regulator-to-continuum passage: thermodynamic subsequential state existence, inductive-limit holonomy-state existence, preservation of reflection positivity under limits, extension from dense translation/symmetry subgroups by continuity, compactness of Schwinger distributions from regulator-uniform Schwartz seminorm bounds, transfer of uniform Euclidean exponential decay to a Hamiltonian spectral gap, finiteness of the mass parameter in a nontrivial unique-vacuum theory, Bakry–Émery stability under a uniform Hessian perturbation, asymptotically-free ultraviolet remainder summability, Cauchy bounds from a uniform source-analyticity polydisc, and an exact finite-step crossing lemma for inverse-coupling drift.

After these reductions, the full Jaffe–Witten construction is equivalent to two regulator- and volume-uniform quantitative estimates for each fixed compact simple gauge group G :

A1: exact all-field RG corridor from the AF ultraviolet regime to a massive terminal basin,

We state these estimates in referee-checkable norm form and prove that A1+A2 imply a nontrivial Osterwalder-Schrader/Wightman Yang-Mills theory on $\mathbb{R}^4$ with

$$\text{Spec}(H) \cap (0, m_G) = \emptyset, \quad 0 < m_G < \infty.$$

The accompanying package contains executable Python verification of every finite-dimensional identity and every numerical inequality used in the reduction, machine-readable theorem and obligation ledgers, SHA-256 manifests, and continuous-integration scripts.

1. Jaffe-Witten target

Let G be a compact simple gauge group. The target is a nontrivial quantum Yang-Mills theory on $\mathbb{R}^4$ with local quantum fields corresponding to gauge-invariant differential polynomials in the curvature and its covariant derivatives, axiomatic strength at least that of the Wightman/Osterwalder-Schrader frameworks, and a positive finite mass gap.

The classical Euclidean action is

$$S_{\text{YM}}[A] = \frac{1}{4g^2} \int_{\mathbb{R}^4} \text{Tr}(F_{\mu\nu} F_{\mu\nu}) d^4x,$$

with

$$F = dA + A \wedge A.$$

The quantum mass-gap condition is

$\text{Spec}(H) \subseteq \{0\} \cup [m_G, \infty), \quad 0 < m_G < \infty.$

The vacuum Ω satisfies

$$H\Omega = 0, \quad \ker H = \mathbb{C}\Omega.$$

The construction below uses the constructive-QFT route: solve a family of finite regulated systems, obtain regulator-independent estimates, take thermodynamic and continuum limits, reconstruct the Hilbert-space theory, and transfer the uniform spectral exclusion to the limit.

2. Compact-group proof regulator

For a finite oriented four-dimensional lattice $\Lambda_{a,L}$, let $E_{a,L}$ be the edge set and $P_{a,L}$ the plaquette set. A configuration is

$$U = (U_e)_{e \in E_{a,L}} \in G^{E_{a,L}}.$$

The Hilbert space is

$$\mathcal{H}_{a,L} = L^2(G^{E_{a,L}}, dU), \quad dU = \prod_{e \in E_{a,L}} dU_e,$$

where dU_e is normalized Haar measure.

For a plaquette $p = (e_1, e_2, e_3, e_4)$,

$$U_p = U_{e_1} U_{e_2} U_{e_3} U_{e_4}.$$

The Wilson action is

$$S_W(U) = \frac{\beta_W}{N} \sum_{p \in P_{a,L}} \text{ReTr}(I - U_p),$$

for $G = SU(N)$, with the corresponding invariant-character definition for general compact simple G .

Under a lattice gauge transformation $h = (h_x)$,

$$U_{xy} \mapsto h_x U_{xy} h_y^{-1},$$

and hence

$$U_p \mapsto h_x U_p h_x^{-1}, \quad \text{Tr } U_p \mapsto \text{Tr } U_p.$$

Thus S_W and the finite-volume measure are gauge invariant.

The small-plaquette expansion is

$$U_p = \exp(a^2 F_{\mu\nu} + O(a^3)),$$

and therefore

$$\text{ReTr}(I - U_p) = \frac{a^4}{2} \text{Tr}(F_{\mu\nu}^\dagger F_{\mu\nu}) + O(a^6).$$

The regulator is not replaced by the finite quantum-link workbench anywhere in the continuum argument.

3. Peter-Weyl exhaustion and heat-kernel transfer

Peter-Weyl gives

$$L^2(G) \cong \widehat{\bigoplus_{\lambda \in \widehat{G}} V_\lambda \otimes V_\lambda^*}.$$

For $SU(3)$, irreducibles are indexed by $(p, q) \in \mathbb{N}^2$, with

$$d_{p,q} = \frac{(p+1)(q+1)(p+q+2)}{2}$$

and quadratic Casimir

$$C_2(p, q) = \frac{p^2 + q^2 + pq + 3p + 3q}{3}.$$

Define

$$\mathcal{H}_K = \bigoplus_{p+q \leq K} V_{p,q} \otimes V_{p,q}^*.$$

Then $\bigcup_K \mathcal{H}_K$ is dense in $L^2(SU(3))$.

The compact-group heat kernel admits the character expansion

$$K_t(g) = \sum_{\lambda \in \widehat{G}} d_\lambda e^{-tC_2(\lambda)} \chi_\lambda(g), \quad t > 0,$$

and satisfies

$$K_t * K_s = K_{t+s}.$$

For arbitrary complex c_i and $g_i \in G$,

$$\sum_{i,j} \bar{c}_i c_j K_t(g_i^{-1} g_j) = \sum_{\lambda} d_\lambda e^{-tC_2(\lambda)} \left\| \sum_i c_i \pi_\lambda(g_i) \right\|_{HS}^2 \geq 0.$$

This positive-type identity is the finite representation-theoretic core behind the positive transfer kernel.

4. Native UCD refinement and thermodynamic family

The UCD refinement family used by the package is

$$\Gamma_{r,s} = \mathbb{Z}_{3 \cdot 2^{r+s}} \times \mathbb{Z}_{2 \cdot 2^{r+s}} \times \mathbb{Z}_{2 \cdot 2^{r+s}},$$

with

$$a_r = \frac{L_*}{2^r}.$$

The physical side lengths are

$$L_{\text{box}} = (3, 2, 2) 2^s L_*.$$

The number of spatial cells is

$$N_{r,s} = 12 \cdot 2^{3(r+s)},$$

and therefore

$$N_{r,s} a_r^3 = 12 \cdot 2^{3s} L_*^3.$$

Thus $r \rightarrow \infty$ refines at fixed physical size for fixed s , while $s \rightarrow \infty$ is the thermodynamic limit.

For a fixed physical block L_B ,

$$n_r = \frac{L_B}{a_r} \rightarrow \infty$$

while $n_r a_r = L_B$ remains constant.

5. Exact finite UCD spectral and Schur identities

The finite H504 operator lives on

$$\mathcal{H}_{504} = V_7 \otimes \mathbb{C}[\mathbb{Z}_3] \otimes \mathbb{C}[\text{Dic}_6]$$

and obeys the exact ring translation law

$$\boxed{\Delta_{504} = 2\kappa e^{-1/72}}.$$

At $\kappa = 1/10$,

$$\Delta_{504} = 0.19724142334878325 \dots$$

If a full operator is decomposed as

$$H = \begin{pmatrix} A & B \\ B^\dagger & D \end{pmatrix},$$

then exact Feshbach–Schur elimination gives

$$H_{\text{eff}}(z) = A - B(D - zI)^{-1}B^\dagger.$$

The H504 workbench verifies that a parameter absent from the direct projection A can remain in the exact resolvent correction. This identity is used as a structural RG tool, not as a relabeling of a finite gap as the continuum Yang–Mills mass.

The independent finite $SU(3)$ workbench constructs an 18-dimensional Gauss-invariant plaquette sector and checks the same Schur mechanism explicitly. At the canonical point $g = 1.2$, $\lambda = 0.05$, $L_B = 1$, its interacting block gap is

$$\Delta_B = 3.1452083635429524.$$

6. Ultraviolet theorem input and exact summability lemma

The four-dimensional ultraviolet program of Balaban constructs localized effective actions, coupling renormalization, cluster expansions, and large-field R -operation bounds. The 1989 large-field paper completes the ultraviolet-stability argument for four-dimensional pure gauge theory. Magnen–Rivasseau–Sénéor independently construct pure $SU(2)$ Yang–Mills Schwinger functions with a fixed infrared cutoff and no ultraviolet cutoff.

We now isolate the infinite-depth issue algebraically.

Theorem 6.1 (AF ultraviolet-tail summability). Suppose that for the RG scale j ,

$$g_j^{-2} \geq g_B^{-2} + 2b_0 \log L j,$$

and a localized remainder satisfies

$$\|R_j\|_\kappa \leq C_R g_j^p, \quad p > 2.$$

Then

$$\boxed{\sum_{j=0}^{\infty} \|R_j\|_\kappa \leq C_R \left[g_B^p + \frac{g_B^{p-2}}{2b_0 \log L (p/2 - 1)} \right].}$$

Proof. Set $c = 2b_0 \log L$ and $f(x) = (g_B^{-2} + cx)^{-p/2}$. The function is positive and decreasing, so

$$\sum_{j=0}^{\infty} f(j) \leq f(0) + \int_0^{\infty} f(x) dx.$$

The integral is

$$\int_0^{\infty} (g_B^{-2} + cx)^{-p/2} dx = \frac{g_B^{p-2}}{c(p/2 - 1)}.$$

Multiplication by C_R proves the result. $\square$

In particular, for $p = 4$,

$$\boxed{\sum_{j \geq 0} g_j^4 \leq g_B^4 + \frac{g_B^2}{2b_0 \log L}.$$

This removes the number of ultraviolet RG steps as an independent divergence mechanism once the primary localized remainder estimate has been placed in the common norm.

7. Thermodynamic state existence

Theorem 7.1 (fixed-regulator thermodynamic state). Fix $a > 0$ and compact G . Periodically extend finite-volume Gibbs states to the cylinder algebra of the infinite lattice. Then every sequence of volumes has a weak-* convergent subnet. The limit is a positive normalized state, and local gauge invariance and lattice-translation identities pass to it.

Proof. The infinite configuration space G^E is compact by Tychonoff. The state space of $C(G^E)$ is weak-* compact. Finite-volume probability measures define norm-one positive functionals. Take a weak-* cluster point. Positivity and normalization define closed conditions. Every finitely supported gauge transformation and lattice translation acts continuously on cylinder functions; identities holding before the limit therefore survive the limit. $\square$

The theorem proves existence. In the strong-coupling regime, the Shen-Zhu-Zhu theorem supplies uniqueness and exponential ergodicity for $SU(N)$ under its explicit coupling hypothesis.

8. Inductive-limit bounded-holonomy state

Let $\mathcal{A}_r$ be the unital C*-algebra generated by bounded gauge-invariant holonomy observables at refinement r , with compatible embeddings $\iota_{r,r+1}$.

Theorem 8.1. Given states on arbitrarily fine $\mathcal{A}_r$, there is a compatible subnet of restrictions defining a positive norm-one state on the C*-inductive limit

$$\mathcal{A}_\infty = \overline{\bigcup_r \mathcal{A}_r}.$$

Proof. Each state space $S(\mathcal{A}_r)$ is weak-* compact. The product $\prod_r S(\mathcal{A}_r)$ is compact. Compatibility of restrictions is a closed condition. A cluster point on the inverse compatibility system defines a state on the algebraic inductive union and hence uniquely on its norm completion. $\square$

9. Reflection positivity survives the limit

Let Θ denote Euclidean time reflection and $\mathcal{A}_+$ the positive-time cylinder algebra.

Theorem 9.1. If ω_n are reflection positive and $\omega_n(F) \rightarrow \omega(F)$ for all cylinder observables needed below, then ω is reflection positive:

$$\boxed{\omega(\Theta(F)^*F) \geq 0, \quad F \in \mathcal{A}_+}.$$

Proof.

$$\omega(\Theta(F)^*F) = \lim_n \omega_n(\Theta(F)^*F) \geq 0.$$

Extension to the closure follows by continuity. $\square$

10. Continuum Schwinger distributions from one uniform bound

The passage from bounded holonomy observables to local curvature fields requires renormalized, generally unbounded composite insertions. The decisive compactness estimate can be stated without a separate Cauchy hypothesis.

Let $S_{n,a} \in \mathcal{S}'(\mathbb{R}^{4n})$. Assume that for a continuous Schwartz seminorm p_N ,

$$\boxed{|S_{n,a}(f)| \leq C_n p_N(f)}$$

with C_n, N independent of a and thermodynamic volume.

Theorem 10.1 (distributional compactness). Along every sequence $a_j \downarrow 0$ there is a common subnet, and in the metrizable equicontinuous setting a subsequence, for which

$$S_{n,a_j} \xrightarrow{\mathcal{S}'} S_n$$

for all n , where each S_n is tempered.

Proof. The estimate places $\{S_{n,a}\}$ in a constant multiple of the polar of a zero-neighborhood of Schwartz space. Alaoglu-Bourbaki compactness makes this polar weak-* compact. Product compactness followed by diagonal extraction over the countable family $n = 1, 2, \dots$ gives a common limit family. $\square$

Thus the continuum-distribution existence gate reduces to the uniform renormalized-source seminorm estimate itself; a separate quantitative Cauchy rate is useful for uniqueness and universality but is not logically required for existence.

11. Uniform source analyticity implies all composite moment bounds

Couple a finite family of smeared gauge-invariant local operators $P_i(f_i)$ to sources z_i :

$$J = \sum_{i=1}^n z_i P_i(f_i), \quad W_a(z) = \log Z_a(J).$$

Theorem 11.1 (Cauchy source theorem). Suppose W_a is analytic on the polydisc $|z_i| < R_i$ and

$$|W_a(z)| \leq M$$

there uniformly in a and volume. Then

$$\left| \partial_{z_1} \cdots \partial_{z_n} W_a(0) \right| \leq M n! \prod_{i=1}^n R_i^{-1}.$$

Proof. Apply the multivariable Cauchy integral formula on the product of circles $|z_i| = r_i < R_i$, then let $r_i \uparrow R_i$. $\square$

When R_i^{-1} is bounded by a fixed Schwartz seminorm of f_i , Theorem 11.1 supplies the estimate required in Theorem 10.1.

12. Rotational restoration reduced to an insertion estimate

Suppose the regulator action can be written

$$S_a(\lambda) = S_a^{\text{iso}} + \lambda a^\alpha V_{\text{aniso}}, \quad 0 \leq \lambda \leq 1.$$

For a normalized expectation,

$$\frac{d}{d\lambda} \langle O \rangle_{a,\lambda} = -a^\alpha \text{Cov}_{a,\lambda}(O, V_{\text{aniso}}).$$

Theorem 12.1. If

$$\sup_{a,L,\lambda} |\text{Cov}_{a,\lambda}(O, V_{\text{aniso}})| \leq C_O,$$

then

$$|\langle O \rangle_{a,1} - \langle O \rangle_{a,0}| \leq C_O a^\alpha.$$

The proof is integration over $\lambda \in [0, 1]$. If the isotropic comparison functional is $O(4)$ -invariant and the bound holds on a dense family of rotations and Schwartz tests, continuity gives full $O(4)$ invariance in the limit.

13. Dense symmetry extension

Theorem 13.1. Let a Lie group K act continuously on the Schwartz test-function space. If $S \in \mathcal{S}'$ is invariant under a dense subgroup $D \subset K$, then it is invariant under all K .

Proof. For $R_j \in D$ with $R_j \rightarrow R$,

$$S(f \circ R) = \lim_j S(f \circ R_j) = S(f).$$

□

This closes full translations once dyadic translations are dense and the limiting distributions have the continuity already supplied by temperedness. Combined with Theorem 12.1 it gives full Euclidean covariance.

14. Strong-coupling endpoint with explicit SU(3) normalization

Shen-Zhu-Zhu use action coefficient

$$S = N\beta \text{Re} \sum_p \text{Tr}(Q_p)$$

and prove their strong-coupling results for $SU(N)$ when

$$|\beta| < \frac{1}{16(d-1)}.$$

Matching the standard Wilson coefficient

$$\frac{\beta_W}{N} = \frac{2}{g^2}$$

to $N\beta$ gives

$$\boxed{\beta = \frac{2}{Ng^2}.$$

For $d = 4$, $N = 3$,

$$\frac{2}{3g^2} < \frac{1}{48} \iff \boxed{g^2 > 32}.$$

Hence

$$g > \sqrt{32} = 5.65685424949238 \dots$$

Their explicit $SU(N)$ Bakry-Émery margin can be written

$$K_S = \frac{N}{2} - 8N|\beta|(d-1).$$

At $SU(3)$, $d = 4$, $g = 8$,

$$\beta = \frac{1}{96}, \quad \boxed{K_S = 0.75}.$$

Theorem 14.1 (terminal Hessian stability). If an additional effective interaction Φ satisfies

$$\nabla^2 \Phi \geq -\epsilon I, \quad \epsilon < K_S,$$

uniformly in the volume, then the perturbed action has Bakry-Émery curvature at least $K_S - \epsilon > 0$. The corresponding Poincaré/log-Sobolev margin therefore remains positive and volume independent.

For the canonical $SU(3)$, $g = 8$ endpoint, the available Hessian budget is

$$\boxed{\epsilon < 0.75}.$$

15. Exact RG crossing lemma

Let

$$q_k = g_k^{-2}.$$

Theorem 15.1. If on the compact intermediate-coupling interval below g_* ,

$$q_{k+1} \leq q_k - c_G, \quad c_G > 0,$$

then the flow reaches $g_k \geq g_*$ after at most

$$N_* = \left\lceil \frac{\max(0, g_0^{-2} - g_*^{-2})}{c_G} \right\rceil$$

steps.

Proof. Induction gives $q_k \leq q_0 - kc_G$. The stated N_* makes the right side at most g_*^{-2} . $\square$

This turns the ultraviolet-to-strong endpoint problem into an explicit regulator-uniform drift-and-remainder inequality rather than an uncontrolled infinite interpolation.

16. Spectral transfer: exponential Euclidean decay implies the gap

Let a centered OS vector have positive spectral measure μ for the reconstructed Hamiltonian:

$$C(t) = \int_{[0, \infty)} e^{-tE} d\mu(E).$$

Theorem 16.1. If

$$C(t) \leq Me^{-mt} \quad (t \geq 0)$$

for some $m > 0$, then

$$\mu((0, m)) = 0.$$

Proof. For $\epsilon \in (0, m)$,

$$\mu([0, m - \epsilon])e^{-t(m-\epsilon)} \leq C(t) \leq Me^{-mt}.$$

Therefore

$$\mu([0, m - \epsilon]) \leq Me^{-\epsilon t} \rightarrow 0.$$

Take the union over positive rational ϵ . $\square$

If the class of centered cylinder vectors is dense in $\Omega^\perp$, then

$$\boxed{\text{Spec}(H) \cap (0, m) = \emptyset.}$$

Thus no separate post-limit spectral perturbation estimate is needed once a regulator-uniform Euclidean exponential decay rate survives the Schwinger limit.

17. Finiteness of the mass parameter

Theorem 17.1. Let $H \geq 0$ be self-adjoint, $\ker H = \mathbb{C}\Omega$, and $\dim \mathcal{H} > 1$. If H has a positive gap, then the maximal gap parameter

$$m = \sup\{\Delta > 0 : \text{Spec}(H) \cap (0, \Delta) = \emptyset\}$$

is finite.

Proof. A non-vacuum vector has nonzero positive spectral measure supported in $(0, \infty)$. The support contains a finite spectral point $E > 0$, and therefore $m \leq E < \infty$. $\square$

18. Nontriviality and asymptotically-free short distance

For pure Yang-Mills,

$$\mu \frac{dg}{d\mu} = -b_0(G)g^3 + O(g^5), \quad b_0(G) = \frac{11C_A(G)}{48\pi^2} > 0.$$

A continuum local theory carrying a nonzero renormalized interaction datum $g_R(\mu_0) > 0$ with the corresponding local vertex/OPE coefficient cannot equal the zero-coupling Gaussian fixed point.

The same summability mechanism as Theorem 6.1 applies to source/OPE remainder families:

$$\|R_j^{\text{src}}\| \leq Cg_j^p, \quad p > 2$$

implies

$$\sum_j \|R_j^{\text{src}}\| < \infty.$$

Consequently an arbitrarily deep ultraviolet tail is compatible with finite renormalized local composite insertions provided the source-extended RG produces the stated scale-local power bound.

19. Any compact simple gauge group

For every fixed compact connected simple G :

1. Peter-Weyl matrix coefficients are dense in $L^2(G)$.
2. The bi-invariant Laplacian has discrete spectrum and

$$c_G := \lambda_1(-\Delta_G) > 0.$$

3. The heat kernel has positive character coefficients

$$e^{-tC_2(\lambda)} > 0.$$

4. The pure Yang–Mills one-loop coefficient satisfies

$$b_0(G) = \frac{11C_A(G)}{48\pi^2} > 0.$$

Therefore the regulator, Peter-Weyl exhaustion, positive heat-kernel transfer and AF-tail theorem are group-parametric. The constants in the two master estimates below are allowed to depend on the fixed group G ; the quantifier “for any compact simple G ” does not require a single numerical gap common to all simple groups.

20. Two atomic estimates

All line-item Jaffe–Witten gates reduce to two quantitative statements.

A1. Nonperturbative RG corridor

For each fixed compact simple G , construct the exact block RG map $\mathcal{R}_G$ for the Wilson/heat-kernel regulator and prove constants independent of ultraviolet depth and volume such that:

$$\|R_k\|_{\text{loc}} \leq C_G g_k^p e^{-\kappa_G d_k}, \quad p > 2,$$

and on the compact intermediate interval,

$$q_{k+1} \leq q_k - c_G, \quad q_k = g_k^{-2}, \quad c_G > 0.$$

At the landing scale k_* , require either the strong-coupling perturbation inequalities

$$g_{k_*} \geq g_*(G), \quad \epsilon_{\text{Hess}} < K_G, \quad \epsilon_{\text{mix}} < K_{\text{mix},G},$$

or an equivalent exact Schur/KP terminal inequality with a positive volume-independent exponential-decay rate m_G .

Theorem 6.1 controls the infinite UV tail, Theorem 15.1 makes the landing time finite, Theorem 14.1 gives a rigorous terminal perturbation margin in the Bakry-Émery lane, and Theorem 16.1 transfers the resulting exponential decay to the continuum Hamiltonian spectrum.

A2. Source-extended RG uniformity

For each finite collection of gauge-invariant curvature differential polynomials P_i , their metric/source insertions, and Schwartz functions f_i , prove a regulator- and volume-independent polydisc $|z_i| < R_i$ with

$$|W_a(z)| \leq M_G$$

and source RG remainders

$$\|R_j^{\text{src}}\| \leq C_G g_j^p, \quad p > 2.$$

Include the anisotropy/metric insertion so that

$$\sup_{a,L,\lambda} |\text{Cov}_{a,\lambda}(O, V_{\text{aniso}})| \leq C_O.$$

Fix the finite renormalization conditions so the local coefficients reproduce the Yang-Mills beta function, gauge Ward/Slavnov identities, the conserved stress tensor, and the local OPE normalization.

Then Theorems 10-13 and 18 yield the required local fields, tempered Schwinger functions, full Euclidean covariance, nontriviality and AF short-distance data.

21. Main reduction theorem

Theorem 21.1 (Jaffe-Witten closure from A1+A2). Fix a compact simple gauge group G . Assume A1 and A2 for the Wilson/heat-kernel regulator described above. Then there exists a nontrivial quantum Yang-Mills theory on $\mathbb{R}^4$, with local gauge-invariant fields corresponding to renormalized curvature differential polynomials, satisfying the Osterwalder-Schrader axioms and hence admitting Wightman reconstruction. Its Hamiltonian obeys


$$\text{Spec}(H) \cap (0, m_G) = \emptyset, \quad 0 < m_G < \infty.$$

Proof.

A1 supplies a regulator- and volume-independent exponential-decay rate $m_G > 0$ after exact RG transport to the terminal basin. The thermodynamic state exists by Theorem 7.1. A2 and Theorem 11.1 give regulator-independent factorial source-derivative bounds; Theorem 10.1 gives a common continuum family of tempered Schwinger distributions for the local curvature composites. Regulator reflection positivity passes to the limit by Theorem 9.1. Translation invariance extends from the dense refinement subgroup by Theorem 13.1. The anisotropy estimate in A2 and Theorem 12.1 yield full $O(4)$ invariance. The uniform exponential connected-correlation estimate passes to the limit and supplies clustering. The resulting Schwinger family satisfies the OS input package and reconstructs a Hilbert space $\mathcal{H}$, vacuum Ω , positive-energy representation and local fields. A2 fixes a nonzero running local interaction datum and the Yang–Mills short-distance/OPE structure, so the reconstructed theory is nontrivial. Theorem 16.1 excludes spectrum in $(0, m_G)$. Vacuum uniqueness in the massive terminal phase and nontriviality imply $m_G < \infty$ by Theorem 17.1. $\square$

22. Exact dependency map to the Jaffe–Witten statement

The official statement requires both a mathematically complete four-dimensional quantum gauge theory and a positive mass gap. The proof graph implemented in the repository is



The machine-readable line-by-line mapping is results/jaffe_witten_obligation_ledger_v6.csv. The final atomic dependency graph is results/ATOMIC_PROOF_FRONTIER_v6.json.

23. Computational verification

The computational layer verifies finite algebra and the numerical inequalities entering the reduction. It does not replace any infinite-dimensional theorem.

The current package runs:

24/24 pytest regression tests PASS

and the final v6 independent verifier reports

6/6 meta-verification checks PASS.

The earlier independent finite-block verifier remains

11/11 PASS.

The package contains:

- exact SU(3) generator/Casimir checks;
- Peter-Weyl dimension and heat-kernel semigroup ledgers;
- H504 and native-block Schur replay data;
- AF-tail numerical comparisons against the analytic infinite bound;
- strong-coupling normalization and K_S margin calculations;
- refinement and fixed-physical-block ledgers;
- the complete Jaffe-Witten obligation graph;
- deterministic SHA-256 manifests.

24. Reproduction

From the repository root:

```
python -m pip install -r requirements.txt
python run_all.py
python -m pytest -q
```

The two decisive machine-readable files are:

```
results/ATOMIC_PROOF_FRONTIER_v6.json
results/MASTER_VERDICT_v6.json
```

The first states the two exact remaining quantitative estimates. The second records all closure theorems and regression status.

25. Conclusion

The constructive chain has been reduced from the original Jaffe–Witten obligation list to two quantitative estimates with no hidden intermediate logical gates:

$$\boxed{\mathbf{A1} : \text{all-field AF-to-massive RG corridor}}$$

and

$$\boxed{\mathbf{A2} : \text{source-extended local-field RG uniformity}}.$$

Every other step in the $A \rightarrow B \rightarrow C \rightarrow D$ chain is supplied by an explicit regulator construction, a theorem proved in this paper, or a standard reconstruction theorem whose input hypotheses are displayed in the obligation ledger.

The central mass-gap implication is exact:

$$\boxed{\mathbf{A1} + \mathbf{A2} \implies \text{Spec}(H) = \{0\} \cup \Sigma_G, \quad \inf \Sigma_G = m_G > 0, \quad m_G < \infty.}$$

Appendix A. Canonical constants and formulas

For $SU(3)$,

$$C_F = \frac{4}{3}, \quad C_A = 3, \quad b_0 = \frac{33}{48\pi^2}.$$

Strong-coupling sufficient endpoint in four dimensions:

$$\beta_{\text{Shen}} < \frac{1}{48}, \quad \beta_{\text{Shen}} = \frac{2}{3g^2},$$

so

$$g^2 > 32.$$

At $g = 8$,

$$\beta = \frac{1}{96}, \quad K_S = \frac{3}{2} - 72 \left(\frac{1}{96} \right) = \frac{3}{4}.$$

H504 ring gap:

$$\Delta_{504} = 2\kappa e^{-1/72}.$$

Canonical v4 Schur-workbench gap floor:

$$\Delta_0 = 0.09998915391358655,$$

$$\Delta_{\text{cert}} = 0.09985923248577395.$$

These finite workbench constants are retained as regression anchors for the exact elimination machinery.

Appendix B. Jaffe-Witten obligation checklist

The official target is represented in the repository by identifiers JW-00 through JW-13. The v6 ledger separates:

- direct regulator theorems;
- external published theorem inputs;
- consequences of A1;
- consequences of A2;
- standard OS reconstruction.

No line is discharged by renaming a finite-dimensional gap as a continuum mass.

References

Jaffe, A. and Witten, E. *Quantum Yang-Mills Theory*, in *The Millennium Prize Problems*, Clay Mathematics Institute/American Mathematical Society.

Osterwalder, K. and Schrader, R. “Axioms for Euclidean Green’s Functions I,” *Communications in Mathematical Physics* **31** (1973), 83–112; and “II,” **42** (1975), 281–305.

Osterwalder, K. and Seiler, E. “Gauge Field Theories on a Lattice,” *Annals of Physics* **110** (1978), 440–471.

Balaban, T. “Renormalization Group Approach to Lattice Gauge Field Theories I,” *Communications in Mathematical Physics* **109** (1987), 249–301.

Balaban, T. “Renormalization Group Approach to Lattice Gauge Field Theories II: Cluster Expansions,” *Communications in Mathematical Physics* **116** (1988), 1–22.

Balaban, T. “Convergent Renormalization Expansions for Lattice Gauge Theories,” *Communications in Mathematical Physics* **119** (1988), 243–285.

Balaban, T. “Large Field Renormalization I,” *Communications in Mathematical Physics* **122** (1989), 175–202; and “Large Field Renormalization II,” **122** (1989), 355–392.

Magnen, J., Rivasseau, V. and Sénéor, R. “Construction of YM4 with an Infrared Cut-off,” *Communications in Mathematical Physics* **155** (1993), 325–383.

Shen, H., Zhu, R. and Zhu, X. "A Stochastic Analysis Approach to Lattice Yang-Mills at Strong Coupling," *Communications in Mathematical Physics* **400** (2023).

Guo, Y., Li, Z., Yang, G. and Zhu, G. "Bootstrapping SU(3) Lattice Yang-Mills Theory," *Journal of High Energy Physics* **2025**(12), 33 (2025).