

Adelic Synthesis: The Pattern-Particle Correspondence and the Complete Arithmetic Theory of Anyons

Contents

Adelic Synthesis: The Pattern-Particle Correspondence and the Complete Arithmetic Theory of Anyons	2
Abstract	2
1. Introduction: From Places to the Adelic Whole	2
1.1 Ostrowski's Theorem and the Incompleteness of Archimedean Anyon Theory	2
1.2 The Adelic Perspective: A Brief Primer	3
1.3 The Pattern-Particle Correspondence (Informal Statement)	3
2. The Adelic Braid Group	4
2.1 Building on Phase 1	4
2.2 Restricted Product of Braid Groups	4
2.3 Adelic Configuration Space	5
3. The Adelic Temperley-Lieb Algebra and Jones Polynomial	5
3.1 Building on Phase 2	5
3.2 Adelic Temperley-Lieb Algebra	5
4. Adelic Anyon Fusion via Quantum Groups	6
4.1 Building on Phase 3	6
4.2 The Adelic Quantum Group	7
4.3 Adelic Anyon Types	7
5. The Adelic Verlinde Algebra and Fusion Category	8
5.1 Adelic Fusion Rules	8
5.2 Adelic Fusion Category	8
5.3 Adelic Braiding: Place-Crossing Phenomena	9
6. Physical Implications	9
6.1 p-Adic Measurements and the Ultrametric Hierarchy	9
6.2 Topological Entanglement Entropy: Adelic Contributions	10
6.3 The p-Adic Hierarchy and Gate Compilation	10
6.4 The Complete Adelic Anyon Table	10
7. Adelic Topological Quantum Computation (ATQC)	11
7.1 The ATQC Architecture	11
7.2 Place-Multiplexed Gate Set	11
7.3 Adelic CNOT via Place Crossing	11
8. Open Questions and Future Directions	12
8.1 Experimental Realization	12
8.2 Adelic Chern-Simons Theory	12
8.3 Langlands Program and Anyons	12
9. Conclusion: The Complete Picture	13
References	13

Adelic Synthesis: The Pattern-Particle Correspondence and the Complete Arithmetic Theory of Anyons

Author: Rowan Brad Quni-Gudzinis | **Date:** 2026-07-05 | **Version:** v1.0 **License:** QNFO Unified License Agreement (QNFO-ULA): <https://legal.qnfo.org/> **Project:** QLoF Extension — Program D: p-adic Anyons (Phase 4 — Adelic Synthesis) **Cross-References:** Phase 1 [DOI: 10.5281/zenodo.21208366] | Phase 2 [DOI: 10.5281/zenodo.21208368] | Phase 3 [DOI: 10.5281/zenodo.21208491] | Research Plan [DOI: 10.5281/zenodo.21208370]

Abstract

The prior three phases of this program constructed p-adic anyon theories at individual finite places: p-adic braid groups on Bruhat-Tits buildings (Phase 1), the Temperley-Lieb parameter as p-adic cyclotomic units with the p-adic Jones polynomial (Phase 2), and anyon fusion and braiding via restricted quantum groups at roots of unity over p-adic fields (Phase 3). Each phase treated a single non-archimedean place in isolation. This capstone paper constructs the **adelic synthesis**: the unified framework that treats an “anyon” as an adelic object — a single arithmetic pattern that manifests differently at each completion of $\mathbb{Q}$, with the familiar archimedean Fibonacci anyon being merely the ∞ -place avatar of a richer adelic entity.

We construct the **adelic braid group** $\mathbb{B}_n(\mathbb{A})$ on the adèle ring $\mathbb{A}$, the **adelic Temperley-Lieb algebra** $\mathbb{TL}(\mathbb{A})$, and the **adelic anyon fusion category** $\mathcal{F}(\mathbb{A})$, proving that all three are restricted products over all places of $\mathbb{Q}$ of their place-specific counterparts. The central result — termed the **Pattern-Particle Correspondence** — states that an anyon type is an adelic representation of $U_q(\mathfrak{sl}_2)$ evaluated at the cyclotomic place $q = \zeta_{p^k}$ for each p , together with the archimedean specialization $q = e^{i\pi/(k+2)}$. The adelic Verlinde algebra factorizes as a restricted tensor product: $\mathcal{V}(\mathbb{A}) \cong \bigotimes_p' \mathcal{V}(\mathbb{Q}_p) \otimes \mathcal{V}(\mathbb{R})$.

Physical implications are profound: the adelic framework predicts that the “same” anyon type carries distinct observable signatures at different places — topological entanglement entropy, braid phases, and fusion multiplicities all acquire p-adic filtration structures absent from the purely archimedean theory. We propose the **Adelic Topological Quantum Computation (ATQC)** paradigm in which computational gates exploit place-crossing transitions (adelic Hecke operators) rather than continuous braiding, potentially eliminating the Solovay-Kitaev overhead entirely.

1. Introduction: From Places to the Adelic Whole

1.1 Ostrowski’s Theorem and the Incompleteness of Archimedean Anyon Theory

Ostrowski’s theorem [Ostrowski1918] establishes that every non-trivial absolute value on $\mathbb{Q}$ is equivalent to either the standard archimedean absolute value $|\cdot|_\infty$ or a p-adic absolute value $|\cdot|_p$ for some prime p . This theorem partitions the world of number theory into infinitely many “places” — one archimedean and one for each prime.

The standard theory of anyons [Kitaev2003; Nayak2008] — including the Chern-Simons effective field theory, the Jones polynomial via the Kauffman bracket, and the Fibonacci anyon model — is constructed entirely at the archimedean place. Every Hilbert space is complex, every path integral is over real manifolds, and every braid is a continuous path in $\mathbb{R}^3$. Ostrowski’s theorem implies that this construction is **incomplete**: it captures only one of infinitely many equally fundamental completions of $\mathbb{Q}$.

Prior work in this program has rectified this incompleteness, one place at a time:

Phase	Construction	Place
Phase 1 [QuniGudzi- nas2026a]	p-adic braid group $B_n(\mathbb{Q}_p)$ on Bruhat-Tits building	Arbitrary p
Phase 2 [QuniGudzi- nas2026b]	p-adic Temperley-Lieb parameter $\delta_p \in \mathbb{Z}_p[\zeta_{p^k}]$; p-adic Jones polynomial	Arbitrary p
Phase 3 [QuniGudzi- nas2026c]	p-adic anyon models via $U_q(\mathfrak{sl}_2)$ at roots of unity; fusion and braiding	Arbitrary p

The remaining task — this paper — is to **unify** these place-specific constructions into a single adelic entity. The question is not merely “what happens at each place?” but “what is the complete object whose projections are these place-specific avatars?”

1.2 The Adelic Perspective: A Brief Primer

The adele ring $\mathbb{A}$ of $\mathbb{Q}$ is the restricted direct product of all completions of $\mathbb{Q}$:

$$\mathbb{A} = \mathbb{R} \times \prod_p' \mathbb{Q}_p$$

where the restricted product means that for all but finitely many primes p , the component lies in the ring of integers $\mathbb{Z}_p$. An adele is thus a tuple $x = (x_\infty, x_2, x_3, x_5, \dots)$ with $x_p \in \mathbb{Z}_p$ for almost all p .

The idele group $\mathbb{A}^\times$ is the restricted product of the multiplicative groups $\mathbb{Q}_p^\times$ together with $\mathbb{R}^\times$. The key structural feature is the **diagonal embedding** $\mathbb{Q} \hookrightarrow \mathbb{A}$ sending $q \mapsto (q, q, q, \dots)$, which satisfies the **product formula**:

$$\prod_v |q|_v = 1, \quad \forall q \in \mathbb{Q}^\times$$

where the product runs over all places $v \in \{\infty, 2, 3, 5, \dots\}$. This product formula is the arithmetic identity that forces the adelic point of view to be the natural framework for any construction over $\mathbb{Q}$.

The crucial insight of this paper is that **anyons are arithmetic objects defined over $\mathbb{Q}$** , and therefore their natural habitat is the adele ring, not $\mathbb{R}$ alone.

1.3 The Pattern-Particle Correspondence (Informal Statement)

Central Thesis: An “anyon” is not a particle per se, but an *adelic pattern* — an irreducible representation of a quantum group defined over $\mathbb{Q}$, evaluated at each completion simultaneously. The archimedean Fibonacci anyon ($\tau = \frac{1+\sqrt{5}}{2}$) is the ∞ -place avatar; the p-adic Fibonacci anyon (Phase 3) is the p -place avatar. They are the same arithmetic object, viewed through different absolute values.

This perspective resolves a tension in the standard theory: why does the fusion category for $SU(2)_k$ Chern-Simons theory require $q = e^{i\pi/(k+2)}$ — a complex root of unity? The answer from the adelic perspective is that q is not “a complex number” but a **cyclotomic integer** living in $\bar{\mathbb{Q}} \subset \bar{\mathbb{Q}}_p$ for every p , and its complex evaluation is just its image under the unique archimedean embedding. The full pattern includes the p-adic evaluations as well.

2. The Adelic Braid Group

2.1 Building on Phase 1

Phase 1 [QuniGudzin2026a] constructed the p -adic braid group $B_n(\mathbb{Q}_p)$ as the fundamental group of the configuration space of n distinct vertices in the Bruhat-Tits tree $\mathcal{T}_p$ (for SL_2) or more generally on the building $\mathcal{B}(\mathrm{SL}_n, \mathbb{Q}_p)$. Key results:

1. **Geometric realization:** Braid generators σ_i correspond to geodesic swaps of marked vertices on $\mathcal{T}_p$. The $(p+1)$ -regular tree structure makes braiding inherently discrete — there is no continuous deformation of braids.
2. **Bruhat-Tits presentation:** The generators satisfy the standard braid relations $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ and $\sigma_i \sigma_j = \sigma_j \sigma_i$ for $|i-j| > 1$, but acquire additional relations from the apartment structure: any word representing a closed geodesic in an apartment acts as the identity on that apartment level.
3. **Archimedean limit:** $\lim_{p \rightarrow \infty} B_n(\mathbb{Q}_p) \cong B_n$ (the classical braid group), where the limit is taken in the sense of the generic fiber over $\mathbb{C}_p \cong \mathbb{C}$.

2.2 Restricted Product of Braid Groups

We now define the **adelic braid group** as the restricted product of place-specific braid groups:

$$\mathbb{B}_n(\mathbb{A}) = B_n(\mathbb{R}) \times \prod_p' B_n(\mathbb{Q}_p)$$

where the restricted product condition means: for all but finitely many primes p , the p -component braid word is **unramified** — i.e., it corresponds to the unique geodesic path of length zero in the subtree of $\mathcal{T}_p$ where all marked vertices lie in a single apartment. Equivalently, the braid is trivial on the mod- p building reduction for almost all p .

Construction 2.1 (Adelic braid word): An element of $\mathbb{B}_n(\mathbb{A})$ is a tuple $\mathbf{b} = (b_\infty, \{b_p\}_p)$ where: - $b_\infty \in B_n(\mathbb{R})$ is a classical braid word - $b_p \in B_n(\mathbb{Q}_p)$ is a p -adic braid word on $\mathcal{T}_p$ - For almost all p , $b_p = \mathrm{id}$ (the trivial braid)

Theorem 2.2 (Diagonal embedding of $B_n(\mathbb{Q})$): There exists a canonical injection

$$\iota : B_n(\mathbb{Q}) \hookrightarrow \mathbb{B}_n(\mathbb{A})$$

defined by sending a braid word with coefficients in $\mathbb{Q}$ to its image under each completion. Concretely, if a braid is defined by algebraic equations over $\mathbb{Q}$, its solutions embed diagonally into all completions simultaneously.

Proof. Since the braid group B_n is finitely presented, any braid word with $\mathbb{Q}$ -coefficients defines, via each place's completion, a braid in $B_n(K_v)$ for every place v . The condition that almost all components are unramified follows from the fact that a $\mathbb{Q}$ -defined braid has denominators in only finitely many primes, so its image in $\mathbb{Q}_p$ lies in $\mathbb{Z}_p$ for almost all p , hence lives in the maximal compact subgroup of the Bruhat-Tits building automorphism group — i.e., the trivial braid at the level of mod- p reduction for all but finitely many p . $\square$

Theorem 2.3 (Strong Approximation for Adelic Braids): The image of ι is dense in $\mathbb{B}_n(\mathbb{A})$ in the adelic topology.

This is the adelic analog of the standard strong approximation theorem for SL_n . It means that any adelic braid can be approximated arbitrarily well (at finitely many places simultaneously) by a $\mathbb{Q}$ -rational braid.

2.3 Adelic Configuration Space

The classical braid group is $B_n = \pi_1(\mathrm{Conf}_n(\mathbb{C}))$. The adelic analog requires an adelic configuration space.

Definition 2.4 (Adelic configuration space): For each place v , let X_v be the configuration space of n distinct points in the geometric realization at place v : - $X_\infty = \mathrm{Conf}_n(\mathbb{C})$ (classical) - $X_p = \mathrm{Conf}_n(\mathcal{T}_p)$ (vertices of the Bruhat-Tits tree, Phase 1)

The adelic configuration space is the restricted product:

$$\mathbb{X}_n = X_\infty \times \prod'_p X_p$$

Theorem 2.5 (Adelic fundamental group): $\mathbb{B}_n(\mathbb{A}) \cong \pi_1^{\mathrm{\acute{e}t}}(\mathbb{X}_n)$, the étale fundamental group of the adelic configuration space.

3. The Adelic Temperley-Lieb Algebra and Jones Polynomial

3.1 Building on Phase 2

Phase 2 [QuniGudzin2026b] established that the Temperley-Lieb algebra parameter $\delta = -A^2 - A^{-2}$ at a p -adic place takes values in the cyclotomic units $\mathbb{Z}_p[\zeta_{p^k}]^\times$. The Jones polynomial $V_L(t)$ at a p -adic place is then a p -adic Laurent polynomial:

$$V_L^{(p)}(t) \in \mathbb{Z}_p[\zeta_{p^k}][t, t^{-1}]$$

Key results from Phase 2: 1. **Cyclotomic identification:** $A_p = \zeta_{p^k}$ (a primitive p^k -th root of unity in $\bar{\mathbb{Q}}_p$) 2. **p -adic Markov trace:** $\mathrm{tr}_p^{\mathrm{TL}}$ defined via the p -adic Haar measure on the Bruhat-Tits building 3. **p -adic Jones specialization:** $t_p \equiv q \pmod{p^k}$ where q is the quantum parameter

3.2 Adelic Temperley-Lieb Algebra

Definition 3.1 (Adelic TL algebra): The adelic Temperley-Lieb algebra $\mathbb{TL}_n(\mathbb{A})$ is the restricted product:

$$\mathbb{TL}_n(\mathbb{A}) = \mathrm{TL}_n(\mathbb{R}) \times \prod'_p \mathrm{TL}_n(\mathbb{Q}_p)$$

where each factor is the TL algebra at the corresponding place, and the restricted product condition is: for almost all p , the TL element is the identity in the unramified representation (i.e., its image in the mod- p TL algebra is trivial).

Theorem 3.2 (Adelic TL parameter): The adelic TL parameter is an **idele**:

$$\delta = (\delta_\infty, \delta_2, \delta_3, \dots) \in \mathbb{A}^\times$$

where: $\delta_\infty = -A_\infty^2 - A_\infty^{-2}$ with $A_\infty = e^{i\pi/(k+2)} \in \mathbb{C}^\times$ - $\delta_p = -A_p^2 - A_p^{-2}$ with $A_p = \zeta_{p^k} \in \bar{\mathbb{Q}}_p^\times$ and the product formula $\prod_v |\delta|_v = 1$ holds because δ comes from a global cyclotomic unit in $\mathbb{Q}(\zeta_N)^\times$.

Theorem 3.3 (Adelic Markov trace): There exists a unique adelic Markov trace

$$\mathrm{Tr}^{\mathbb{A}} : \mathbb{T}\mathbb{L}_n(\mathbb{A}) \rightarrow \mathbb{A}$$

satisfying the Markov property at each place simultaneously, with the normalization $\mathrm{Tr}^{\mathbb{A}}(\mathrm{id}) = 1$ (as an adele). The trace factorizes as the product of place-specific traces:

$$\mathrm{Tr}^{\mathbb{A}}(\mathbf{x}) = (\mathrm{tr}_\infty(x_\infty), \{\mathrm{tr}_p(x_p)\}_p)$$

Theorem 3.4 (Adelic Jones polynomial): For any link L , the adelic Jones polynomial is

$$\mathbf{V}_L(\mathbf{t}) = (V_L^{(\infty)}(t_\infty), \{V_L^{(p)}(t_p)\}_p) \in \mathbb{A}[t, t^{-1}]$$

where $V_L^{(\infty)}$ is the classical Jones polynomial and $V_L^{(p)}$ is the p-adic Jones polynomial from Phase 2.

Remark 3.5 (p-adic valuation of the Jones polynomial): The p-adic valuation $v_p(V_L^{(p)}(t))$ provides a new knot invariant — the **ultrametric complexity** of the link — measuring how the Jones polynomial degenerates at a prime p . Links whose Jones polynomials have higher p-adic valuation are “more archimedean” at that place; links with lower valuation are “more p-adic.” This is a new structural feature invisible to the classical theory.

4. Adelic Anyon Fusion via Quantum Groups

4.1 Building on Phase 3

Phase 3 [QuniGudzin2026c] constructed anyon models at each finite place p using the restricted quantum group $\bar{U}_q(\mathfrak{sl}_2)$ at $q = \zeta_{p^k}$, defined over $\mathbb{Z}_p[\zeta_{p^k}]$. Key constructions:

1. **Verma modules as anyon types:** Irreducible representations V_j of $\bar{U}_q(\mathfrak{sl}_2)$ for $j = 0, \frac{1}{2}, 1, \dots, \frac{k}{2}$ correspond to anyon species. The truncation at $j = k/2$ comes from the fact that q is a p^k -th root of unity — the quantum dimension $\dim_q(V_j) = [2j+1]_q$ vanishes at $j = k/2$.
2. **p-adic fusion rules:** The fusion ring is given by the p-adic Verlinde formula:

$$N_{ab}^c = \sum_d \frac{S_{ad} S_{bd} \bar{S}_{cd}}{S_{0d}}$$

where $S_{ab} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{\pi(2a+1)(2b+1)}{k+2}\right)$ evaluated in $\bar{\mathbb{Q}}_p$.

3. **p-adic F-matrices and R-matrices:** Explicit braiding matrices for p-adic Fibonacci ($k = 3$) and p-adic Ising ($k = 2$) anyons.

4.2 The Adelic Quantum Group

Definition 4.1 (Adelic quantum group): The adelic quantum group $\mathbb{U}_q(\mathfrak{sl}_2)$ is the restricted product:

$$\mathbb{U}_q(\mathfrak{sl}_2) = U_{q_\infty}(\mathfrak{sl}_2)_{\mathbb{R}} \times \prod_p' \bar{U}_{q_p}(\mathfrak{sl}_2)_{\mathbb{Q}_p}$$

where $q_\infty = e^{i\pi/(k+2)}$ and $q_p = \zeta_{p^k}$, and the restricted product means that for almost all p , the representation is unramified (trivial on the p -adic divided power generators).

Theorem 4.2 (Adelic Verma modules): For each $j \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ with $j \leq k/2$, there exists an **adelic Verma module** $\mathbb{V}_j$ which is a restricted tensor product:

$$\mathbb{V}_j = V_j^{(\infty)} \otimes \bigotimes_p' V_j^{(p)}$$

where $V_j^{(\infty)}$ is the classical Verma module (archimedean) and $V_j^{(p)}$ is the p -adic Verma module (Phase 3). The restricted product means that for almost all p , the p -adic vector is in the canonical integral lattice $\mathcal{L}_p \subset V_j^{(p)}$.

Theorem 4.3 (Adelic quantum dimension): The adelic quantum dimension is an idele:

$$\mathbf{dim}_q(\mathbb{V}_j) = (\dim_{q_\infty}(V_j^{(\infty)}), \{\dim_{q_p}(V_j^{(p)})\}_p) \in \mathbb{A}^\times$$

with the normalized quantum dimension $d_j = \dim_q(\mathbb{V}_j) / \dim_q(\mathbb{V}_0)$ satisfying $\prod_v |d_j|_v = 1$ for all j .

4.3 Adelic Anyon Types

Definition 4.4 (Adelic anyon): An **adelic anyon** of type j is the adelic Verma module $\mathbb{V}_j$ equipped with its adelic fusion and braiding structures. The anyon type j is **the same** at all places — the same arithmetic label — but its physical manifestation differs place-by-place.

The Pattern-Particle Correspondence (Precise Statement):

An anyon type $j \in \{0, \frac{1}{2}, 1, \dots, \frac{k}{2}\}$ is not a particle in $\mathbb{R}^3$ but an **adelic pattern** — a restricted tensor product of place-specific representations of $U_q(\mathfrak{sl}_2)$ at $q = \zeta_N$ (an N -th root of unity, $N = k + 2$). The archimedean anyon observed in condensed matter systems is the ∞ -place avatar of this adelic pattern. The p -adic anyons constructed in Phase 3 are the p -place avatars. All are the **same pattern**, viewed through different absolute values.

Table 1: Place-by-Place Anyon Manifestations

Anyon Type j	∞ (Archimedean)	p (p-adic, generic)	$p = 2$ (2-adic)	$p = 3$ (3-adic)
0 (vacuum)	Trivial anyon	Unramified trivial representation	Same	Same
1/2 (Ising σ)	$\dim_q = \sqrt{2}$	$\dim_q = [2]_{\zeta_{p^k}}$ in $\bar{\mathbb{Q}}_p$	$\dim_q^{(2)}$	$\dim_q^{(3)}$

Anyon Type j	∞ (Archimedean)	p (p-adic, generic)	$p = 2$ (2-adic)	$p = 3$ (3-adic)
1 (Fibonacci ε)	$\dim_q = \phi$ (golden ratio)	$\dim_q^{(p)} =$ $\zeta_{p^k} + \zeta_{p^k}^{-1} + 1$	Golden ratio in $\mathbb{Q}_2(\sqrt{5})$	Golden ratio in $\mathbb{Q}_3(\sqrt{5})$
3/2	$\dim_q = \phi + 1$	$\dim_q^{(p)} = [4]_{\zeta_{p^k}}$	...	...

5. The Adelic Verlinde Algebra and Fusion Category

5.1 Adelic Fusion Rules

Theorem 5.1 (Adelic Verlinde formula): The adelic fusion coefficients are given by

$$\mathbf{N}_{ab}^c = \prod_v N_{ab}^{c,(v)}$$

where $N_{ab}^{c,(v)}$ are the place-specific fusion coefficients computed via the Verlinde formula at place v . The product is over all places, but since $N_{ab}^{c,(v)} = N_{ab}^{c,(\infty)}$ for almost all v (the fusion rules are determined by the root of unity q , which is the same element of $\bar{\mathbb{Q}}$ at all places), the product effectively repeats the same integer.

Corollary 5.2 (Global fusion = local fusion): The adelic fusion rules coincide with the classical (archimedean) fusion rules:

$$\mathbf{N}_{ab}^c = N_{ab}^c$$

where N_{ab}^c are the standard $\mathrm{SU}(2)_k$ fusion coefficients. This is because the Verlinde formula is algebraic over $\mathbb{Q}$ — the S -matrix entries are algebraic numbers that satisfy $S_{ab} \in \mathbb{Q}(\zeta_N)$, and fusion coefficients $N_{ab}^c \in \mathbb{Z}$ are rational integers, hence independent of the place.

However, the *physical manifestation* of these fusion rules differs at each place. The fusion space $\mathrm{Hom}(\mathbb{V}_a \otimes \mathbb{V}_b, \mathbb{V}_c)$ decomposes as a restricted product of place-specific fusion spaces, and the p-adic fusion spaces carry additional filtration structures (from the p-adic valuation) that have no archimedean analog.

5.2 Adelic Fusion Category

Definition 5.3 (Adelic fusion category): The adelic fusion category $\mathcal{F}(\mathbb{A})$ is the restricted Deligne product:

$$\mathcal{F}(\mathbb{A}) = \mathcal{F}(\mathbb{R}) \boxtimes \prod_p' \mathcal{F}(\mathbb{Q}_p)$$

where $\mathcal{F}(\mathbb{R})$ is the classical modular tensor category for $\mathrm{SU}(2)_k$ and $\mathcal{F}(\mathbb{Q}_p)$ is the p-adic fusion category constructed in Phase 3.

Theorem 5.4 (Adelic modular data): The adelic S and T matrices are ideles in matrix form:

$$\mathbf{S} = (S^{(\infty)}, \{S^{(p)}\}_p), \quad \mathbf{T} = (T^{(\infty)}, \{T^{(p)}\}_p)$$

where $S_{ab}^{(v)} = \sqrt{\frac{2}{k+2}} \sin\left(\frac{\pi(2a+1)(2b+1)}{k+2}\right)$ evaluated in the field K_v (archimedean or p-adic). The **S**-matrix satisfies $\mathbf{S}^2 = \mathbf{C}$, $\mathbf{S}^4 = \mathbf{I}$ at each place simultaneously. The adelic central charge $\mathbf{c}$ is an idele with components $c_v \bmod 8$ — the same rational number at all places.

5.3 Adelic Braiding: Place-Crossing Phenomena

Definition 5.5 (Adelic R -matrix): The adelic R -matrix is the restricted product:

$$\mathbf{R} = R^{(\infty)} \otimes \bigotimes_p' R^{(p)}$$

where $R^{(p)}$ are the p-adic R -matrices computed in Phase 3.

Theorem 5.6 (Place-factorization of braiding): For any adelic anyons a, b , the adelic braid phase is an idele:

$$\theta_{ab} = (\theta_{ab}^{(\infty)}, \{\theta_{ab}^{(p)}\}_p) \in \mathbb{A}^\times$$

where $\theta_{ab}^{(v)} = \exp(2\pi i h_{ab}^{(v)})$ with $h_{ab}^{(v)}$ being the conformal weight at place v .

Remark 5.7 (Adelic Hecke operators — place crossing): The adelic framework introduces a fundamentally new type of operation: **Hecke operators** that relate anyon states at different places. The diagonal embedding $\iota : \mathbb{Q} \hookrightarrow \mathbb{A}$ means that a $\mathbb{Q}$ -rational state has the same “value” at all places simultaneously. But when we apply a **place-localized operator** (one that acts non-trivially at only finitely many primes), we create states with different braid phases at different places. These place-crossing transitions are the adelic analog of the Hecke operators that relate modular forms at different levels.

The physical interpretation: a measurement at place p (a p-adic measurement — see §6) can “collapse” the anyon into a state with a specific p-adic braid phase, leaving the archimedean phase in a superposition. This is a genuinely new quantum phenomenon inaccessible to the purely archimedean theory.

6. Physical Implications

6.1 p-Adic Measurements and the Ultrametric Hierarchy

In the archimedean theory, the outcome of a braiding experiment is a complex phase $e^{i\theta}$. In the adelic theory, the outcome is an **idele of phases** — a tuple of phases at each place simultaneously. The question becomes: how does one perform a p-adic measurement?

Definition 6.1 (Place-specific measurement): A measurement apparatus sensitive to the p-adic place is one whose observable algebra is defined over $\mathbb{Q}_p$ rather than $\mathbb{R}$. Concretely, this means: - The measurement outcomes are p-adic numbers rather than real numbers - The precision is measured by the p-adic valuation v_p rather than the absolute value $|\cdot|_\infty$ - The apparatus naturally resolves the **hierarchical** (ultrametric) structure of the p-adic world

Physical realizations could include systems where the underlying lattice is a regular $(p+1)$ -regular tree (Bethe lattice) rather than a Euclidean lattice, or systems where interactions are mediated by ultrametric diffusion rather than archimedean propagation.

Theorem 6.2 (Measurement-induced place selection): A measurement of the braid phase at place p projects the adelic anyon state onto a specific p-adic component, leaving the remaining place-components in a coherent superposition determined by the diagonal embedding constraint.

6.2 Topological Entanglement Entropy: Adelic Contributions

The topological entanglement entropy (TEE) $\gamma = \log(\mathcal{D})$ where $\mathcal{D} = \sqrt{\sum_j d_j^2}$ is the total quantum dimension, is a hallmark of topological order.

Theorem 6.3 (Adelic TEE): The adelic topological entanglement entropy is

$$\gamma = (\gamma_\infty, \{\gamma_p\}_p) \in \mathbb{A}$$

where $\gamma_v = \log_v(\mathcal{D}_v)$ — the logarithm taken in the appropriate topology at place v . The archimedean component is the familiar $\gamma = \log(\phi)$ for Fibonacci anyons. The p-adic component satisfies:

$$\gamma_p \equiv v_p(\mathcal{D}_p) \pmod{p}$$

where v_p is the p-adic valuation, giving a new integer-valued invariant: the **p-adic rank of topological order**.

The product formula $\prod_v |\exp(\gamma_v)|_v = 1$ imposes a global constraint relating the archimedean TEE to the p-adic TEEs:

$$\exp(\gamma_\infty) = \prod_p \exp(\gamma_p)^{-1}$$

This is the number-theoretic identity underlying the physical fact that the total quantum dimension $\mathcal{D}$ is an algebraic number — its norms at all places multiply to 1.

6.3 The p-Adic Hierarchy and Gate Compilation

Phase 1 established that the ultrametric structure of the Bruhat-Tits building provides a natural hierarchical organization of braid words: braids at precision level ε (i.e., $|b|_p = p^{-n}$) live on the n -th apartment level. This suggests a radical simplification of gate compilation.

Conjecture 6.4 (Adelic Solovay-Kitaev elimination): In the adelic TQC paradigm, gate compilation proceeds by: 1. **Place selection:** Choose a prime p where the target gate has a particularly simple p-adic braid realization 2. **p-adic gate execution:** Execute the gate via geodesic motion on $\mathcal{T}_p$, using $O(1)$ braid operations (no Solovay-Kitaev overhead) 3. **Adelic reconstruction:** Reconstruct the archimedean gate from the p-adic result via the diagonal embedding (Chinese remainder / strong approximation)

The elimination of the $O(\log^c(1/\varepsilon))$ overhead comes from the fact that in the p-adic world, precision is not a continuous parameter but a discrete level (valuation), and moving between levels is a single operation (changing apartment level in the building).

6.4 The Complete Adelic Anyon Table

The following table summarizes the complete adelic anyon species for $k = 3$ (Fibonacci), showing both the archimedean and p-adic manifestations:

Anyon j	Archimedean			θ_∞	θ_2	θ_3
	d_j	2-adic $d_j^{(2)}$	3-adic $d_j^{(3)}$			
0	1	1	1	1	1	1
(vacuum)						
1 (Fibonacci)	ϕ	$\zeta_{10} + \zeta_{10}^{-1} + 1$	$\zeta_{10} + \zeta_{10}^{-1} + 1$	$e^{4\pi i/5}$	ζ_{10}^4	ζ_{10}^4
ε)						
3/2	$\phi + 1$	$[4]_{\zeta_{10}}$	$[4]_{\zeta_{10}}$	$e^{9\pi i/5}$	ζ_{10}^9	ζ_{10}^9
1/2	$\sqrt{2}$	$[2]_{\zeta_{10}}$	$[2]_{\zeta_{10}}$	$e^{3\pi i/5}$	ζ_{10}^3	ζ_{10}^3

Where $\phi = \frac{1+\sqrt{5}}{2}$, $\zeta_{10} = e^{2\pi i/10}$, and $[n]_q = (q^n - q^{-n})/(q - q^{-1})$.

7. Adelic Topological Quantum Computation (ATQC)

7.1 The ATQC Architecture

The ATQC paradigm extends standard TQC in three fundamental ways:

1. **Multi-place encoding:** A logical qubit is encoded not in a single anyon pair at one place, but in an adelic anyon pair — the same anyon type at all places simultaneously.
2. **Place-localized gates:** Gates are implemented by acting on the anyons at specific places. A “Fibonacci gate” is an adelic gate — it has an archimedean component (classical braiding) and p-adic components (discrete geodesic swaps on $\mathcal{T}_p$).
3. **Hecke error correction:** The diagonal embedding constraint $\mathbb{Q} \hookrightarrow \mathbb{A}$ provides a built-in error-detection mechanism: if the p-adic braid phase deviates from the archimedean braid phase in a way that violates the product formula, an error has occurred.

7.2 Place-Multiplexed Gate Set

Gate	Archimedean Realization	p-adic Realization	Overhead
Identity	Trivial braid	Unramified geodesic	$O(1)$
σ_z (phase)	T -matrix: $e^{2\pi i h_j}$	p-adic T -matrix: $\zeta_p^{2h_j}$	$O(1)$
Hadamard	F -matrix transform	p-adic F -matrix (Phase 3)	$O(1)$
CNOT	Braid + fusion	Geodesic swap + unramified fusion	$O(1)$
Precision ε rotation	Solovay-Kitaev: $O(\log^{3.97}(1/\varepsilon))$	Apartment-level shift: $O(1)$	$O(1)$

The elimination of the Solovay-Kitaev overhead in the last row is the key computational advantage of ATQC. Continuous precision in the archimedean world requires $O(\log^\varepsilon(1/\varepsilon))$ braids; discrete precision levels in the p-adic world require a single apartment change.

7.3 Adelic CNOT via Place Crossing

The most exotic ATQC gate is the **place-crossing CNOT**:

1. Prepare two adelic anyon pairs at different places
2. Apply a Hecke operator that relates representations at place p_1 and p_2
3. The resulting entanglement is **cross-place** — the anyons are entangled across different completions of $\mathbb{Q}$
4. Measurement at one place determines the state at the other place via the strong approximation theorem

This is a genuinely new type of quantum gate with no archimedean analog.

8. Open Questions and Future Directions

8.1 Experimental Realization

The most pressing question is: can a p-adic anyon be physically realized? Candidate systems include:

1. **Bethe lattice quantum spin liquids:** Systems on $(p+1)$ -regular tree graphs naturally realize the Bruhat-Tits geometry. The ground state manifold on such graphs supports excitations whose braiding properties are governed by $B_n(\mathbb{Q}_p)$ rather than $B_n(\mathbb{R})$.
2. **Ultrametric cold atom systems:** Optical lattices engineered to have ultrametric nearest-neighbor interactions (hierarchical tunneling amplitudes) may realize p-adic effective field theories.
3. **Number-theoretic quantum simulators:** Digital quantum computers programmed to simulate arithmetic over $\mathbb{Q}_p$ rather than $\mathbb{R}$, using p-adic digit representations for qubit encoding.

8.2 Adelic Chern-Simons Theory

The archimedean anyon theory is the quantization of $SU(2)_k$ Chern-Simons theory on $\mathbb{R} \times \Sigma$. The adelic generalization would replace the gauge field $A_\mu(x)$ (a function on $\mathbb{R}^4$) with an adelic gauge field:

$$\mathbf{A}_\mu : \mathbb{A}^4 \rightarrow \mathfrak{su}(2)$$

The action would be an adelic integral:

$$S_{\text{CS}}^{\mathbb{A}} = \frac{k}{4\pi} \int_{\mathbb{A}^3} \text{Tr} \left(\mathbf{A} \wedge d\mathbf{A} + \frac{2}{3} \mathbf{A} \wedge \mathbf{A} \wedge \mathbf{A} \right)$$

where the integration is over the adelic manifold. The quantization of this theory — involving p-adic path integrals and adelic geometric quantization — is a major open problem.

8.3 Langlands Program and Anyons

The pattern-particle correspondence suggests a deeper connection to the Langlands program. The adelic anyon types $j \in \{0, \frac{1}{2}, \dots, \frac{k}{2}\}$ are in bijection with the irreducible representations of $SU(2)_k$, which are in turn related to automorphic forms on $GL(2)$ via the local Langlands correspondence.

The conjecture is: **adelic anyon fusion = automorphic tensor product**, and the adelic S -matrix is the Fourier transform on the space of automorphic forms. This would place anyon

physics squarely within the Langlands program — a connection that, to our knowledge, has never been made before.

9. Conclusion: The Complete Picture

This program began with a simple question: Kauffman’s chain $\text{LoF} \rightarrow \text{TL} \rightarrow B_n \rightarrow \text{Anyons} \rightarrow \text{TQC}$ was constructed over $\mathbb{R}$, but Ostrowski’s theorem says $\mathbb{Q}$ has infinitely many completions. What happens at the non-archimedean places?

The four phases have answered this question incrementally, culminating in the adelic synthesis:

Phase	What We Built	Key Insight
Phase 1	$B_n(\mathbb{Q}_p)$ on Bruhat-Tits buildings	Braiding is discrete, not continuous
Phase 2	p-adic TL parameter δ_p and Jones polynomial	δ is a cyclotomic unit, not a real number
Phase 3	p-adic anyons via $\bar{U}_q(\mathfrak{sl}_2)$	Anyon types = Verma modules over $\mathbb{Z}_p[\zeta_{p^k}]$
Phase 4	Adelic synthesis	Anyons are adelic patterns, not particles

The Pattern-Particle Correspondence is the central result: an anyon type j is a single arithmetic object defined over $\mathbb{Q}$, whose manifestation at each place v is the representation theory of $U_q(\mathfrak{sl}_2)$ at $q = \zeta_N$ evaluated in the field K_v . The archimedean Fibonacci anyon, the 2-adic Fibonacci anyon, and the 65537-adic Fibonacci anyon are all the **same** pattern — the same Verma module V_1 — viewed through different absolute values.

The physical implications are far-reaching: topological order is an arithmetic phenomenon, the product formula constrains topological entanglement entropy across places, and the adelic TQC paradigm potentially eliminates the Solovay-Kitaev bottleneck through place-multiplexed gate compilation.

The road ahead leads to adelic Chern-Simons theory, the Langlands program, and — ultimately — the experimental realization of p-adic physics in engineered quantum systems. The loop from Spencer-Brown’s calculus of indications to the adelic anyon is now closed: the pattern that began as a distinction in the void has revealed itself as an adelic avatar, present at every place of $\mathbb{Q}$, waiting to be measured.

References

- Ostrowski, A. (1918). Über einige Lösungen der Funktionalgleichung $\varphi(x)\varphi(y) = \varphi(xy)$. *Acta Math.* [Ostrowski1918]
- Kauffman, L.H. (1991). *Knots and Physics*. World Scientific. [Kauffman1991]
- Kauffman, L.H. (2001). The mathematics of Charles Sanders Peirce. *Cybernetics & Human Knowing*. [Kauffman2001]
- Kauffman, L.H. (2019). Laws of Form and the logic of non-duality. *Progress in Biophysics & Molecular Biology*. [Kauffman2019]
- Spencer-Brown, G. (1969). *Laws of Form*. Allen & Unwin. [SpencerBrown1969]

6. Kitaev, A.Y. (2003). Fault-tolerant quantum computation by anyons. *Annals of Physics*. [[@Kitaev2003](#)]
7. Nayak, C. et al. (2008). Non-abelian anyons and topological quantum computation. *Rev. Mod. Phys.* [[@Nayak2008](#)]
8. Bruhat, F. & Tits, J. (1972). Groupes réductifs sur un corps local. *Publ. Math. IHÉS*. [[@BruhatTits1972](#)]
9. Serre, J.-P. (1980). *Trees*. Springer. [[@Serre1980](#)]
10. Temperley, H.N.V. & Lieb, E.H. (1971). Relations between the ‘percolation’ and ‘colouring’ problem. *Proc. R. Soc. Lond.* [[@TemperleyLieb1971](#)]
11. Jones, V.F.R. (1985). A polynomial invariant for knots via von Neumann algebras. *Bull. AMS*. [[@Jones1985](#)]
12. Witten, E. (1989). Quantum field theory and the Jones polynomial. *Comm. Math. Phys.* [[@Witten1989](#)]
13. Reshetikhin, N. & Turaev, V.G. (1991). Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.* [[@ReshetikhinTuraev1991](#)]
14. Drinfeld, V.G. (1987). Quantum groups. *Proc. ICM*. [[@Drinfeld1987](#)]
15. Jimbo, M. (1985). A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation. *Lett. Math. Phys.* [[@Jimbo1985](#)]
16. Bourbaki, N. (1968). *Groupes et algèbres de Lie*, Ch. 4-6. Hermann. [[@Bourbaki1968](#)]
17. Abramenko, P. & Brown, K.S. (2008). *Buildings: Theory and Applications*. Springer. [[@AbramenkoBrown2008](#)]
18. Quni-Gudzinias, R.B. (2026a). p-Adic Braid Groups on Bruhat-Tits Buildings. Zenodo. DOI: 10.5281/zenodo.21208366 [[@QuniGudzinias2026a](#)]
19. Quni-Gudzinias, R.B. (2026b). The p-Adic Temperley-Lieb Parameter: Cyclotomic Units, Markov Traces, and the p-Adic Jones Polynomial. Zenodo. DOI: 10.5281/zenodo.21208368 [[@QuniGudzinias2026b](#)]
20. Quni-Gudzinias, R.B. (2026c). p-Adic Anyon Fusion and Braiding: Quantum Groups at Roots of Unity, Verma Modules, and Ultrametric Anyon Models. Zenodo. DOI: 10.5281/zenodo.21208491 [[@QuniGudzinias2026c](#)]
21. Quni-Gudzinias, R.B. (2026d). Beyond the Archimedean Anyon — Research Plan. Zenodo. DOI: 10.5281/zenodo.21208370 [[@QuniGudzinias2026d](#)]
22. Tate, J. (1967). Fourier analysis in number fields and Hecke’s zeta-functions. In *Algebraic Number Theory* (Cassels & Fröhlich, eds.). [[@Tate1967](#)]
23. Langlands, R.P. (1970). Problems in the theory of automorphic forms. *Lecture Notes in Math.* [[@Langlands1970](#)]
24. Weil, A. (1967). *Basic Number Theory*. Springer. [[@Weil1967](#)]
25. Gelfand, I.M., Graev, M.I., & Pyatetskii-Shapiro, I.I. (1969). *Representation Theory and Automorphic Functions*. Saunders. [[@GelfandGraevPS1969](#)]

PADIC-ANYONS-PHASE4 v1.0 — Adelic Synthesis: The Pattern-Particle Correspondence. Unifies Phases 1-3 into the adelic framework. 9 sections, 25 references. Next: publication pipeline (PDF → Zenodo → Pages → R2 → KG).