Lean formalization of an Assumptions of Physics technical brief: log

This log records which statements of the technical brief in this directory were checked in Lean, where the formal version differs from the brief, and any problems found in the brief. The brief is available at `https://assumptionsofphysics.org/resources/briefs`.

Setup

- Lean `v4.36.0-rc2`; Mathlib at commit `1b1ccf4b`; physlib (`master`) and TauCeti (commit `e7e7f52`, on the same Mathlib) as further dependencies.
- Build: `lake build` in the project root.
- Every file is checked with `#print axioms` on its main results. A result is recorded here only if it depends on no axioms beyond `propext`, `Classical.choice` and `Quot.sound`, and so has no `sorry`.
- Imports are kept to the Mathlib modules actually needed.
- Material that belongs in Mathlib is kept in `ForMathlib/`, in Mathlib's directory layout (library `ForMathlib`), and imported by the brief files.

1 Unphysicality of Hilbert spaces for quantum mechanics

Files.

- File: `briefs/HilbertUnphysicality/HilbertUnphysicality.lean`
- Library: `HilbertUnphysicality`
- Imports: `InconsistentConvergence`, one new `ForPhyslib` file:
  `ForPhyslib/QuantumMechanics/UnitaryEquivalence.lean` (a unitary carries the quantum model),
  and five new `ForMathlib` files:
  `ForMathlib/Analysis/InnerProductSpace/l2Diagonal.lean` (diagonal operators on ℓ^2),
  `ForMathlib/Analysis/SpecialFunctions/Gaussian/Erf.lean` (the error function),
  `ForMathlib/Analysis/InnerProductSpace/l2Space.lean` (separable Hilbert spaces are $\ell^2(\mathbb{N})$),
  `ForMathlib/MeasureTheory/Function/L2Space.lean` (L^2 is infinite-dimensional),
  `ForMathlib/MeasureTheory/Function/LpSpace/ChangeOfVariables.lean` (the unitary of a change of variables on $L^2(\mathbb{R})$)

Setting. Theorem 5 says that the Hilbert-space formulation QSTO:H is inconsistent with three physical conditions. The brief proves four mathematical propositions (Sec. 2) and then argues each part of Theorem 5 from one of them (Sec. 4). The propositions are checked in Lean, and Theorem 5 is stated and proved as `theorem5`, with each condition in its minimum form (see the open problems, "Precise readings").

Results.

Brief	Lean
Proposition 1	
$\mathcal{D}(A) \neq \mathcal{H}$ (Hellinger–Toeplitz)	`bounded_of_isSymmetric`, `not_isSymmetric_of_unbounded`
A is not continuous on $\mathcal{D}(A)$	`not_continuous_of_unbounded`
Proposition 2	
N with domain $\mathcal{D}(N)$ is self-adjoint	`numberOp`, `coe_domain_numberOp`, `isSelfAdjoint_numberOp` (from `lp.isSelfAdjoint_diagonal`)
N is unbounded, so not continuous on $\mathcal{D}(N)$	`exists_lt_norm_numberOp`, `not_continuous_numberOp`
$\mathcal{D}(N)$, $E_N(x) = \langle x\|Nx\rangle / \|x\|^2$	`numberDomain`, `expN`, `expN_eq_inner`
$x_j \to e_0$ in $\mathcal{D}(N) \setminus \{0\}$, $E_N(x_j) = 1$, $E_N(e_0) = 0$	`expN_not_continuousWithinAt`, `not_continuousOn_expN` (sequence from the inconsistent-convergence brief)
Proposition 3	
erf, $\|\operatorname{erf}\| < 1$, erf $\to \pm 1$	`Real.erf`, `Real.abs_erf_lt_one`, `Real.tendsto_erf_atTop`
$f(x) = \tan(\frac{\pi}{2} \operatorname{erf} x)$ is increasing	`frameChange`, `strictMono_frameChange`
f is a bijection of $\mathbb{R}$, $g = f^{-1}$, $g' > 0$	`surjective_frameChange`, `frameIso`, `hasDerivAt_frameIso_symm`
$U\psi = \phi$, at the level of densities	`cauchyDensity_frameChange_mul_deriv`: $\|\phi(f(x))\|^2 f'(x) = \|\psi(x)\|^2$
the same, integrated	`arctan_frameChange`: $\arctan(f(x))/\pi = \operatorname{erf}(x)/2$
ψ normalized, $\psi \in \mathcal{D}(X)$	`integral_gaussDensity`, `integrable_sq_mul_gaussDensity`
ϕ normalized, $\phi \notin \mathcal{D}(Y)$	`integral_cauchyDensity`, `not_integrable_sq_mul_cauchyDensity`
the unitary U, $(U\psi)(y) = \psi(g(y))\sqrt{g'(y)}$	`frameUnitary` (from `Lp.compDiffeo`), `coeFn_frameUnitary`
$\mathcal{D}(X) = \mathcal{D}(Y) = \{\psi : x\psi \in L^2\}$	`positionDomain`
$\psi \in \mathcal{D}(X)$, $U\psi = \phi$, $U\psi \notin \mathcal{D}(Y)$	`gaussState_mem_positionDomain`, `frameUnitary_gaussState`, `frameUnitary_gaussState_not_mem`
the minimum form of [B]WDQ fails	`not_forall_frameUnitary_mem_positionDomain`

Brief	Lean
Proposition 4	
separable infinite-dimensional Hilbert spaces are isomorphic	`nonempty_linearIsometryEquiv_of_separable`
$L^2(\mathbb{R}^n)$ is separable and infinite-dimensional	`L2.separableSpace_euclidean`, `L2.not_finiteDimensional_euclidean`
$L^2(\mathbb{R}^{n+1}) \cong L^2(\mathbb{R}^{m+1})$	`L2_euclidean_iso`
Theorem 5, [B]STDIST	
a unitary induces bijections between rays, density operators, self-adjoint operators	`mapRays`, `isDensityOperator_conjStarAlgEquiv_iff`, `bijective_conjLinearPMap`, `isSelfAdjoint_conjLinearPMap_iff`
and preserves the Born rule and the entropy	`born_map`, `IsDensityOperator.vonNeumannEntropy_conjStarAlgEquiv`
$L^2(\mathbb{R}^{n+1})$ and $L^2(\mathbb{R}^{m+1})$ carry the same QSTO:H model	`carriesQSTO`, `exists_carriesQSTO`
Theorem 5	
QSTO:H is inconsistent with [B]COBS, [B]WDQ and [B]STDIST	`theorem5`

Differences from the brief.

- *Proposition 1.* Both parts are stated for linear maps, without Mathlib's partially defined operators. Discontinuity is proved for an unbounded linear map on any normed space, which includes $\mathcal{D}(A)$ with the norm inherited from $\mathcal{H}$. Hellinger–Toeplitz is proved for an everywhere-defined symmetric map; self-adjointness is not needed.

- *Proposition 2.* The number operator `numberOp` is the diagonal operator with entries n on $\ell^2(\mathbb{N})$, on its maximal domain, as a partially defined operator (`LinearPMap`); its domain is `numberDomain` $= \{x : \sum_n n^2|x_n|^2 < \infty\}$. Self-adjointness is proved for every diagonal operator with real entries (`lp.diagonal`, in `ForMathlib`). The expectation `expN` $x = \sum_n n\,|x_n|^2/\|x\|^2$ is defined on all of ℓ^2 and equals $\mathrm{Re}\,\langle x|Nx\rangle\,/\|x\|^2$ on $\mathcal{D}(N)$ (`expN_eq_inner`). Discontinuity is proved within $\mathcal{D}(N)\setminus\{0\}$ with the subspace topology, as in the minimum form of [B]COBS. The sequence x_j is the brief's.

- *Proposition 3.* The wave functions are $\psi = (e^{-x^2}/\sqrt{\pi})^{1/2}$ (`gaussState`) and $\phi = (1/(\pi(1+y^2)))^{1/2}$, as elements of $L^2(\mathbb{R})$. Both frames use the same space $L^2(\mathbb{R})$ with Lebesgue measure. The position operators are represented by their common maximal domain `positionDomain` $= \{\psi : x\psi \in L^2\}$; the operators themselves, and their self-adjointness, are not formalized. That f is strictly increasing and onto $\mathbb{R}$, and that $g = f^{-1}$ has $g' > 0$, are proved; smoothness is not, only differentiability, which is all the unitary needs. The unitary U is `Lp.compDiffeo`, proved unitary for any bijection g of $\mathbb{R}$ that is differentiable with a differentiable inverse, with weight $\sqrt{|g'|}$; its inverse V is the same construction for g^{-1}. The operator UXU^{-1} and its domain are not formalized. Mathlib has no error function; `Real.erf` is defined in `ForMathlib`.

- *Proposition 4.* The classification of separable infinite-dimensional Hilbert spaces is new in `ForMathlib`. The positive integers n, m appear as $n+1, m+1$.

- *Theorem 5.* Stated as `theorem5`, a conjunction of three parts, one per condition, each for one particular system: the oscillator (Proposition 2), one spatial degree of freedom (Proposition 3), and any two numbers $n+1$, $m+1$ of continuous degrees of freedom (Proposition 4). "Inconsistent" is read as the failure of the minimum form of the condition for that system. The [B]COBS part states that N is self-adjoint on $\mathcal{D}(N)$, that $E_N(x) = \operatorname{Re}\langle x|Nx\rangle / \|x\|^2$ there, and that E_N is not continuous on $\mathcal{D}(N) \setminus \{0\}$. The [B]WDQ part states $\psi \in \mathcal{D}(X)$, $U\psi = \phi$ a.e. and $U\psi \notin \mathcal{D}(Y)$, for the specific unitary U of the change of frame (for an arbitrary unitary the statement would be trivial). The [B]STDIST part states that a unitary $L^2(\mathbb{R}^{n+1}) \to L^2(\mathbb{R}^{m+1})$ exists and carries the QSTO:H model. For [B]STDIST, that a unitary carries rays, density operators, self-adjoint operators, the Born rule and the entropy is proved for any unitary between complex Hilbert spaces (`carriesQSTO`). Rays are elements of `Projectivization`; density operators and the entropy are those of `ForPhyslib` (positive, trace class, trace one; entropy in $[0, \infty]$); self-adjoint operators are partially defined (`LinearPMap`). The closing remarks on classical cutoffs and on Schwartz spaces are not formalized.

- The additional remarks of Section 5 are not formalized.

Problems found in the brief.

- Cross-references lack their label: the proof of Theorem 5 has "This satisfies the conditions for 2", "As shown in 3" and "By 4" for Propositions 2, 3 and 4.

- "Schwartz spaces present the same problem" is asserted without proof, and without saying which structure on $\mathcal{S}(\mathbb{R}^n)$ is meant (see the open problems).

- QSTO:H still allows any separable Hilbert space, finite-dimensional ones included. This no longer affects Theorem 5, whose proof reads the inconsistency as the failure of a condition for one particular system, but the brief could say so.