Lean formalization of an Assumptions of Physics technical brief: log

This log records which statements of the technical brief in this directory were checked in Lean, where the formal version differs from the brief, and any problems found in the brief. The brief is available at `https://assumptionsofphysics.org/resources/briefs`.

Setup

- Lean `v4.36.0-rc2`; Mathlib at commit `1b1ccf4b`; physlib (`master`) and TauCeti (commit `e7e7f52`, on the same Mathlib) as further dependencies.
- Build: `lake build` in the project root.
- Every file is checked with `#print axioms` on its main results. A result is recorded here only if it depends on no axioms beyond `propext`, `Classical.choice` and `Quot.sound`, and so has no `sorry`.
- Imports are kept to the Mathlib modules actually needed.
- Material that belongs in Mathlib is kept in `ForMathlib/`, in Mathlib's directory layout (library `ForMathlib`), and imported by the brief files.

1 Quantum pure states are not enough

Files.

- File: `briefs/PureStatesNotEnough/PureStatesNotEnough.lean`
- Library: `PureStatesNotEnough`
- Imports: six new `ForMathlib` files:
  `ForMathlib/Analysis/Convex/Combination.lean`: a map that preserves the convex combinations of two points of a convex set preserves all finite convex combinations (`Convex.map_sum_of_map_add`), and a convex set contains them (`Convex.sum_mem_of_semifield`), with scalars in any ordered semifield such as $\mathbb{R}_{\geq 0}$.
  `ForMathlib/MeasureTheory/Constructions/HaarToSphere.lean`: Archimedes' hat-box theorem. The map $(z, \theta) \mapsto (\sqrt{1-z^2}\cos\theta, \sqrt{1-z^2}\sin\theta, z)$ is injective on $(-1, 1) \times (-\pi, \pi]$ with image the sphere minus its poles and area Jacobian 1 (`euclideanHausdorffMeasure_image_cylinderToSphere`, by TauCeti's area formula), so Mathlib's surface measure is $dz\, d\theta$ (`Measure.map_toSphere_volume_eq_map_cylinderToSphere`) and the height is distributed as $2\pi\, dz$ (`Measure.map_toSphere_volume_height`).

`ForMathlib/Topology/Compactification/OnePoint/SimplyConnected.lean`: the one-point compactification of a real vector space of dimension at least two is simply connected (`OnePoint.simplyConnectedSpace_of_two_le_finrank`), by Mathlib's homeomorphism with the sphere and TauCeti's simple connectivity of spheres; in particular the Riemann sphere is.
`ForMathlib/Topology/Algebra/Projectivization.lean`: a continuous linear equivalence $e : V \simeq W$ induces a homeomorphism $P(V) \simeq P(W)$, $[v] \mapsto [ev]$, of the quotient topologies (`Projectivization.mapHomeomorph`).
`ForMathlib/Geometry/Manifold/Instances/HopfMap.lean`: the Hopf map $S^{2n+1} \to \mathbb{C}P^n$, $x \mapsto [x]$ (`Projectivization.hopfMap`), from the unit sphere of $\mathbb{C}^{n+1}$ to $\mathbb{C}P^n = P(\mathbb{C}^{n+1})$ with TauCeti's real manifold structure, is onto, smooth (`Projectivization.contMDiff_hopfMap`) and a submersion (`Projectivization.isSubmersion_hopfMap`): in the affine chart $x_i \neq 0$ it is $x \mapsto (x_j/x_i)_{j\neq i}$, and its derivative is onto because $z \mapsto (x_i/|x_i|)\,\hat{z}/\|\hat{z}\|$, with $\hat{z}$ the vector z with 1 inserted at i, is a smooth local section through x.
`ForMathlib/Geometry/Symplectic/Manifold/FubiniStudy.lean`: the Fubini–Study form on $\mathbb{C}P^n$ by symplectic reduction. The standard form $\omega(u, w) = \mathrm{Im}\langle u, w\rangle = \langle iu, w\rangle_{\mathbb{R}}$ of $\mathbb{C}^{n+1}$ restricts to a closed form σ on the unit sphere (`Projectivization.sphereForm`, `Projectivization.isClosed_sphereForm`). At a unit vector x the kernel of σ and the kernel of the derivative of the Hopf map are both the line spanned by the tangent vector ix (`Projectivization.ker_bilinFormAt_sphereForm`): the first because the tangent space is the real orthogonal complement of x, the second by counting dimensions, $2n + 1 - 2n = 1$, since the Hopf map is constant along $e^{it}x$. Unit scalars preserve ω and commute with the Hopf map, so σ is basic (`Projectivization.isBasic_sphereForm`). TauCeti's symplectic descent then gives the Fubini–Study form `Projectivization.fubiniStudyForm`, which pulls back to σ (`Projectivization.mpullback_fubiniStudyForm`) and is symplectic (`Projectivization.isSymplectic_fubiniStudyForm`). A unitary map U of $\mathbb{C}^{n+1}$ induces a smooth map $[v] \mapsto [Uv]$ of $\mathbb{C}P^n$ (`Projectivization.unitaryMap`) that commutes with the Hopf map, and since U preserves the standard form, uniqueness of the descended form shows that it preserves the Fubini–Study form (`Projectivization.mpullback_unitaryMap_fubiniStudyForm`).
Five new `ForPhyslib` files on the qubit, in `ForPhyslib/QuantumMechanics/Qubit/`:
`BlochSphere.lean`: the Bloch point of a ray (`blochPoint`), injective, onto the unit sphere, and a homeomorphism `blochSphere` : $P(\mathbb{C}^2) \simeq S^2$; $|\psi\rangle\langle\psi| = \frac{1}{2}(1 + a \cdot \sigma)$ (`hermitianProjector_eq`).
`BlochRotation.lean`: a unitary U acts on Bloch points by TauCeti's rotation `pauliRotation` U (`blochPoint_unitaryAction`); every rotation is the action of an element of $SU(2)$ (`exists_unitaryAction_eq`); two unitaries act alike iff they differ by a phase (`unitaryAction_eq_iff`); complex conjugation of the components is the reflection $a_1 \mapsto -a_1$ (`blochPoint_conjAction`) and no unitary action (`conjAction_ne_unitaryAction`).
`AntisymplecticReflection.lean`: complex conjugation `conjSphere` of the Riemann sphere is smooth, preserves the Fubini–Study density (`fubiniStudyDensity_conj`), pulls the Fubini–Study form back to its negative (`mpullback_conjSphere`), so no map that preserves the form is complex conjugation (`ne_conjSphere_of_mpullback_eq`), nor any time-t map of a Hamiltonian flow (`globalFlow_ne_conjSphere`); it corresponds to `conjAction` (`equivProjectivization_conjSphere`).
`HamiltonianIsotopy.lean`: on a simply connected manifold, a smooth vector field with closed contraction $\iota_X\sigma$ is Hamiltonian (`SmoothTwoForm.mem_symplecticVectorFields_iff_exists_isHamiltonianVectorField`), so for a closed σ the fields with $L_X\sigma = 0$

are the Hamiltonian ones; a time-dependent Hamiltonian flow preserves a closed form (`SmoothTwoForm.mpullback_timeDependentFlowOf_of_isHamiltonianVectorField`), and on a simply connected manifold a jointly smooth time-dependent flow preserves it exactly when every velocity field is Hamiltonian (`SmoothTwoForm.forall_mpullback_timeDependentFlowOf_eq_iff`); a smooth isotopy with jointly smooth inverses is the time-dependent flow of its generating field $X_t = \dot{\varphi}_t \circ \varphi_t^{-1}$ (`isotopyField`, `contMDiff_isotopyField`, `timeDependentFlowOf_isotopyField`), so an isotopy from the identity preserves a closed form exactly when every X_t is Hamiltonian (`SmoothTwoForm.forall_mpullback_isotopy_eq_iff`); on the Riemann sphere, with completeness from compactness, no time map of a time-dependent Hamiltonian flow is complex conjugation (`timeDependentFlowOf_ne_conjSphere`), nor any map of a Hamiltonian isotopy (`isotopy_ne_conjSphere`).
`Precession.lean`: every Hermitian H is $c + \frac{1}{2} n \cdot \sigma$ (`exists_eq_smul_one_add_pauli`); along a solution of $i\psi' = H\psi$ the Bloch vector satisfies $a' = n \times a$ (`hasDerivAt_blochVector`).
These use TauCeti's Pauli map, double cover $SU(2) \to SO(3)$, Hermitian projectors, topology of projective space, its real manifold structure and affine charts, submersions and their criterion by surjective derivatives, symplectic descent along surjective submersions, Fubini–Study form on `OnePoint` $\mathbb{C}$, global Hamiltonian flows, area formula and identification of `Measure.toSphere` with Hausdorff measure, simple connectivity of spheres, vanishing of H^1_{dR} on simply connected manifolds, symplectic vector fields and the class $[\iota_X \sigma]$, and time-dependent flows with the Lie derivative of forms along them.
Existing files: `ForMathlib/Analysis/InnerProductSpace/l2Space.lean` (`Orthonormal.countable`), `ForPhyslib/QuantumMechanics/DensityOperator.lean` (`IsDensityOperator`, `posTrace`), `ForPhyslib/QuantumMechanics/Wigner/Density.lean` (`IsDensity`, `proj`), `ForPhyslib/ProbabilisticTheory/HilbertSpace/TraceClass/CanonicalExpansion.lean` (`IsTraceClass.exists_hasSum_rankOne`) and `ForPhyslib/QuantumMechanics/Kadison.lean` (`isPure_iff_isExtremeDensityOperator`).

Setting. The ensembles of CST over a measurable space X with Liouville measure μ are the probability measures $\nu \ll \mu$ (`classicalEnsembles` μ). Their mixtures are $a\nu_1 + b\nu_2$ with $a, b \in \mathbb{R}_{\geq 0}$, $a + b = 1$. `HasManyDisjointSets` μ says that for every N there are N disjoint measurable sets of positive finite measure. The density operators of QE-DENS are `IsDensity` (positive, trace one) on a finite-dimensional space and `IsDensityOperator` (positive, trace class, trace one) on a separable complex Hilbert space $\mathcal{H}$ of any dimension. The pure states (`pureStates`) are the projections $|\psi\rangle\langle\psi|$ with $\|\psi\| = 1$, a copy of $P(\mathcal{H})$ inside the bounded operators, with the Borel σ-algebra of the operator norm. An identification of two ensemble spaces is modelled as a map that preserves mixtures of two ensembles.

Results.

Brief	Lean
CST, the ensemble space	`classicalEnsembles`; it is convex: `Convex.classicalEnsembles`
(used for Theorem 3) a Darboux chart has N disjoint sets of positive finite Liouville measure for every N	`hasManyDisjointSets_restrict`: any nonempty open set in a nontrivial finite-dimensional real normed space, for any measure that is positive on open sets and finite on compact sets
(used for Theorem 3) the CST ensemble space is infinite-dimensional	`not_injOn_of_map_add`: no mixture-preserving map from it into a finite-dimensional real vector space is injective
Theorem 3, CST and QE-DENS are inconsistent	`cst_qeDens_inconsistent`, for $\dim \mathcal{H} < \infty$
Theorem 1, QP-RAY and CST are consistent iff $1 \leq \dim V < \infty$	only in part: the "if", without the Liouville measure, and the edges below
(Theorem 1, "if") $P(V)$ with the Fubini–Study form is a symplectic manifold, for $V = \mathbb{C}^{n+1}$	`existsUnique_isSymplectic_projectivization`: $\mathbb{C}P^n$ has exactly one symplectic form whose pullback along the Hopf map is the restricted standard form of $\mathbb{C}^{n+1}$; for $n = 0$ it lives on a point
(Theorem 1, "if") the same for every V of finite dimension $n + 1$	`basisHomeomorph`: an orthonormal basis identifies $P(V)$ with $\mathbb{C}P^n$ as topological spaces; `contMDiff_basisHomeomorph_symm_trans`, `mpullback_basisHomeomorph_symm_trans`: the change of basis is smooth and preserves the Fubini–Study form, so the structure carried over does not depend on the basis
(Theorem 1) $\dim V = 0$	`isEmpty_projectivization`: $P(V)$ is empty; `nonempty_projectivization_iff`
(Theorem 1) $\dim V = 1$	`subsingleton_projectivization`: $P(V)$ is a single point
(Theorem 1) $2 \leq \dim V$	`nontrivial_projectivization_iff`: in finite dimension $P(V)$ has two points iff $2 \leq \dim V$
(Theorem 3 at $\dim \mathcal{H} = 1$) a point with Liouville measure δ	`not_hasManyDisjointSets_dirac`; `cst_qeDens_consistent_of_finrank_one`: an injective mixture-preserving map exists
(Theorem 1, "only if") $P(\ell^2)$ versus the sphere	`mk_projectivization`: $P(V)$ has $\mathfrak{c}$ points; `nonempty_projectivization_equiv`: $P(\ell^2) \simeq P(\mathbb{C}^2)$ as sets
Remark after Example 2, the Born rule is not $\delta_{\psi\phi}$	`born_ne_delta`: e_0 and $(e_0 + e_1)/\sqrt{2}$ are distinct rays with Born probability $\frac{1}{2}$
Proof of Theorem 5, the pure states are the extreme points of $D(\mathcal{H})$	`pureStates_eq_extremePoints`
Theorem 5, the map $r : M^1(P(\mathcal{H})) \to D(\mathcal{H})$	`barycenter` $r(\mu) = \int \rho \, d\mu(\rho)$; `isDensityOperator_barycenter`, `posTrace_barycenter`; `toDensity`
Theorem 5, r is affine	`barycenter_add`
Theorem 5, $D(\mathcal{H}) \equiv M^1(P(\mathcal{H}))/\sim$	`toDensity_surjective`, `densityEquiv`
Remark after Theorem 5, a class records the different preparations of one ensemble	`exists_barycenter_eq_and_ne`: $\frac{1}{2}\delta_{e_0} + \frac{1}{2}\delta_{e_1}$ and $\frac{1}{2}\delta_{e_+} + \frac{1}{2}\delta_{e_-}$ have the same barycenter but give e_0 different probabilities

Brief	Lean
Example 2, $P(\mathbb{C}^2)$ is a two-sphere	`nonempty_homeomorph_sphere`, from `blochSphere`: a homeomorphism onto the unit sphere of $\mathbb{R}^3$
Example 2, Darboux coordinates z, θ_{xy}, the Liouville measure is the area	`toSphere_eq_dz_dtheta`: the surface measure is $dz\,d\theta$ on $(-1,1)\times(-\pi,\pi]$; the height is distributed as $2\pi\,dz$
Example 2, Hamiltonian evolutions are the volume-preserving diffeomorphisms	`conjSphere_preserves_area_reverses_form`, `globalFlow_ne_conjSphere`, `timeDependentFlowOf_ne_conjSphere`: complex conjugation of the Riemann sphere preserves the area but is not Hamiltonian (see "Problems found"); with the area form, `lieDeriv_areaForm_eq_zero_iff_hamiltonian`: the vector fields that preserve it are the Hamiltonian ones, `preserves_areaForm_iff_hamiltonian`: the time-dependent flows that preserve it are the Hamiltonian evolutions, and `isotopy_preserves_areaForm_iff_hamiltonian`: so are the isotopies from the identity that preserve it
Example 2, quantum evolutions are exactly the rotations	`exists_unitaryAction_eq_iff`: a map of rays is the action of a unitary iff it is a rotation on Bloch vectors; `unitaryAction_eq_iff`: unitaries act alike iff they differ by a phase
Example 2, linear Hamiltonians $h = n \cdot x$ give rotations about n	`exists_blochVector_precession`: quantum side only; every Hamiltonian $c + \frac{1}{2} n \cdot \sigma$ gives $a' = n \times a$

Differences from the brief.

- Theorem 3 compares the whole ensemble spaces, not those supported on a two-dimensional subspace A (see "Problems found"). It assumes $\dim \mathcal{H} < \infty$, which the brief derives from Theorem 1. Only the finite dimension of the real vector space of operators is used, not positivity or trace. "The spaces correspond" becomes a map from the CST ensembles to the density operators that is injective and preserves mixtures of two ensembles. The proof normalizes μ on $\dim + 2$ disjoint sets, takes an affine dependence among the images, and separates its positive and negative parts into two different mixtures with the same image.

- The Liouville measure enters only through `HasManyDisjointSets`, which holds on every Darboux chart. The symplectic structure itself is not used.

- Theorem 5 is proved for every separable complex Hilbert space, not only in finite dimension. $r(\mu)$ is the Bochner integral of the pure states in the operator norm. It is positive because the positive operators form a closed convex cone, and it has trace one by Parseval and monotone convergence over a countable Hilbert basis. r is onto by the canonical expansion $\rho = \sum_i \lambda_i |e_i\rangle\langle e_i|$, represented by $\sum_i \lambda_i \delta_{e_i}$. No Choquet theory is used. "$\equiv$" is the bijection between the quotient by $r(\mu_1) = r(\mu_2)$ and $D(\mathcal{H})$. That the pure states are the extreme points of $D(\mathcal{H})$, where the brief's proof starts, is checked separately, in Mathlib's sense of extreme points of a real convex set; the rest of the proof does not use it.

- Theorem 1, "if", is checked for $V = \mathbb{C}^{n+1}$ (`Fin` $(n+1) \to \mathbb{C}$), with $P(V)$ the real manifold of dimension $2n$ given by TauCeti's affine charts. For a complex inner product space V of

dimension $n+1$, $P(V)$ is not given charts of its own in Lean. An orthonormal basis identifies it with $\mathbb{C}P^n$, and the statement is that the identification is unique up to a smooth map of $\mathbb{C}P^n$ that preserves the Fubini–Study form; the structure carried over is therefore well defined. QP-RAY asks only for a complex vector space; a Hermitian inner product is needed to single out the Fubini–Study form. "X is a symplectic manifold" becomes a closed, nondegenerate smooth two-form on $P(V)$. The Fubini–Study form is defined by reduction, as the descent of the standard form of $\mathbb{C}^{n+1}$ restricted to the unit sphere, not by a Kähler potential; in this normalization a projective line should have area π (not checked), while TauCeti's form on the Riemann sphere, used in Example 2, has total area 1. The two are not compared in Lean. The Liouville measure ω^n, which CST also requires, is not constructed.

- Example 2 is checked at the level of topology and measure, and its symplectic part on the Riemann sphere `OnePoint` $\mathbb{C}$ with TauCeti's Fubini–Study form. The sphere is the unit sphere of $\mathbb{R}^3$ with Mathlib's surface measure, which TauCeti identifies with the Hausdorff area; "Darboux coordinates" becomes the statement that this measure is $dz\,d\theta$. The area-preserving map that is not Hamiltonian is complex conjugation of the Riemann sphere, which corresponds to the Bloch reflection $a_1 \mapsto -a_1$; area preservation is invariance of the Fubini–Study density, since there is no measure on `OnePoint` $\mathbb{C}$ yet. Quantum evolution is the action of a unitary on rays, and for a Hamiltonian the Schrödinger equation is a hypothesis on ψ; the precession $a' = n \times a$ is not integrated to a finite rotation. "Hamiltonian evolution" is checked for vector fields and for evolutions: the time-dependent flows of jointly smooth time-dependent vector fields, which preserve the Fubini–Study form exactly when each velocity field X_t is the Hamiltonian vector field of a smooth H_t, and the smooth isotopies from the identity with jointly smooth inverses, which preserve it exactly when each generating field $X_t = \dot{\varphi}_t \circ \varphi_t^{-1}$ is Hamiltonian. The H_t are found one time at a time, not as one jointly smooth $H(t, x)$.

- Not formalized:
  - in Theorem 1: the Liouville measure of the Fubini–Study form, and the "only if";
  - in Example 2: the Liouville measure of the Fubini–Study form and the symplectic form $dz \wedge d\theta$ in the chart; the diffeomorphism $\mathbb{C}P^1 \cong S^2$ with the comparison of the Fubini–Study and round area forms; ensembles as densities $\rho(z, \theta)$ (Radon–Nikodym, not restated); the classical side of linear Hamiltonians (moment maps); that every orientation-preserving, area-preserving diffeomorphism of the sphere is the end of a Hamiltonian evolution, which needs Smale's theorem $\mathrm{Diff}^+(S^2) \simeq SO(3)$;
  - Insight 4, which is informal.

Problems found in the brief. Theorem 5 holds, and Theorem 3 holds for $\dim \mathcal{H} \geq 2$, but their proofs have gaps. Example 2 holds for evolutions once "volume-conserving" is read as "area-form-preserving". At $\dim \mathcal{H} = 1$ Theorem 3 depends on a convention of Theorem 1. Of Theorem 1 only the "if" is checked, without the Liouville measure.

- Theorem 1 states $1 \leq \dim V$, while its proof uses $2 \leq \dim X$. For $\dim V = 1$ the state space $P(V)$ is a single point (`subsingleton_projectivization`), a manifold of dimension 0. So either 0-dimensional symplectic manifolds count, and the proof's bound should be $0 \leq \dim X$, or they do not, and the theorem should say $2 \leq \dim V < \infty$. The choice also decides Theorem 3. If a point counts, then for $\dim \mathcal{H} = 1$, which QE-DENS allows, the CST ensembles of the point are $\{\delta\}$ and the density operators are $\{1\}$, and the constant map identifies the two (`cst_qeDens_consistent_of_finrank_one`): Theorem 3 fails. The

Lean proof of Theorem 3 avoids this through `HasManyDisjointSets`, which fails for a point (`not_hasManyDisjointSets_dirac`). So either Theorem 1 says $2 \leq \dim V$, or Theorem 3 needs $\dim \mathcal{H} \geq 2$.

- For $\dim V = 0$, $P(V)$ is empty (`isEmpty_projectivization`), and the empty set is a symplectic manifold of every dimension. Either lower bound, 1 or 2, needs CST to exclude it, for instance by requiring an ensemble: there is no probability measure on the empty set.

- The "only if" of Theorem 1 needs the identification of X with $P(V)$ to respect the topology. As sets, $P(\ell^2)$ and $P(\mathbb{C}^2)$, the sphere, are in bijection: both have $\mathfrak{c}$ points (`nonempty_projectivization_equiv`). Carrying the symplectic structure of the sphere over along such a bijection would make $P(\ell^2)$ a symplectic manifold. The theorem holds when V is Hausdorff and X has the quotient topology of $P(V)$; for a non-Hausdorff V, $P(V)$ need not be Hausdorff even in finite dimension, and the "if" fails. "Since X is a symplectic manifold, $\dim X < \infty$" holds only for finite-dimensional symplectic manifolds, and the projective space of an infinite-dimensional Hilbert space is an infinite-dimensional one. The finiteness comes from CST itself: the Liouville measure ω^n needs a finite n.

- In the proof of Theorem 3, for $\dim \mathcal{H} \geq 3$ the sphere $P(A)$ has dimension $2 < 2(\dim \mathcal{H} - 1)$ and therefore Liouville measure zero. No CST ensemble has support in $P(A)$, so the classical side is empty, not infinite-dimensional. The argument works as written only for $\dim \mathcal{H} = 2$. Comparing the whole spaces repairs it.

- The proof of Theorem 5 cites Choquet theory, which concerns compact convex sets, but $D(\mathcal{H})$ is not compact when $\dim \mathcal{H} = \infty$. The proof also never shows that every density operator is represented, which $D(\mathcal{H}) \equiv M^1/\sim$ needs. Both gaps are closed by the canonical expansion. "Every continuous linear functional $f : D(\mathcal{H}) \to \mathbb{R}$" should read continuous affine function, since $D(\mathcal{H})$ is not a vector space.

- Example 2: area-preserving diffeomorphisms of the sphere include the orientation-reversing ones, such as $z \mapsto -z$, which are not Hamiltonian. So Hamiltonian evolution preserves the area form, not only the area. Checked in Lean for complex conjugation of the Riemann sphere, the reflection $a_1 \mapsto -a_1$ of the Bloch sphere: it preserves the Fubini–Study density but reverses the form, and no time map of a Hamiltonian flow, autonomous or time-dependent, or of a Hamiltonian isotopy equals it (`conjSphere_preserves_area_reverses_form`, `globalFlow_ne_conjSphere`, `timeDependentFlowOf_ne_conjSphere`, `isotopy_ne_conjSphere`). With the area form, the claim holds for evolutions: a time-dependent flow, or an isotopy from the identity, preserves the form exactly when it is Hamiltonian (`preserves_areaForm_iff_hamiltonian`, `isotopy_preserves_areaForm_iff_hamiltonian`). The obstruction would lie in $H^1_{dR}(S^2) = 0$, so no flux homomorphism is involved. Along an evolution from the identity, preserving the area and preserving the form agree, since the sign of the pulled-back form cannot jump (not checked in Lean); only single maps tell them apart.

- CST says $p(\xi_1 \mid \xi_2) = 0$ "for all $\xi_1, \xi_2 \in X$", which includes $p(\xi \mid \xi) = 0$. It should say $\xi_1 \neq \xi_2$, as the Remark after Example 2 ($p(\psi \mid \phi) = \delta_{\psi\phi}$) has it.

- Typos: "non necessarily" should be "not necessarily"; "r defined about" should be "r defined above".