Lean formalization of an Assumptions of Physics technical brief: log

This log records which statements of the technical brief in this directory were checked in Lean, where the formal version differs from the brief, and any problems found in the brief. The brief is available at `https://assumptionsofphysics.org/resources/briefs`.

Setup

- Lean `v4.36.0-rc2`; Mathlib at commit `1b1ccf4b`; physlib (`master`) and TauCeti (commit `e7e7f52`, on the same Mathlib) as further dependencies.
- Build: `lake build` in the project root.
- Every file is checked with `#print axioms` on its main results. A result is recorded here only if it depends on no axioms beyond `propext`, `Classical.choice` and `Quot.sound`, and so has no `sorry`.
- Imports are kept to the Mathlib modules actually needed.
- Material that belongs in Mathlib is kept in `ForMathlib/`, in Mathlib's directory layout (library `ForMathlib`), and imported by the brief files.

1 Physical vector spaces for quantum mechanics

Files.

- Files: `briefs/QuantumVectorSpaces/QuantumVectorSpaces.lean`, and `briefs/QuantumVectorSpaces/QuantumVectorSpaces/Countable.lean` (a countable generating set of observables, Sec. 5)
- Library: `QuantumVectorSpaces`
- Imports: `SchwartzNorms` (the L^2 seminorm family on $\mathcal{S}(\mathbb{R})$) and new `ForMathlib` files: `ForMathlib/Analysis/InnerProductSpace/Defs.lean` (the seminorm $\sqrt{\mathrm{re}\langle x, x\rangle}$ of a `PreInnerProductSpace.Core`, and continuity of the inner product against it), `ForMathlib/Analysis/CStarAlgebra/Matrix.lean` (`Matrix.toEuclideanLin` is multiplicative), `ForMathlib/Analysis/Distribution/SchwartzSpace/Complete.lean` (`SchwartzMap.instCompleteSpace`: $\mathcal{S}(E, F)$ is complete for complete F), `ForMathlib/Analysis/Distribution/SchwartzSpace/SupNorm.lean` (`SchwartzMap.exists_norm_le_toLp_two`: a Sobolev bound for the values of a Schwartz function, from TauCeti's `SchwartzMap.exists_integral_norm_fourierInv_le`),

ForMathlib/Analysis/InnerProductSpace/MultilinearBasis.lean
(ContinuousMultilinearMap.norm_le_sum_orthonormalBasis: a multilinear map is bounded by its values on an orthonormal basis),
ForMathlib/Analysis/LocallyConvex/FirstCountable.lean
(WithSeminorms.exists_countable: a family of seminorms inducing a first-countable topology has a countable subfamily inducing it)

Setting. V is a complex vector space with an inner product given as a PreInnerProductSpace.Core. Unlike Mathlib's InnerProductSpace, this puts no norm or topology on V. The defining observables are a family X_i ($i \in \iota$) in Module.End, and algebra is the unital algebra they generate (Algebra.adjoin). The seminorms $p_A(v) = \sqrt{\langle Av, Av\rangle}$ form seminormFamily, and topology is the topology they generate. A topology is IsAdmissible if it makes V a topological vector space and makes the X_i and the inner product (jointly) continuous. For Example 3, $Xf = xf$ and $Pf = -i\hbar f'$ act on Mathlib's $\mathcal{S}(\mathbb{R}, \mathbb{C})$ (posOp, momOp), with the L^2 inner product l2Core. For Example 4, $S_i = (\hbar/2)\sigma_i$ acts on $\mathbb{C}^2$ as EuclideanSpace (spin). In n dimensions, $X^i f = x_i f$ and $P_i f = -i\hbar\,\partial_{e_i} f$ act on $\mathcal{S}(\mathbb{R}^n, \mathbb{C})$, with $\mathbb{R}^n$ as EuclideanSpaceâD̊I(Finn) (posOpN, momOpN, indexed by ObsNn), and the L^2 inner product l2CoreN.

Definition 1 is IsQuantumVectorSpacecXÄğC: the X_i are symmetric, they satisfy $[X_i, X_j] = i\hbar \sum_k C_{ijk} X_k$ with finitely many nonzero C_{ijk} for each i, j, and V with topology is complete and second-countable. Completeness refers to uniformSpace, the uniform structure that every topological vector space carries.

Results.

Brief	Lean
Corollary 2, coarsest topology	`isGreatest_topology` (from `isAdmissible_topology` and `IsAdmissible.continuous_algebra`)
Corollary 2, locally convex	`locallyConvexSpace`
finite dimension (Example 4)	`topology_eq_moduleTopology`, `topology_qubit`
Example 3, $[X, P] = i\hbar$	`posOp_mul_momOp_sub`
Example 3, X, P, I symmetric	`isSymmetric_obs`
Example 3, Remark: the topology is that of $\mathcal{S}$	`topology_schwartz`, `withSeminorms_schwartz`, `isAdmissible_schwartz` (for $\hbar \neq 0$)
Example 4, $[S_i, S_j] = i\hbar\,\epsilon_{ijk}S_k$	`spin_comm`
Example 4, S_i symmetric	`isSymmetric_spin`
Example 4 is a quantum vector space	`isQuantumVectorSpace_qubit`
Example 3, second countability	`secondCountableTopology_schwartz`
Example 3, completeness	`SchwartzMap.instCompleteSpace`
Example 3 is a quantum vector space	`isQuantumVectorSpace_schwartz` (for $\hbar \neq 0$)
Example 3 in n dimensions, commutation relations	`posOpN_comm`, `momOpN_comm`, `posOpN_comm_momOpN`
Example 3 in n dimensions, symmetric observables	`isSymmetric_obsN`
Example 3 in n dimensions, Remark: the topology is that of $\mathcal{S}$	`topology_schwartzN` (for $\hbar \neq 0$)
Example 3 in n dimensions, second countability	`secondCountableTopology_schwartzN`
Example 3 in n dimensions is a quantum vector space	`isQuantumVectorSpace_schwartzN` (for $\hbar \neq 0$)
Theorem 5, [B]COBS	`IsAdmissible.continuous_expectation`
Sec. 5, a countable subset of observables generates the topology	`exists_countable_topology_eq`, `exists_countable_topology_eq_iff` (under first countability)
Sec. 5, the same for a quantum vector space	`IsQuantumVectorSpace.exists_countable`
Sec. 5, separability is not enough	`Diagonal.separableSpace`, `Diagonal.not_firstCountableTopology`, `Diagonal.not_exists_countable`

Differences from the brief.

- "Coarsest" is `IsGreatest` in Mathlib's order on topologies, where coarser is larger.
- The inner product is only assumed positive semidefinite in Corollary 2; definiteness is used only in finite dimension, to make the topology Hausdorff.
- Example 3 is formalized twice. For $\mathcal{S}(\mathbb{R})$, the proof that the topologies agree has two halves. (a) Each seminorm $\|x^a\partial^b f\|_2$ equals $|\hbar|^{-b}p_{X^aP^b}(f)$. (b) Every element of the algebra is a combination of the normally ordered monomials X^aP^b (`algebra_le_monoSpan`, from $PX^{n+1} = X^{n+1}P - (n+1)i\hbar X^n$), so every p_A is bounded by finitely many $\|x^a\partial^b f\|_2$. With `withSeminorms_l1_l2` from the previous brief, this gives the Schwartz topology.

- For $\mathcal{S}(\mathbb{R}^n)$, one half is that the Schwartz topology is admissible (`isAdmissible_schwartzN`), hence finer than the topology of Corollary 2. For the other half, each Schwartz seminorm $\sup_x \|x\|^k \|D^m f(x)\|$ is bounded by finitely many p_A (`exists_seminorm_le`). The derivative $D^m f(x)$ is bounded by its values on the unit vectors, $\|x\|^k \le (1+\|x\|^2)^k$ is the polynomial $(1+\sum_i (X^i)^2)^k$, and the Sobolev bound $|g(x)| \le C(\|g\|_2 + \sum_i \|\partial_i^{n+1} g\|_2)$ turns the supremum into L^2 norms of elements of the algebra applied to f. No normal ordering is needed, since the operators $\partial_i^{n+1}(1+\sum_j (X^j)^2)^k \partial^r$ are themselves in the algebra.

- In finite dimension the topology of Corollary 2 is the unique Hausdorff vector-space topology, whatever the defining observables (`topology_eq_moduleTopology`). This is how Example 4 is checked.

- Second countability in n dimensions comes from a general fact (`secondCountableTopology_topology`). With countably many defining observables, the p_A for the words $A = X_{i_1} \cdots X_{i_k}$ already generate the topology, so $v \mapsto (Av)_A$ embeds V into a countable product of copies of L^2.

- Second countability of $\mathcal{S}(\mathbb{R})$ is proved by embedding it into a countable product of copies of $L^2(\mathbb{R})$ through $f \mapsto (x^a \partial^b f)_{a,b}$ (`l2Embedding`). The topology of $\mathcal{S}(\mathbb{R})$ is induced by this map, because it is given by the L^2 seminorms (`withSeminorms_l1_l2`).

- Completeness of $\mathcal{S}(E,F)$, for any complete F, is proved in `ForMathlib`, since neither Mathlib (at this pin or on master) nor TauCeti nor physlib has it. The proof is the classical one (Becnel and Sengupta, *Mathematics* 3 (2015), §3.5): along a Cauchy sequence the weighted derivatives $\|x\|^k D^n u_m$ converge uniformly; the limits are the derivatives of the limit function, by the theorem on uniform limits of derivatives (`hasFDerivAt_of_tendstoUniformly`); and the bounds pass to the limit. For the qubit, completeness and second countability come from finite dimension.

- The claim of Sec. 5 that a countable subset of the observables generates the topology is proved under first countability, which second countability implies; it is in fact equivalent to first countability (`exists_countable_topology_eq_iff`). The proof (`WithSeminorms.exists_countable`) takes a countable basis of balls at 0; each ball involves finitely many words $X_{i_1} \cdots X_{i_k}$, and their letters form the countable subset. For a quantum vector space the subset is enlarged to be closed under the structure constants, so that it satisfies Definition 1 itself (`IsQuantumVectorSpace.exists_countable`).

- Separability is not enough. On the finitely supported sequences, with the ℓ^2 inner product and every multiplication operator $(M_f v)_n = f_n v_n$ by a real sequence f as a defining observable, the observables are symmetric and commute (`Diagonal.isSymmetric_mulOp`, `Diagonal.mulOp_comm`), and the space is separable, as the union of countably many finite-dimensional subspaces (`Diagonal.separableSpace`). But the topology is not first countable: for countably many balls U_n at 0, choose $t_n > 0$ with $t_n e_n \in U_n$; then $\|M_f(t_n e_n)\| = 1$ for $f_n = 1/t_n$, so no U_n lies in the ball of p_{M_f} (`Diagonal.not_firstCountableTopology`). Hence no countable subset of the observables generates the topology (`Diagonal.not_exists_countable`). The topology is the finest locally convex one, so the space is also complete; this is not formalized.

- Not formalized: the "iff $n = m$" part of Example 3, which needs a notion of isomorphism of quantum vector spaces; [B]WDQ and [B]STDIST in Theorem 5 and the other remarks of section 5, which are informal.

Problems found in the brief. One claim is false, and two steps are left implicit.

- Sec. 5 says that "separability/second countability would allow us to find a countable subset of observables from a bigger set that generates the same topology". This holds for second countability (first countability suffices), but not for separability: see the counterexample under "Differences".
- The Remark after Example 3 says the L^2 seminorms and the sup seminorms dominate each other. But the seminorms of Corollary 2 are $\|Af\|$ for every A in the algebra, including non-normally-ordered products such as PX. Reducing them to $\|x^a\partial^b f\|$ needs the commutation relation (normal ordering), which the brief does not mention.
- Example 3 needs $\hbar \neq 0$: for $\hbar = 0$ we get $P = 0$, and the topology is just that of L^2.