Lean formalization of an Assumptions of Physics technical brief: log

This log records which statements of the technical brief in this directory were checked in Lean, where the formal version differs from the brief, and any problems found in the brief. The brief is available at `https://assumptionsofphysics.org/resources/briefs`.

Setup

- Lean `v4.36.0-rc2`; Mathlib at commit `1b1ccf4b`; physlib (`master`) and TauCeti (commit `e7e7f52`, on the same Mathlib) as further dependencies.
- Build: `lake build` in the project root.
- Every file is checked with `#print axioms` on its main results. A result is recorded here only if it depends on no axioms beyond `propext`, `Classical.choice` and `Quot.sound`, and so has no `sorry`.
- Imports are kept to the Mathlib modules actually needed.
- Material that belongs in Mathlib is kept in `ForMathlib/`, in Mathlib's directory layout (library `ForMathlib`), and imported by the brief files.

1 Inconsistent convergence in Hilbert space

Files.

- File: `briefs/InconsistentConvergence/InconsistentConvergence.lean`
- Library: `InconsistentConvergence`
- Imports: `Mathlib.Analysis.Normed.Lp.lpSpace`
- Second file, for the oscillator: module `InconsistentConvergence.Oscillator`, in the subdirectory `InconsistentConvergence/`. It is separate from the main file, so that the comparator lane, which certifies the main file, is unchanged. Its imports:
  - `InconsistentConvergence`;
  - `ForMathlib.Analysis.Distribution.SchwartzSpace.HermiteReal`;
  - `TauCeti.Analysis.SpecialFunctions.Hermite.Function.Oscillator`.

  The `ForMathlib` file identifies $\mathcal{S}(\mathbb{R})$, with its topology, with the rapidly decreasing Hermite series. It transfers the multi-index Hermite expansions on $\mathbb{R}^1$, in the same directory, to $\mathbb{R}$.

- Third file, for the operators: module `InconsistentConvergence.NumberOperator`, in the same subdirectory and separate from the main file for the same reason. Its imports:
  - `InconsistentConvergence`;
  - `ForMathlib.Analysis.InnerProductSpace.l2Diagonal`;
  - `ForMathlib.LinearAlgebra.LinearPMap`.

  The `ForMathlib` files provide the diagonal operators on ℓ^2, self-adjoint on their maximal domains, and the composition of unbounded operators on its natural domain.

Setting. $H = \ell^2(\mathbb{N})$ (`lp (fun _ : ℕ => ℂ) 2`), with $|n\rangle = \delta_n$ the oscillator eigenstates and

$$|e^j\rangle = \sqrt{\tfrac{j-1}{j}}\,|0\rangle + \sqrt{\tfrac{1}{j}}\,|j\rangle\,.$$

In the oscillator file, U is the unitary $\ell^2(\mathbb{N}) \to L^2(\mathbb{R})$ of the Hermite basis (`TauCeti.hermiteHilbertBasis`), with $U\,|n\rangle = \psi_n$, the n-th Hermite function ($m = \omega = \hbar = 1$). $S(\mathbb{N})$ is described by its seminorms $\sup_n (n+1)^K |x_n|$, $K \in \mathbb{N}$.

Results.

Brief	Lean
eq. (2): $\|e^j - \|0\rangle\|^2 = \left(\sqrt{(j-1)/j} - 1\right)^2 + 1/j$	`norm_sub_ground_sq`
$e^j \to \|0\rangle$ in ℓ^2	`tendsto_e`
eq. (3): $\sum_n n^2 \|e^j_n\|^2 = j \to \infty$	`secondMoment_e`, `tendsto_secondMoment_e`
the $\|j\rangle$ alone have no limit in ℓ^2	`not_cauchySeq_level`: $\|\,\|j\rangle - \|k\rangle\,\|^2 = 2$ for $j \neq k$
eq. (4): $\langle e^j \|N\| e^j\rangle = 1$ for $j \geq 1$	`numberExp_e`
eq. (5), the inconsistency	`not_continuousAt_numberExp`: $\langle N\rangle$ is not continuous at $\|0\rangle$ in ℓ^2

Results: the oscillator.

Brief	Lean
$U\,\|n\rangle = \psi_n$	`hermiteUnitary_level`; the coordinates of $U^{-1}g$ are $\langle\psi_n\|g\rangle$ (`hermiteUnitary_symm_apply`)
$\|n\rangle$ is the oscillator eigenstate with n quanta	`oscillator_level`: $-\psi_n''/2 + x^2\psi_n/2 = (n+\frac{1}{2})\psi_n$
$S(\mathbb{N})$ is the Schwartz space	`exists_schwartz_iff`: $Ux \in \mathcal{S}(\mathbb{R})$ iff $\sup_n (n+1)^K \|x_n\| < \infty$ for every K
and has its topology	`tendsto_schwartz_iff`: $f_j \to g$ in $\mathcal{S}(\mathbb{R})$ iff $U^{-1}f_j \to U^{-1}g$ in every seminorm of $S(\mathbb{N})$
the e^j are Schwartz states	`exists_schwartz_e`: $Ue^j \in \mathcal{S}(\mathbb{R})$
$e^j \to \|0\rangle$, but not in $S(\mathbb{N})$	`tendsto_hermiteUnitary_e`: $Ue^j \to \psi_0$ in $L^2(\mathbb{R})$; `not_tendsto_schwartz_e`: Ue^j has no limit in $\mathcal{S}(\mathbb{R})$

Results: the number operator. In the operator file, a and $a^\dagger$ act by $(ax)_n = \sqrt{n+1}\, x_{n+1}$, $(a^\dagger y)_0 = 0$ and $(a^\dagger y)_{n+1} = \sqrt{n+1}\, y_n$, and N by $(Nx)_n = n\, x_n$, each on its maximal domain: $D(a) = \{x \mid (\sqrt{n}\, x_n)_n \in \ell^2\}$ and $D(N) = \{x \mid (n\, x_n)_n \in \ell^2\}$.

Brief	Lean
$a\,\lvert n+1\rangle = \sqrt{n+1}\,\lvert n\rangle$, $a\,\lvert 0\rangle = 0$, $a^\dagger\,\lvert n\rangle = \sqrt{n+1}\,\lvert n+1\rangle$	`annihilation_level_succ`, `annihilation_level_zero`, `creation_level`
$a^\dagger$ is the adjoint of a	`adjoint_annihilation`
N is self-adjoint, with $N\,\lvert n\rangle = n\,\lvert n\rangle$	`isSelfAdjoint_numberOp`, `numberOp_level`; domain: `mem_domain_numberOp`
$N = a^\dagger a$	`numberOp_eq_adjoint_compMax`, with the composition on $\{x \in D(a) \mid ax \in D(a^\dagger)\}$
$\langle x\lvert N\rvert x\rangle = \sum_n n\,\lvert x_n\rvert^2$	`inner_numberOp`, on $D(N)$; `numberExp_eq_norm_sq`: $\sum_n n\,\lvert x_n\rvert^2 = \lVert ax\rVert^2$ on $D(a)$
eq. (4) for the operator	`inner_numberOp_e`: $e^j \in D(N)$ and $\langle e^j\lvert N\rvert e^j\rangle = 1$ for $j \geq 1$

Differences from the brief.

- Indexing. Proposition 1 indexes sequences from 1, with the ground term at $i = 1$. The formalization indexes from 0, matching the oscillator of Section 3. The divergence is the same; with the brief's indexing the sum is $(j-1)/j + j$ instead of j.
- No limit in $S(\mathbb{N})$. In the main file, what is proved is that the seminorm $\sum_n n^2 |x_n|^2$ is unbounded along e^j, which is the brief's argument. The Schwartz sequence space $S(\mathbb{N})$ is not constructed as a topological space. The oscillator file describes it by its seminorms $\sup_n (n+1)^K |x_n|$, identifies it with $\mathcal{S}(\mathbb{R})$ through U, and proves that Ue^j has no limit in $\mathcal{S}(\mathbb{R})$.
- In the main file, $\langle N\rangle$ is defined on all of ℓ^2 as the weighted sum $\sum_n n\,\lvert x_n\rvert^2$, which has a default value outside the domain of N. The discontinuity is still meaningful, because every e^j is finitely supported. The operator file constructs $N = a^\dagger a$ and shows that the weighted sum is $\langle x\lvert N\rvert x\rangle$ on $D(N)$.

Problems found in the brief. No mathematical problems. Minor: the distance is written $d(x,y) = \sum_i (x_i - y_i)^2$, which is the squared distance and, for complex entries, should read $\sum_i |x_i - y_i|^2$.