

## Exercises on the four fundamental subspaces

**Problem 10.1:** (3.6 #11. *Introduction to Linear Algebra*: Strang)  $A$  is an  $m$  by  $n$  matrix of rank  $r$ . Suppose there are right sides  $\mathbf{b}$  for which  $A\mathbf{x} = \mathbf{b}$  has no solution.

- a) What are all the inequalities ( $<$  or  $\leq$ ) that must be true between  $m$ ,  $n$ , and  $r$ ?
- b) How do you know that  $A^T \mathbf{y} = \mathbf{0}$  has solutions other than  $\mathbf{y} = \mathbf{0}$ ?

**Solution:**

- a) The rank of a matrix is always less than or equal to the number of rows and columns, so  $r \leq m$  and  $r \leq n$ . The second statement tells us that the column space is not all of  $\mathbb{R}^n$ , so  $r < n$ .
- b) These solutions make up the left nullspace, which has dimension  $m - r > 0$  (that is, there are nonzero vectors in it).

**Problem 10.2:** (3.6 #24.)  $A^T \mathbf{y} = \mathbf{d}$  is solvable when  $\mathbf{d}$  is in which of the four subspaces? The solution  $\mathbf{y}$  is unique when the \_\_\_\_\_ contains only the zero vector.



**Solution:** It is solvable when  $\mathbf{d}$  is in the row space, which consists of all vectors  $A^T \mathbf{y}$ . The solution  $\mathbf{y}$  is unique when the left nullspace contains only the zero vector.



MIT OpenCourseWare  
<http://ocw.mit.edu>

18.06SC Linear Algebra  
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.