

CMPSC 465

Data Structures and Algorithms

Spring 2022

Instructor: Chunhao Wang

Linear Programming

(Textbook, Section 7.1)

Please consider taking

CMPSC 497 — Quantum Computation in Fall 2022

if you are interested in learning **Quantum Computing**

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$,
subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$, subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

- If no structures of f or C are known: general purpose constraint optimization

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$, subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

- If no structures of f or C are known: general purpose constraint optimization
- Given some restrictions on f or C , e.g., f is convex and C is a convex region: convex optimization

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$, subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

- If no structures of f or C are known: general purpose constraint optimization
- Given some restrictions on f or C , e.g., f is convex and C is a convex region: convex optimization
- Lots of stuff in between: quadratic programming, 2nd-order cone programming (SOCP), semidefinite programming (SDP)

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$, subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

- If no structures of f or C are known: general purpose constraint optimization
- Given some restrictions on f or C , e.g., f is convex and C is a convex region: convex optimization
- Lots of stuff in between: quadratic programming, 2nd-order cone programming (SOCP), semidefinite programming (SDP)
- Simplest non-trivial (but still powerful) case: f and C are linear functions, e.g., $f(\mathbf{x}) = a_1x_1 + a_2x_2 + \cdots + a_nx_n$

Background

Optimization: we want to maximize some function $f(\mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^n$, subject to constraints

$$C(\mathbf{x}) \leq \mathbf{b}, \text{ for } \mathbf{b} \in \mathbb{R}^n$$

- If no structures of f or C are known: general purpose constraint optimization
- Given some restrictions on f or C , e.g., f is convex and C is a convex region: convex optimization
- Lots of stuff in between: quadratic programming, 2nd-order cone programming (SOCP), semidefinite programming (SDP)
- Simplest non-trivial (but still powerful) case: f and C are linear functions, e.g., $f(\mathbf{x}) = a_1x_1 + a_2x_2 + \cdots + a_nx_n$
— **Linear Programming**

Example

Resource allocation: 168 hours in a week

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to pass classes: $S \geq 60$

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to pass classes: $S \geq 60$
- to stay sane: $P + E \geq 70$

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to pass classes: $S \geq 60$
- to stay sane: $P + E \geq 70$
- $2S + E - 3P \geq 150$: need more study time if had too much fun or not enough sleep

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to to pass classes: $S \geq 60$
- to stay sane: $P + E \geq 70$
- $2S + E - 3P \geq 150$: need more study time if had too much fun or not enough sleep
- happiness: $2P + E$ **objective function**

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to pass classes: $S \geq 60$
- to stay sane: $P + E \geq 70$
- $2S + E - 3P \geq 150$: need more study time if had too much fun or not enough sleep
- happiness: $2P + E$ **objective function**
i.e., $f(S, P, E) = 2P + E$

Example

Resource allocation: 168 hours in a week

S : study time; P : fun/party time; E : everything else

- to survive: $E \geq 56$
- to pass classes: $S \geq 60$
- to stay sane: $P + E \geq 70$
- $2S + E - 3P \geq 150$: need more study time if had too much fun or not enough sleep
- happiness: $2P + E$ **objective function**
i.e., $f(S, P, E) = 2P + E$

How to allocate your time?

Maximize happiness: LP formulation:

Maximize happiness: LP formulation:

$$\text{maximize} \quad 2P + E$$

LP formulation

Maximize happiness: LP formulation:

$$\begin{array}{ll}\text{maximize} & 2P + E \\ \text{subject to} & \end{array}$$

LP formulation

Maximize happiness: LP formulation:

$$\begin{array}{ll} \text{maximize} & 2P + E \\ \text{subject to} & E \geq 56 \end{array}$$

Maximize happiness: LP formulation:

$$\begin{array}{ll} \text{maximize} & 2P + E \\ \text{subject to} & E \geq 56 \\ & S \geq 60 \end{array}$$

LP formulation

Maximize happiness: LP formulation:

$$\begin{array}{ll} \text{maximize} & 2P + E \\ \text{subject to} & E \geq 56 \\ & S \geq 60 \\ & 2S + E - 3P \geq 150 \end{array}$$

LP formulation

Maximize happiness: LP formulation:

$$\begin{array}{llll} \text{maximize} & 2P + E & & \\ \text{subject to} & E \geq 56 & & \\ & S \geq 60 & & \\ & 2S + E - 3P \geq 150 & & \\ & S, P, E \geq 0 & & \end{array}$$

LP formulation

Maximize happiness: LP formulation:

$$\begin{array}{llll} \text{maximize} & 2P + E & & \\ \text{subject to} & E \geq 56 & & \\ & S \geq 60 & & \\ & 2S + E - 3P \geq 150 & & \\ & S, P, E \geq 0 & & \\ & S + P + E \leq 168 & & \end{array}$$

How to solve an LP


Consider a simpler LP:

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$

How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:


$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



How to solve an LP

Consider a simpler LP:

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



Optimal solution: $x + 2y = 70$

Algorithm for solving LP

Observation: (search for an optimal solution)

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex.

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists.

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary.

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary. At the boundary, moving in any direction will

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary. At the boundary, moving in any direction will

(a) Decrease objective function \rightarrow don't go this way

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary. At the boundary, moving in any direction will

- (a) Decrease objective function \rightarrow don't go this way
- (b) Increase objective function \rightarrow follow to a vertex

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary. At the boundary, moving in any direction will

- (a) Decrease objective function \rightarrow don't go this way
- (b) Increase objective function \rightarrow follow to a vertex
- (c) Objective function stays constant \rightarrow follow to a vertex

Algorithm for solving LP

Observation: (search for an optimal solution)

Objective function is linear, and feasible region is convex. So a unique direction of maximal increase of objective function exists. Follow it and you will run into the boundary. At the boundary, moving in any direction will

- (a) Decrease objective function \rightarrow don't go this way
- (b) Increase objective function \rightarrow follow to a vertex
- (c) Objective function stays constant \rightarrow follow to a vertex

Theorem

For an LP with bounded, nonempty feasible region, the maximum value will be attained at some vertex of the feasible region

Algorithm idea

The hill climbing approach (the simplex method)

Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function


$$\begin{array}{ll} \text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0 \end{array}$$

Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$




Algorithm idea

The hill climbing approach (the simplex method)

Start at a vertex, look at adjacent vertices, move in the direction of largest increase to the objective function

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$



Standard forms

LP solvers, such as MOSEK, Gurobi, CVX, and COIN are implementations of the simplex method. They require the LP to be in certain standard form

Standard forms

LP solvers, such as MOSEK, Gurobi, CVX, and COIN are implementations of the simplex method. They require the LP to be in certain standard form

Standard form 1

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

Standard forms

LP solvers, such as MOSEK, Gurobi, CVX, and COIN are implementations of the simplex method. They require the LP to be in certain standard form

Standard form 1

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

Example:

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array}$$

Standard forms

LP solvers, such as MOSEK, Gurobi, CVX, and COIN are implementations of the simplex method. They require the LP to be in certain standard form

Standard form 1

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

Example:

$$\begin{array}{ll}\text{maximize} & x + 2y \\ \text{subject to} & x \leq 20 \\ & y \leq 30 \\ & x + y \leq 40 \\ & x, y \geq 0\end{array} \quad \equiv \quad \begin{array}{ll}\text{maximize} & (1, 2) \begin{pmatrix} x \\ y \end{pmatrix} \\ \text{subject to} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \leq \begin{pmatrix} 20 \\ 30 \\ 40 \end{pmatrix} \\ & x, y \geq 0\end{array}$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \max & -\mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array}$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \max & -\mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array}$$

- Equality to inequality

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 = 7 \end{array} \quad \equiv$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \max & -\mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array}$$

- Equality to inequality

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 = 7 \end{array} \quad \equiv \quad \begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 \leq 7 \\ & x_1 + x_2 \geq 7 \end{array}$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \max & -\mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array}$$

- Equality to inequality

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 = 7 \end{array} \quad \equiv \quad \begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 \leq 7 \\ & x_1 + x_2 \geq 7 \end{array}$$

- Wrong inequality direction

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 \geq 7 \end{array} \quad \equiv$$

Convert to the standard form

- Minimization to maximization

$$\begin{array}{ll} \min & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array} \quad \equiv \quad \begin{array}{ll} \max & -\mathbf{c}^T \mathbf{x} \\ \text{s. t.} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq 0 \end{array}$$

- Equality to inequality

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 = 7 \end{array} \quad \equiv \quad \begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 \leq 7 \\ & x_1 + x_2 \geq 7 \end{array}$$

- Wrong inequality direction

$$\begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & x_1 + x_2 \geq 7 \end{array} \quad \equiv \quad \begin{array}{ll} \max & \mathbf{c}^T \mathbf{x} \\ \text{s. t.} & -x_1 - x_2 \leq -7 \end{array}$$

- Missing nonnegative constraints

$$\begin{array}{llll} \max & x_1 + 2x_2 & & \\ \text{s. t.} & x_1 & \leq & 20 \\ & x_1 + x_2 & \leq & 40 \\ & x_1 & \geq & 0 \end{array} \quad \equiv$$

- Missing nonnegative constraints

$$\begin{array}{ll}
 \max & x_1 + 2x_2 \\
 \text{s. t.} & x_1 \leq 20 \\
 & x_1 + x_2 \leq 40 \\
 & x_1 \geq 0
 \end{array}
 \quad \equiv \quad
 \begin{array}{ll}
 \max & x_1 + 2(x_2^+ - x_2^-) \\
 \text{s. t.} & x_1 \leq 20 \\
 & x_1 + (x_2^+ - x_2^-) \leq 40 \\
 & x_1 \geq 0 \\
 & x_2^+ \geq 0 \\
 & x_2^- \geq 0
 \end{array}$$

rewrite $x_2 = x_2^+ - x_2^-$

Another Standard form

Standard form 2

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} = \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

Another Standard form

Standard form 2

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} = \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

- Inequality to equality: use **slack variables**

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & x_1 \leq 20 \\ & x_1 \geq 0\end{array}$$

Another Standard form

Standard form 2

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} = \mathbf{b} \\ & \mathbf{x} \geq 0 \\ & \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}\end{array}$$

- Inequality to equality: use **slack variables**

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & x_1 \leq 20 \\ & x_1 \geq 0\end{array}$$

20 is bigger than x_1 by some positive amount, call it s

Another Standard form

Standard form 2

$$\begin{aligned} & \text{maximize} && \mathbf{c}^T \mathbf{x} \\ & \text{subject to} && A\mathbf{x} = \mathbf{b} \\ & && \mathbf{x} \geq 0 \\ & && \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n} \end{aligned}$$

- Inequality to equality: use **slack variables**

$$\begin{array}{lll} \text{maximize} & \mathbf{c}^T \mathbf{x} & \\ \text{subject to} & x_1 \leq 20 & \equiv \\ & x_1 \geq 0 & \end{array} \quad \begin{array}{ll} \text{maximize} & \mathbf{c}^T \mathbf{x} + 0 \cdot s \\ \text{subject to} & x_1 + s = 20 \\ & x_1 \geq 0 \\ & s \geq 0 \end{array}$$

20 is bigger than x_1 by some positive amount, call it s

Another Standard form

Standard form 2

$$\begin{aligned} & \text{maximize} && \mathbf{c}^T \mathbf{x} \\ & \text{subject to} && A\mathbf{x} = \mathbf{b} \\ & && \mathbf{x} \geq 0 \\ & && \mathbf{x}, \mathbf{c}, \mathbf{b} \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n} \end{aligned}$$

- Inequality to equality: use **slack variables**

$$\begin{array}{lll} \text{maximize} & \mathbf{c}^T \mathbf{x} & \\ \text{subject to} & x_1 \leq 20 & \equiv \\ & x_1 \geq 0 & \end{array} \quad \begin{array}{ll} \text{maximize} & \mathbf{c}^T \mathbf{x} + 0 \cdot s \\ \text{subject to} & x_1 + s = 20 \\ & x_1 \geq 0 \\ & s \geq 0 \end{array}$$

20 is bigger than x_1 by some positive amount, call it s

The new variable s is call the *slack variable*

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute `shortest_path(s, t)` for given $s, t \in V$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute `shortest_path`(s, t) for given $s, t \in V$

Can we model this as an LP?

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute `shortest_path`(s, t) for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute $\text{shortest_path}(s, t)$ for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute `shortest_path`(s, t) for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

i.e., d_v is the largest value s.t. $d_v \leq d_u + w(u, v)$ for all $(u, v) \in E$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute $\text{shortest_path}(s, t)$ for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

i.e., d_v is the largest value s.t. $d_v \leq d_u + w(u, v)$ for all $(u, v) \in E$

So we have

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute $\text{shortest_path}(s, t)$ for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

i.e., d_v is the largest value s.t. $d_v \leq d_u + w(u, v)$ for all $(u, v) \in E$

So we have

$$\begin{array}{ll} \text{minimize} & d_t \\ \text{subject to} & d_v \leq d_u + w(u, v) \quad \forall (u, v) \in E \\ & d_s = 0 \end{array}$$

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute $\text{shortest_path}(s, t)$ for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

i.e., d_v is the largest value s.t. $d_v \leq d_u + w(u, v)$ for all $(u, v) \in E$

So we have

$$\begin{array}{ll} \text{minimize} & d_t \\ \text{subject to} & d_v \leq d_u + w(u, v) \quad \forall (u, v) \in E \\ & d_s = 0 \end{array}$$

There are $|V|$ variables, $|E|$ constraints

Applications of LP — shortest path

We are given $G = (V, E)$, $w : E \rightarrow \mathbb{R}$

Want to compute $\text{shortest_path}(s, t)$ for given $s, t \in V$

Can we model this as an LP?

Recall Bellman-Ford: we calculate d_v for all $v \in V$, s.t. $d_v \leq d_u + w(u, v)$

So we had the greatest lower bound: $d_v = \min_{u \text{ s.t. } (u,v) \in E} \{d_u + w(u, v)\}$

i.e., d_v is the largest value s.t. $d_v \leq d_u + w(u, v)$ for all $(u, v) \in E$

So we have

$$\begin{array}{ll} \text{minimize} & d_t \\ \text{subject to} & d_v \leq d_u + w(u, v) \quad \forall (u, v) \in E \\ & d_s = 0 \end{array}$$

There are $|V|$ variables, $|E|$ constraints

We just **reduced** shortest_path to LP

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u, v) = \sum_{(v,w) \in E} f(v, w)$

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u, v) = \sum_{(v,w) \in E} f(v, w)$

LP formulation

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u, v) = \sum_{(v,w) \in E} f(v, w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u, v) = \sum_{(v,w) \in E} f(v, w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

We just **reduced** max_flow to LP

Application of LP — network flow


We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$. Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u,v) = \sum_{(v,w) \in E} f(v,w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

We just **reduced** max_flow to LP



Application of LP — network flow


We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u,v) = \sum_{(v,w) \in E} f(v,w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

We just **reduced** max_flow to LP



Application of LP — network flow


We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u,v) = \sum_{(v,w) \in E} f(v,w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

We just **reduced** max_flow to LP



Application of LP — network flow

We are given $G = (V, E)$, $s, t \in V$, capacity c_e for all $e \in E$ Find a flow $f : E \rightarrow \mathbb{R}^{\geq 0}$ s.t.

- $0 \leq f(e) \leq c_e$
- $\sum_{(u,v) \in E} f(u,v) = \sum_{(v,w) \in E} f(v,w)$

LP formulation

$$\begin{array}{ll} \max & \sum_{(s,u) \in E} f_{s,u} \\ \text{s.t.} & f_e \leq c_e \quad \forall e \in E \\ & \sum_{(u,v) \in E} f_{u,v} - \sum_{(v,w) \in E} f_{v,w} = 0 \\ & f_e \geq 0 \end{array}$$

We just **reduced** max_flow to LP

