

We need to show the following three things:

We need to show the following three things:

- Running time

We need to show the following three things:

- Running time
- FORD-FULKERSON outputs a flow

We need to show the following three things:

- Running time
- FORD-FULKERSON outputs a flow
- FORD-FULKERSON outputs the max flow

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

$$\begin{array}{cc} \frac{1}{3} & \frac{1}{4} \\ \downarrow & \downarrow \\ 4 & 3 \end{array}$$

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## **Fact (Fact 1)**

*In every step of the algorithm, the flow and the residual capacities are all integers*

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## Fact (Fact 1)

*In every step of the algorithm, the flow and the residual capacities are all integers*

## Fact (Fact 2)

*Let  $f$  be a flow in  $G$  and  $P$  be a simple  $s$ - $t$  path in  $G_f$ . Then*

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## Fact (Fact 1)

*In every step of the algorithm, the flow and the residual capacities are all integers*

## Fact (Fact 2)

*Let  $f$  be a flow in  $G$  and  $P$  be a simple  $s$ - $t$  path in  $G_f$ . Then*

$$v(f') = v(f) + \text{bottleneck}(P, f)$$



# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## Fact (Fact 1)

*In every step of the algorithm, the flow and the residual capacities are all integers*

## Fact (Fact 2)

*Let  $f$  be a flow in  $G$  and  $P$  be a simple  $s$ - $t$  path in  $G_f$ . Then*

$$v(f') = v(f) + \text{bottleneck}(P, f)$$

## Proof.

The first edge of  $P$  leaves  $s$ , and  $P$  doesn't revisit  $s$  again.

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## Fact (Fact 1)

*In every step of the algorithm, the flow and the residual capacities are all integers*

## Fact (Fact 2)

*Let  $f$  be a flow in  $G$  and  $P$  be a simple  $s$ - $t$  path in  $G_f$ . Then*

$$v(f') = v(f) + \text{bottleneck}(P, f)$$

## Proof.

The first edge of  $P$  leaves  $s$ , and  $P$  doesn't revisit  $s$  again. Moreover, it's a forward edge. So

# Running time analysis (I)

For simplicity, we assume the capacities are all integers

## Fact (Fact 1)

*In every step of the algorithm, the flow and the residual capacities are all integers*

## Fact (Fact 2)

*Let  $f$  be a flow in  $G$  and  $P$  be a simple  $s$ - $t$  path in  $G_f$ . Then*

$$v(f') = v(f) + \text{bottleneck}(P, f)$$

## Proof.

The first edge of  $P$  leaves  $s$ , and  $P$  doesn't revisit  $s$  again. Moreover, it's a forward edge. So

$$v(f) = \sum_{e \text{ out of } s} f(e)$$
$$v(f') = v(f) + \text{bottleneck}(P, f)$$

$f'(e) = f(e) + b$

□

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

### Corollary

*The Ford-Fulkerson algorithm performs  $\leq C$  iterations*

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

### Corollary

*The Ford-Fulkerson algorithm performs  $\leq C$  iterations*

### Proof.

All capacities are integers. Every iteration increase the value by  $\geq 1$   $\square$



## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

### Corollary

*The Ford-Fulkerson algorithm performs  $\leq C$  iterations*

### Proof.

All capacities are integers. Every iteration increase the value by  $\geq 1$   $\square$

Finding an  $s$ - $t$  path takes  $O(|V| + |E|) = O(|E|)$  time (BFS)

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

### Corollary

*The Ford-Fulkerson algorithm performs  $\leq C$  iterations*

### Proof.

All capacities are integers. Every iteration increase the value by  $\geq 1$   $\square$

Finding an  $s$ - $t$  path takes  $O(|V| + |E|) = O(|E|)$  time (BFS)

Augmentation takes  $O(|V|)$  time

## Running time analysis (II)

Since  $\text{bottleneck}(P, f) \geq 1$ ,

$$v(f') \geq v(f) + 1$$

Let  $C = \sum_{e \text{ out of } s} c_e$ . We have

### Corollary

*The Ford-Fulkerson algorithm performs  $\leq C$  iterations*

### Proof.

All capacities are integers. Every iteration increase the value by  $\geq 1$   $\square$

Finding an  $s$ - $t$  path takes  $O(|V| + |E|) = O(|E|)$  time (BFS)

Augmentation takes  $O(|V|)$  time

So total running time is  $O(C \cdot |E|)$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

**Proof.**

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- Capacity constraint.

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$   
Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$



# Ford-Fulkerson outputs a flow

## Lemma

Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$

- if  $e$  is a forward edge, then  $\text{residual capacity: } c_e - f(e)$

$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

$$\text{So, } 0 \leq f'(e) \leq c_e$$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$   
Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$ 
  - if  $e$  is a forward edge, then
$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$

- if  $e$  is a forward edge, then

$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

$$\text{So } 0 \leq f'(e) \leq c_e$$

# Ford-Fulkerson outputs a flow

## Lemma

Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$

- if  $e$  is a forward edge, then

$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

$$\text{So } 0 \leq f'(e) \leq c_e$$

- if  $e$  is a backward edge, then

residual capacity:  $f(e) \geq \text{bottleneck}(P, f)$

$$\underline{c_e} \geq f(e) \geq \underline{f'(e) = f(e) - \text{bottleneck}(P, f)} \geq f(e) - f(e) = \underline{0}$$

$$\Rightarrow 0 \leq f'(e) \leq c_e$$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$

- if  $e$  is a forward edge, then

$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

So  $0 \leq f'(e) \leq c_e$

- if  $e$  is a backward edge, then

$$c_e \geq f(e) \geq f'(e) = f(e) - \text{bottleneck}(P, f) \geq f(e) - f(e) = 0$$

# Ford-Fulkerson outputs a flow

## Lemma

*Let  $f'$  be the function obtained after augmenting. Then  $f'$  is a flow*

## Proof.

- **Capacity constraint.** It suffices to consider edges of  $P$

Let  $e = (u, v) \in P$ .  $\text{bottleneck}(P, f)$  is at most the residual capacity of  $e$

- if  $e$  is a forward edge, then

$$0 \leq f(e) \leq f'(e) = f(e) + \text{bottleneck}(P, f) \leq f(e) + (c_e - f(e)) = c_e$$

$$\text{So } 0 \leq f'(e) \leq c_e$$

- if  $e$  is a backward edge, then

$$c_e \geq f(e) \geq f'(e) = f(e) - \text{bottleneck}(P, f) \geq f(e) - f(e) = 0$$

$$\text{So } 0 \leq f'(e) \leq c_e$$

- **Conservation condition.** It suffices to observe that for every vertex, additional amount of flow, 0, or  $\text{bottleneck}(P, f)$  entering this vertex equals the additional amount of flow, 0, or  $\text{bottleneck}(P, f)$  leaving it




# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition

An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$





# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition

An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$

### Definition

The **capacity** of the cut is

$$c(A, B) = \sum_{e \text{ out of } A} c_e$$



# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition

An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$

### Definition

The **capacity** of the cut is

$$c(A, B) = \sum_{e \text{ out of } A} c_e$$

How does a cut help?

# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition


An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$

### Definition

The **capacity** of the cut is

$$c(A, B) = \sum_{e \text{ out of } A} c_e$$

How does a cut help?  $c(A, B) = 22$



# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition


An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$

### Definition

The **capacity** of the cut is

$$c(A, B) = \sum_{e \text{ out of } A} c_e$$

How does a cut help?



The flow must have a value  $\leq 22$

# Correctness of Ford-Fulkerson (I)

## Flow and Cut

### Definition


An **s-t cut** is a partition of  $V$ ,  $(A, B)$  where  $s \in A$  and  $t \in B$

### Definition

The **capacity** of the cut is

$$c(A, B) = \sum_{e \text{ out of } A} c_e$$

How does a cut help?



The flow must have a value  $\leq 22$   
Capacity of a cut put a bound on the flow value

## Correctness of Ford-Fulkerson (II)

### Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

# Correctness of Ford-Fulkerson (II)

## Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

**Proof.**

# Correctness of Ford-Fulkerson (II)

## Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

**Proof.** Notation:



# Correctness of Ford-Fulkerson (II)

## Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

## Correctness of Ford-Fulkerson (II)

### Lemma

Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

$$f^{\text{in}}(A) = \sum_{e \text{ into } A} f(e)$$

$$v(f) = \sum_{e \text{ out of } s} f(e) = f^{\text{out}}(s) = f^{\text{out}}(s) - \underbrace{f^{\text{in}}(s)}_{=0}$$

## Correctness of Ford-Fulkerson (II)

### Lemma

Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

$$f^{\text{in}}(A) = \sum_{e \text{ into } A} f(e)$$

So,  $v(f) = \sum_{e \text{ out of } s} f(e) = f^{\text{out}}(s) = f^{\text{out}}(s) - f^{\text{in}}(s)$  (no edge into  $s$ )

for all  $v \in A - \{s, t\}$   $f^{\text{out}}(v) = f^{\text{in}}(v) \Rightarrow f^{\text{out}}(v) - f^{\text{in}}(v) = 0$

# Correctness of Ford-Fulkerson (II)

## Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

$$f^{\text{in}}(A) = \sum_{e \text{ into } A} f(e)$$

So,  $v(f) = \sum_{e \text{ out of } s} f(e) = f^{\text{out}}(s) = f^{\text{out}}(s) - f^{\text{in}}(s)$  (no edge into  $s$ )

Also, for all  $v \in A - \{s, t\}$ ,  $f^{\text{out}}(v) = f^{\text{in}}(v)$  (flow conservation)

## Correctness of Ford-Fulkerson (II)

### Lemma

Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

$$f^{\text{in}}(A) = \sum_{e \text{ into } A} f(e)$$

So,  $v(f) = \sum_{e \text{ out of } s} f(e) = f^{\text{out}}(s) = \underline{f^{\text{out}}(s) - f^{\text{in}}(s)}$  (no edge into  $s$ )

Also, for all  $v \in A - \{s, t\}$ ,  $f^{\text{out}}(v) = f^{\text{in}}(v)$  (flow conservation)

$\implies \underline{f^{\text{out}}(v) - f^{\text{in}}(v)} = 0$  for all  $v \neq s, t$

$$\begin{aligned} v(f) &= f^{\text{out}}(s) - f^{\text{in}}(s) \quad \text{to to to to to} \dots \text{to} \\ &= \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v)) \end{aligned}$$

## Correctness of Ford-Fulkerson (II)

### Lemma

*Let  $f$  be an  $s$ - $t$  flow,  $(A, B)$  be an  $s$ - $t$  cut. Then  $v(f) \leq c(A, B)$*

**Proof.** Notation:

$$f^{\text{out}}(A) = \sum_{e \text{ out of } A} f(e)$$

$$f^{\text{in}}(A) = \sum_{e \text{ into } A} f(e)$$

So,  $v(f) = \sum_{e \text{ out of } s} f(e) = f^{\text{out}}(s) = f^{\text{out}}(s) - f^{\text{in}}(s)$  (no edge into  $s$ )

Also, for all  $v \in A - \{s, t\}$ ,  $f^{\text{out}}(v) = f^{\text{in}}(v)$  (flow conservation)


$$\implies f^{\text{out}}(v) - f^{\text{in}}(v) = 0 \text{ for all } v \neq s, t$$

So

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

For this expression

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$



For this expression

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

consider every edge  $(v, w)$




For this expression

$$v(f) = \sum_{v \in A} (\underbrace{f^{\text{out}}(v)} - \underbrace{f^{\text{in}}(v)})$$

consider every edge  $(v, w)$

- if  $v, w \in A$ , this edge contributes 0 in the summation




For this expression

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

consider every edge  $(v, w)$

- if  $v, w \in A$ , this edge contributes 0 in the summation
- if  $v \in A, w \notin A$ , this edge contributes  $f(e)$



For this expression


$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

consider every edge  $(v, w)$

- if  $v, w \in A$ , this edge contributes 0 in the summation
- if  $v \in A, w \notin A$ , this edge contributes  $f(e)$
- if  $v \notin A, w \in A$ , this edge contributes  $-f(e)$

$$v(f) = \sum_{e \text{ out of } A} f(e) - \sum_{e \text{ into } A} f(e)$$

$$\leq \sum_{e \text{ out of } A} f(e) = C(A, B)$$



For this expression

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

consider every edge  $(v, w)$

- if  $v, w \in A$ , this edge contributes 0 in the summation
- if  $v \in A, w \notin A$ , this edge contributes  $f(e)$
- if  $v \notin A, w \in A$ , this edge contributes  $-f(e)$

We rewrite the summation as

For this expression

$$v(f) = \sum_{v \in A} (f^{\text{out}}(v) - f^{\text{in}}(v))$$

consider every edge  $(v, w)$

- if  $v, w \in A$ , this edge contributes 0 in the summation
- if  $v \in A, w \notin A$ , this edge contributes  $f(e)$
- if  $v \notin A, w \in A$ , this edge contributes  $-f(e)$

We rewrite the summation as

$$\underline{v(f)} = \sum_{e \text{ out of } A} f(e) - \sum_{e \text{ into } A} f(e) \leq \sum_{e \text{ out of } A} f(e) = \underline{c(A, B)}$$

$$v(f) \leq c(A, B)$$

□

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.**



## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ .

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ . We have the follow facts

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ . We have the follow facts

- $(A^*, B^*)$  is an  $s$ - $t$  cut:  $s \in A^*$ ,  $t \in B^*$  (no  $s$ - $t$  path in  $G_f$ )

## Correctness of Ford-Fulkerson (III)

The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ . We have the follow facts

- $(A^*, B^*)$  is an  $s$ - $t$  cut:  $s \in A^*$ ,  $t \in B^*$  (no  $s$ - $t$  path in  $G_f$ )
- for all edge  $e = (v, v') \in E$  with  $v \in A^*$ ,  $v' \in B^*$ , we have  $f(e) = c_e$

## Correctness of Ford-Fulkerson (III)


The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

### Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*

**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ . We have the follow facts

- $(A^*, B^*)$  is an  $s$ - $t$  cut:  $s \in A^*$ ,  $t \in B^*$  (no  $s$ - $t$  path in  $G_f$ )
- for all edge  $e = (v, v') \in E$  with  $v \in A^*$ ,  $v' \in B^*$ , we have  $f(e) = c_e$   
Otherwise,  $(v, v')$  is an edge in  $G_f$  with capacity  $c_e - f(e) \neq 0$ .



# Correctness of Ford-Fulkerson (III)


The upper bound  $c(A, B)$  is achievable by Ford-Fulkerson

## Lemma

*Let  $f$  be a flow s.t. there's no  $s$ - $t$  path in  $G_f$ . Then there exists an  $s$ - $t$  cut  $(A^*, B^*)$  s.t.  $v(f) = c(A^*, B^*)$*


**Proof.** Let  $A^*$  be the set of vertices reachable from  $s$  in  $G_f$ . Let  $B^*$  be  $V - A^*$ . We have the follow facts

- $(A^*, B^*)$  is an  $s$ - $t$  cut:  $s \in A^*$ ,  $t \in B^*$  (no  $s$ - $t$  path in  $G_f$ )
- for all edge  $e = (v, v') \in E$  with  $v \in A^*$ ,  $v' \in B^*$ , we have  $f(e) = c_e$   
Otherwise,  $(v, v')$  is an edge in  $G_f$  with capacity  $c_e - f(e) \neq 0$ .  
Forward edge. So  $v'$  is reachable from  $s$  in  $G_f$  (contradiction)




- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$

- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
 Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contradiction)






- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
 Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contradiction)



Then


- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contradiction)



Then

$$v(f) = \underbrace{\sum_{e \text{ out of } A^*} f(e)}_{c_e} - \underbrace{\sum_{e \text{ into } A^*} f(e)}_0 \quad (\text{from the proof of } v(f) \leq c(A, B))$$


- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contraction)



Then

$$\begin{aligned}
 v(f) &= \sum_{e \text{ out of } A^*} f(e) - \sum_{e \text{ into } A^*} f(e) \quad (\text{from the proof of } v(f) \leq c(A, B)) \\
 &= \sum_{e \text{ out of } A^*} f(e) - 0
 \end{aligned}$$


- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contraction)



Then

$$\begin{aligned}
 v(f) &= \sum_{e \text{ out of } A^*} f(e) - \sum_{e \text{ into } A^*} f(e) \quad (\text{from the proof of } v(f) \leq c(A, B)) \\
 &= \sum_{e \text{ out of } A^*} f(e) - 0 \\
 &= \sum_{e \text{ out of } A^*} c_e
 \end{aligned}$$

- for all edge  $e = (w', w)$  with  $w' \in B^*$ ,  $w \in A^*$ , we have  $f(e) = 0$   
Otherwise,  $(w, w')$  is a backward edge in  $G_f$ , and  $w'$  would be reachable from  $s$  (contradiction)



Then

$$\begin{aligned}
 v(f) &= \sum_{e \text{ out of } A^*} f(e) - \sum_{e \text{ into } A^*} f(e) \quad (\text{from the proof of } v(f) \leq c(A, B)) \\
 &= \sum_{e \text{ out of } A^*} f(e) - 0 \\
 &= \sum_{e \text{ out of } A^*} c_e = c(A^*, B^*)
 \end{aligned}$$

□

# Summary of Max flow

Some consequences:

# Summary of Max flow

Some consequences:

- The flow returned by Ford-Fulkerson is a maximum flow

# Summary of Max flow

Some consequences:

- The flow returned by Ford-Fulkerson is a maximum flow
- In every flow network, maximum value of a flow = minimum capacity of a cut



# Summary of Max flow

Some consequences:

- The flow returned by Ford-Fulkerson is a maximum flow
- In every flow network, maximum value of a flow = minimum capacity of a cut
- Given a flow of max value, can compute a cut of minimum capacity in  $O(|E|)$  time

# Summary of Max flow

Some consequences:

- The flow returned by Ford-Fulkerson is a maximum flow
- In every flow network, maximum value of a flow = minimum capacity of a cut
- Given a flow of max value, can compute a cut of minimum capacity in  $O(|E|)$  time
- If all capacities of a flow network are integers, then there is a max flow  $f$  s.t.  $f(e)$  is an integer for all  $e$