

A
PS-CVD5 + arm vs base (S32)
filled: q < .05; open: q ≥ .05
1:1, caliper 0.2 (main)
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
1:1, caliper 0.1
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
1:3 matching
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
IPTW
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
Overlap weights
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
Gradient-boosted PS
+ ECG
+ CLMBR-T
+ ECG + CLMBR-T
+ permuted ECG
+ permuted CLMBR-T
−20
0
20
Balance: relative reduction in
mean |SMD|, 54 items, % (95% CI)
B
−0.05
0.00
Agreement: change in mean
|Δ log HR| vs RCT (< 0: closer)
Closer P Perm. P q
20/32 .049 .70 .58
18/32 .030 .93 .51
18/32 .019 .94 .43
16/32 1.00 .16 1.00
19/32 .97 .29 1.00
16/32 .11 .85 .70
16/32 .13 .99 .71
20/32 .037 .99 .51
13/32 .96 .076 1.00
19/32 .95 .34 1.00
21/32 .007 .037 .43
19/32 .036 .89 .51
20/32 .020 .78 .43
16/32 .75 .040 1.00
16/32 .91 .19 1.00
15/32 .69 .96 1.00
17/32 .37 .94 .95
17/32 .38 .91 .95
14/32 .88 .60 1.00
17/32 .87 .95 1.00
20/32 .011 .20 .43
16/32 .13 .98 .74
19/32 .087 .96 .69
15/32 .91 .75 1.00
20/32 .12 .21 .70
17/32 .32 .005 .89
17/32 .25 .99 .81
19/32 .15 .84 .75
14/32 .58 .94 1.00
14/32 .27 .78 .83
C
All 337 S32 contrast rows. No agreement row reaches q < .05 in S32
(numbers: rows with q < .05 for balance | |Δ log HR| | benchmark permutation)
18 | 0 | 0
14 | 0 | 0
18 | 0 | 0
7 | 0 | 0
23 | 0 | 0
18 | 0 | 0
BH q (whole sweep, by family)
q = .05
main
caliper 0.1
1:3
IPTW
overlap
GBM PS
balance (mean |SMD|, 54)
agreement (|Δ log HR|)
benchmark permutation
D
label-perm. P: ECG .98,
CLMBR-T .87
quality tier:
descriptive only
Agreement gain per trial:
|Δ log HR| base − arm (> 0: closer)
Physio-
logic
Not
Excellent
/good
Moderate
6 trial points beyond ±0.3 drawn at the limit
+ ECG
+ CLMBR-T