
A
mean / median
PS-CVD5
PS-CVD5 + ECG
PS-CVD5 + CLMBR-T
PS-CVD5 + ECG + CLMBR-T
PS-Demo + CLMBR-T
PS-Flags (C = 1)
hdPS200
PS-Clinical
0.89 / 0.99
0.88 / 0.97
0.79 / 0.83
0.77 / 0.81
0.80 / 0.83
0.74 / 0.73
0.77 / 0.80
0.85 / 0.91
0.5 0.6 0.7 0.8 0.9 1.0
Matched share of the smaller arm, main
(bar: mean; tick: median over trials)
B
PS-Demo
PS-Demo + CLMBR-T
PS-CVD5
PS-CVD5 + ECG
PS-CVD5 + CLMBR-T
PS-Flags (C = 1)
hdPS200
PS-Clinical
IPTW
overlap
0.0 0.2 0.4 0.6 0.8 1.0
ESS fraction min(ESS_t/n_t, ESS_c/n_c)
(dot: median; tick: minimum over trials)
C
+ ECG: all 32 trials (main)
CLMBR-T retention drop ≤ 0.05 (15)
drop > 0.05 (17)
overlap weights, no trimming (32)
+ CLMBR-T: all 32 trials (main)
CLMBR-T retention drop ≤ 0.05 (15)
drop > 0.05 (17)
overlap weights, no trimming (32)
+ ECG + CLMBR-T: all 32 trials (main)
CLMBR-T retention drop ≤ 0.05 (15)
drop > 0.05 (17)
overlap weights, no trimming (32)
0 10 20
Relative reduction in mean |SMD| vs PS-CVD5, %
per-trial CLMBR-T gain vs its retention loss: Spearman ρ = 0.21, P = .24
D
agreement
perm. P
PS-Flags CV + CLMBR-T vs PS-Flags CV
main .16
1:3 .071
overlap .39
PS-Flags CV + ECG vs PS-Flags CV
main .052
1:3 .032
overlap .037
PS-Flags CV vs PS-Flags C = 1
main .008
1:3 .003
overlap .46
PS-Demo + CLMBR-T vs PS-Flags CV
main .001
1:3 <.001
overlap .70
−15 −10 −5 0 5 10
Relative reduction in mean |SMD|, 54 items, %
(sensitivity rows, outside the FDR sweep)