

MAT237Y1 - Interview Activity

1. *Claim:* Let $S \subseteq \mathbb{R}^n$ be a finite set. Prove that S has zero Jordan measure.

Solution: Fix $\varepsilon > 0$. Set $N = |S|$, the cardinality of the set. Label the elements of S as x_1, x_2, \dots, x_N . For $i \in \{1, \dots, N\}$, define the rectangles

$$R_i = \left[(x_i)_1 - \sqrt[n]{\frac{\varepsilon}{2^{n+1}N}}, (x_i)_1 + \sqrt[n]{\frac{\varepsilon}{2^{n+1}N}} \right] \times \dots \times \left[(x_i)_n - \sqrt[n]{\frac{\varepsilon}{2^{n+1}N}}, (x_i)_n + \sqrt[n]{\frac{\varepsilon}{2^{n+1}N}} \right] .$$

Notice that $x \in R_i$ by the construction of R_i and

$$\text{vol}(R_i) = \left(2 \sqrt[n]{\frac{\varepsilon}{2^{n+1}N}} \right)^n = \left(\sqrt[n]{\frac{\varepsilon}{2N}} \right)^n = \frac{\varepsilon}{2N} .$$

Therefore, $S \subseteq \bigcup_{i=1}^N R_i$ and

$$\sum_{i=1}^N \text{vol}(R_i) = N \frac{\varepsilon}{2N} = \frac{\varepsilon}{2} < \varepsilon$$

as needed.

The skills needed to produce this solution include:

- The ability to recall the definition of a set having zero Jordan measure, which in turn involves the definitions of rectangles, the volume of a rectangle, and the finite union and sum operations.
- The ability to match the structure of the solution with the format of the definition. The steps must be taken in this order for correctness: fix an arbitrary ε , set an appropriate N , construct specific rectangles depending on ε and N , then ensure the conditions of the definition are satisfied.
- The ability to recognize what a finite set is and how to construct a simple rectangle that uses the properties of such a set to meet the conditions of the definition.
- The ability to recognize and avoid potential sources of confusion. Above $(x_i)_1$ was used to avoid confusion with notation such as x_{i1}, \dots, x_{in} , where it would be difficult to distinguish between the different subscripts.