Circle the correct answer for each question. If you're unsure, you can write numbers 1, 2, 3, etc. next to the choices to get partial credit. Questions refer to the nonindented paragraph above unless otherwise stated.

According to the 2019 U.S. Census Bureau estimates, 63.4% of the U.S. self-identifies as Non-Hispanic white, 15.3% as Hispanic and Latino, 13.4% as Black or African American, 5.9% as Asian, and 2.0% as Other. A random sample of 1000 students graduating from California colleges and universities resulted in the following data on self-identified race:

Self-identified Race	Non-Hispanic White	Black or African American	Hispanic or Latino	Asian	Other
Number in Sample	679	51	77	190	3

Use this information to answer questions 1 and 2

1. In the null hypothesis, what does the symbol μ_{Race} represent?


   (a) The proportion of Californians that self-identify as a certain Race
   (b) The proportion of all students graduating from California colleges and universities that self-identify as a certain Race
   (c) The proportion of the U.S. that self-identify as a certain Race
   (d) The proportion of the sample of 1000 students that self-identified as a certain Race

2. Based upon a p-value of less than 0.001, what is an appropriate conclusion for the goodness-of-fit test?

   (a) Strong evidence exists that the distribution of self-identified race among California college and university students is the same as the distribution of self-identified race in the U.S. overall.
   (b) Strong evidence does not exist that the distribution of self-identified race among California college and university students differs from the distribution of self-identified race in the U.S. overall.
   (c) Strong evidence exists that the distribution of self-identified race among California college and university students differs from the distribution of self-identified race in the U.S. overall.
   (d) Strong evidence exists that the probabilities of each self-identified race among California college and university students are not all equal.

A British study examined whether the type of background music playing in a restaurant affected the amount of money that diners spent on their meals. The researchers asked a restaurant to alternate silence, popular music, and classical music on successive nights over eighteen days. Each type of music was played for 6 nights (the order was randomly determined to guard against confounding). The type of music played and the amount spent on food and drinks (in British pounds) for each of 393 customers was recorded. The data are summarized below.

	Sample size	Mean	StDev
Classical	120	24.1	2.30
Pop	142	21.9	2.73
None	131	21.7	3.38
Pooled	393	22.5	2.85



3. All of the following statements are true for this situation, but one is not a legitimate validity condition. Which one is not the real validity condition?

   (a) The sample sizes are all at least 20.
   (b) The standard deviations of the sample are approximately equal.
   (c) The sample means are all at least 20.
   (d) The sample distributions show no strong skewness.


4. The p-value for the ANOVA F-test is less than 0.001. How would you interpret this value?

   (a) In less than 0.1% of all samples, the F-statistic would be as large or larger than the one observed, assuming there is an association between music type and amount spent on food and drinks.
   (b) In less than 0.1% of all samples, the F-statistic would be as small or smaller than the one observed, assuming there is no association between music type and amount spent on food and drinks.
   (c) In less than 0.1% of all samples, the F-statistic would be as large or larger than the one observed, assuming there is no association between music type and amount spent on food and drinks.
   (d) In less than 0.1% of all samples, the F-statistic would be as small or smaller than the one observed, assuming there is an association between music type and amount spent on food and drinks.

5. An experiment is conducted which randomly places students into one of three treatment groups: (1) those students who will take a test that is printed on white paper, (2) those students who will take the same test, but printed on red paper, and (3) those who will take the same test, but printed on blue paper. The researcher analyzes the data using ANOVA in order to compare the average scores in the three groups looking for evidence that the average score is different on at least one of the three tests. Assume that at least one group really is different when answering the following question.
   As the size of the sample in each group increases, all else remaining the same, what will be the impact on the strength of evidence for finding a difference in the average scores on the three exams?

   (a) There will be less evidence of a difference.
   (b) There is not enough information to tell.
   (c) There will be more evidence of a difference.
   (d) There will be no impact on the strength of evidence of a difference.

6. The two graphs, A and B, show dotplots from two different data sets. Suppose I want to compare the means in A and do another comparison of means in B. For each case an F-statistic is computed. Which of the following will be true about these statistics?



   (a) The F-statistic is the same in A and B.
   (b) The F-statistic is larger in A than in B.
   (c) The F-statistic is smaller in A than in B.
   (d) We cannot tell if the F-statistic will be larger in A or B.

Are people equally likely to be born on any day of the seven days of the week? Or are some days more likely to be a person's birthday than other days? To investigate this question, days of birth were recorded for the 147 "noted writers of the present" listed in The World Almanac and Book of Facts 2000. The counts for the seven days of the week are given in the following table.

Mon	Tues	Wed	Thurs	Fri	Sat	Sun
17	26	22	23	18	15	25

Use this information to answer questions 7 and 8

7. Are validity conditions met to conduct a theory-based chi-square goodness-of-fit test?

   (a) No, since the p-values in questions 35 and 37 were different.
   (b) No, since this was not a random sample.
   (c) Yes, since we have at least 10 successes and 10 failures.
   (d) Yes, since all observed counts are at least 10.

8. Based upon a p-value of 0.53, what is the appropriate conclusion for the test?

   (a) Strong evidence does not exist that people are equally likely to be born on any day of the seven days of the week.
   (b) Strong evidence exists that people are not equally likely to be born on any day of the seven days of the week.
   (c) Strong evidence exists that people are equally likely to be born on any day of the seven days of the week.
   (d) Strong evidence does not exist that people are not equally likely to be born on any day of the seven days of the week.

9. When performing an ANOVA F-test, which definition of a p-value is the most accurate?

   (a) The p-value is the probability that the observed F-statistic will occur again.
   (b) If the null hypothesis is true, the p-value is the probability of observing an F-statistic as large or larger than the one actually observed.
   (c) The p-value is the probability that the null hypothesis is true.
   (d) The p-value is the value that an F-statistic must reach in order to be significant under the null hypothesis.