

## Step-1

(a)

Use row operations to simplify and compute the determinant.

Consider the determinant,

$$A = \begin{vmatrix} 101 & 201 & 301 \\ 102 & 202 & 302 \\ 103 & 203 & 303 \end{vmatrix}$$

Properties of determinants:

(i) If matrix,  $A$  has row of zero's, then  $\det A = 0$

(ii) Adding a multiple of one row to another row leaves same determinant.

(iii) If row of matrix,  $A$  is multiplied by constant,  $k$  resulting determinant is  $k|A|$

Subtract first row from second and third rows.

$$A \sim \begin{vmatrix} 101 & 201 & 301 \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{vmatrix}$$

Subtract 2 times of second row from third row.

$$= \begin{vmatrix} 101 & 201 & 301 \\ 1 & 1 & 1 \\ 0 & 0 & 0 \end{vmatrix}$$

$= 0$  Since third row of determinant has zero's

Therefore, the determinant value is  $\boxed{0}$ .

## Step-2

(b)

Consider the determinant,

$$\begin{vmatrix} 1 & t & t^2 \\ t & 1 & t \\ t^2 & t & 1 \end{vmatrix}$$

Subtract the first row from second row and third rows.

$$= \begin{vmatrix} 1 & t & t^2 \\ t-1 & 1-t & t-t^2 \\ t^2-1 & 0 & 1-t^2 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & t & t^2 \\ -(1-t) & 1-t & t(1-t) \\ -(1-t^2) & 0 & 1-t^2 \end{vmatrix}$$

$$= (1-t)(1-t^2) \begin{vmatrix} 1 & t & t^2 \\ -1 & 1 & t \\ -1 & 0 & 1 \end{vmatrix} \text{ Takeout common factors}$$

Add first row to second and third rows.

$$= (1-t)(1-t^2) \begin{vmatrix} 1 & t & t^2 \\ 0 & 1+t & t+t^2 \\ 0 & t & 1+t^2 \end{vmatrix}$$

Subtract third row from second row.

$$= (1-t)(1-t^2) \begin{vmatrix} 1 & t & t^2 \\ 0 & 1 & t-1 \\ 0 & t & 1+t^2 \end{vmatrix}$$

### Step-3

Subtract third row from second row.

$$= (1-t)(1-t^2) \begin{vmatrix} 1 & 0 & -1 \\ 0 & 1 & t-1 \\ 0 & t & 1+t^2 \end{vmatrix}$$

$$= (1-t)(1-t^2) [1(1+t^2) - t(t-1)]$$

$$= (1-t)(1-t^2)(1+t)$$

$$= \boxed{(1-t^2)}.$$