

Step-1

We have to find an orthonormal basis for the plane $x - y + z = 0$, and we have to find the matrix P that project onto the plane, and we have to find the nullspace of P .

Given equation is $x - y + z = 0$

Put $y = k, z = r$

$$\Rightarrow x = y - z$$

$$= k - r$$

Step-2

Therefore

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} k - r \\ k \\ r \end{bmatrix}$$
$$= k \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + r \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

$$a_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, a_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Therefore are two vectors in the plane $x - y + z = 0$

Step-3

$$q_1 = \frac{a_1}{\|a_1\|}$$
$$= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$
$$= \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$$

Step-4

$$q_2 = \frac{\beta}{\|\beta\|} \text{ where } \beta = a_2 - (q_1^T a_2) q_1$$

$$q_1^T a_2 = \frac{1}{\sqrt{2}} (1, 1, 0) \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$= \frac{1}{\sqrt{2}} (-1 + 0 + 0)$$

$$= -\frac{1}{\sqrt{2}}$$

Step-5

$$(q_1^T a_2) q_1 = -\frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$= -\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Step-6

$$\beta = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} + \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} -1/2 \\ 1/2 \\ 1 \end{bmatrix}$$

Step-7

$$\|\beta\| = \sqrt{\frac{1}{4} + \frac{1}{4} + 1}$$

$$= \sqrt{\frac{6}{4}}$$

Step-8

$$\begin{aligned}
 q_2 &= \frac{\beta}{\|\beta\|} \\
 &= \frac{2}{\sqrt{6}} \begin{bmatrix} -1/2 \\ 1/2 \\ 1 \end{bmatrix} \\
 &= \begin{bmatrix} -1/\sqrt{6} \\ 1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix}
 \end{aligned}$$

Step-9

Therefore the required orthonormal basis is

$$\left\{ \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}, \begin{bmatrix} -1/\sqrt{6} \\ 1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix} \right\}$$

Step-10

P = the projection matrix on the plane = $A(A^T A)^{-1} A^T$

$$A = \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Here

$$\begin{aligned}
 A^T A &= \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}
 \end{aligned}$$

Step-11

$$\begin{aligned}
 (A^T A)^{-1} &= \frac{1}{3} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \\
 A(A^T A)^{-1} &= \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{1}{3} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \\
 &= \frac{1}{3} \begin{bmatrix} 1 & -1 \\ 2 & 1 \\ 1 & 2 \end{bmatrix}
 \end{aligned}$$

Step-12

$$\begin{aligned}
 P &= A(A^T A)^{-1} A^T \\
 &= \frac{1}{3} \begin{bmatrix} 1 & -1 \\ 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \\
 &= \frac{1}{3} \begin{bmatrix} 2 & 1 & -1 \\ 1 & 2 & 1 \\ -1 & 1 & 2 \end{bmatrix}
 \end{aligned}$$

This is the required projection matrix.

Step-13

The nullspace of P is obtained from $Px = 0$

$$\frac{1}{3} \begin{bmatrix} 2 & 1 & -1 \\ 1 & 2 & 1 \\ -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Step-14

Apply $R_2 \rightarrow 2R_2 - R_1, R_3 \rightarrow 2R_3 + R_1$

$$\begin{bmatrix} 2 & 1 & -1 \\ 0 & 3 & 3 \\ 0 & 3 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$2x + y - z = 0$$

$$3x + 3y = 0$$

$$\Rightarrow x = -y$$

Step-15

if $y = k$
 $\Rightarrow x = -k$
 $z = 2x + y$
 $= -k$

Step-16

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -k \\ k \\ -k \end{bmatrix}$$

$$= -k \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

The nullspace of P is represented the vector $= \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

^ ^ ^ ^ ^ ^ ^ ^ ^ ^ ^ ^ ^ ^ ^