

1. (20 points, 4 points each) Label the following statements as **True** or **False**. Along with your answer, provide an informal proof, counterexample, or other explanation.

- (a) The sum of two positive operators on a finite-dimensional complex inner product space is positive. **True**
- (b) Let V be a 5-dimensional vector space and $T \in \mathcal{L}(V)$. Then there exists a 3-dimensional subspace U of V invariant under T . **False**
- (c) Any polynomial of degree n with leading coefficients $(-1)^n$ is the characteristic polynomial of some linear operator. **True**
- (d) If x, y , and z are vectors in an inner product space such that $\langle x, y \rangle = \langle x, z \rangle$, then $y = z$. **False**
- (e) Every normal operator is diagonalizable. **False**

2. (20 points) Suppose $T \in \mathcal{L}(\mathbb{R}^3)$ is defined by

$$T(x_1, x_2, x_3) = (2x_1, x_2 - x_3, x_2 + x_3) = (x_1, x_2, x_3) \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & 1 \end{pmatrix}$$

- (a) Determine the eigenspace of T corresponding to each eigenvalue. $\lambda = 2$
- (b) Find the Jordan form and a Jordan basis of T .
- (c) Find the minimal polynomial of T .
- (d) Find the trace of T , $\text{trace } T$.
- (e) Find the determinant of T , $\det T$.

第 1 页/共 2 页

考试科目: 线性代数精讲

2021-2022 学年秋季学期期末考试

3. (10 points) Suppose V is a finite-dimensional inner product space, $T \in \mathcal{L}(V)$ is normal, and U is a subspace of V that is invariant under T . Show that U^\perp is invariant under T . **9.30**
4. (20 points) Let T be a linear operator on a finite-dimensional vector space V , and let v be a nonzero vector in V . The subspace

$$U = \text{span} \{v, Tv, T^2v, \dots\}$$

is called the T -cyclic subspace of V generated by v .

- (a) Show that U is a finite-dimensional invariant subspace of V .
- (b) Let $k = \dim U$. Show that $\{v, Tv, T^2v, \dots, T^{k-1}v\}$ is a basis for U .
- (c) If $a_0v + a_1Tv + a_2T^2v + \dots + a_{k-1}T^{k-1}v + T^k(v) = 0$, show that the characteristic polynomial of $T|_U$ is

$$f(t) = (-1)^k(a_0 + a_1t + \dots + a_{k-1}t^{k-1} + t^k).$$

- (d) Let $g(t)$ be the characteristic polynomial of T , show that $g(T) = 0$, where 0 is the zero operator. That is, T "satisfies" its characteristic equation.

5. (10 points) If $\mathbb{F} = \mathbb{C}$, show that T is an isometry if and only if T is normal and $|\lambda| = 1$ for every eigenvalue λ of T .
6. (20 points) Let $\mathcal{P}_2(\mathbb{R})$ and $\mathcal{P}_1(\mathbb{R})$ be the polynomial spaces with inner products defined by

$$\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx, \quad f, g \in \mathcal{P}_2(\mathbb{R}).$$

Let $T: \mathcal{P}_2(\mathbb{R}) \rightarrow \mathcal{P}_1(\mathbb{R})$ be the linear operator defined by

$$T(f(x)) = f'(x).$$

- (a) Find orthonormal bases $\{v_1, v_2, v_3\}$ for $\mathcal{P}_2(\mathbb{R})$ and $\{u_1, u_2\}$ for $\mathcal{P}_1(\mathbb{R})$.
- (b) Find $p \in \mathcal{P}_1(\mathbb{R})$ that makes

$$\int_{-1}^1 |x^5 - p(x)|^2 dx$$

as small as possible.

- (c) Find the singular values σ_1, σ_2 of T such that $T(v_i) = \sigma_i u_i$, $i = 1, 2$, and $T(v_3) = 0$.

7. (Bonus Question) Let V be a real inner product space. A function $f: V \rightarrow V$ is called a **rigid motion** if

$$\|f(x) - f(y)\| = \|x - y\|$$

for all $x, y \in V$. For example, any **isometry** on a finite-dimensional real inner product space is a **rigid motion**. Another class of rigid motions are the translations. A function $g: V \rightarrow V$, where V is a real inner product space, is called a **translation** if there exists a vector $v_0 \in V$ such that $g(v) = v + v_0$ for all $v \in V$. Let $f: V \rightarrow V$ be a rigid motion on a finite-dimensional real inner product space V , show that there exists a unique isometry T on V and a unique translation g on V such that $f = g \circ T$.