

## Step-1

By choosing the correct vector in  $b$  in the Schwarz inequality, we have to prove that

$(a_1 + a_2 + \dots + a_n)^2 \leq n(a_1^2 + a_2^2 + \dots + a_n^2)$ , and also we have to find that, when does the equality holds.

## Step-2

Let 
$$a = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

Choose  $b = (1, 1, \dots, 1)$

Schwarz inequality is  $|a^T b| \leq \|a\| \|b\|$  (1)

## Step-3

$$a^T b = (a_1, a_2, \dots, a_n) \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix}$$

$$= a_1 + a_2 + \dots + a_n$$

$$\Rightarrow |a^T b| = |a_1 + a_2 + \dots + a_n|$$

$$\Rightarrow |a^T b|^2 = (a_1 + a_2 + \dots + a_n)^2$$

## Step-4

And  $\|a\| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$

$$\|b\| = \sqrt{1^2 + 1^2 + \dots + 1^2} \text{ (n times)}$$

$$= \sqrt{n} \cdot 1$$

## Step-5

$$\Rightarrow \|a\|^2 = a_1^2 + a_2^2 + \dots + a_n^2 \text{ and } \|b\|^2 = n$$

By (1),  $\left|a^T b\right|^2 \leq\left\|a\right\|^2\left\|b\right\|^2$

Hence  $\boxed{\left(a_1+a_2+\cdots+a_n\right)^2 \leq n\left(a_1^2+a_2^2+\cdots+a_n^2\right)}$

## Step-6

If  $a_1=a_2=\cdots=a_n$  then

$$\begin{aligned} & \left(a_1+a_2+\cdots+a_n\right)^2 \\ &= \left(n a_1\right)^2 \\ &= n^2 a_1^2 \end{aligned}$$

## Step-7

And

$$\begin{aligned} & n\left(a_1^2+a_2^2+\cdots+a_n^2\right) \\ &= n\left(n a_1^2\right) \\ &= n^2 a_1^2 \end{aligned}$$

Therefore  $\left(a_1+a_2+\cdots+a_n\right)^2=n\left(a_1^2+a_2^2+\cdots+a_n^2\right)$

Hence the equality holds if  $a_1=a_2=\cdots=a_n$