

Step-1

Consider the following skew-Hermitian matrix:

$$K = \begin{bmatrix} i & i \\ i & i \end{bmatrix}$$

Diagonalize matrix K and compute $e^{Kt} = Se^{\Lambda t}S^{-1}$. Verify that e^{Kt} is unitary. Also find the derivative of e^{Kt} at $t = 0$.

Step-2

First step is to find the Eigen values of matrix K . To calculate the Eigen values do the following calculations:

$$K - \lambda I = \begin{bmatrix} i - \lambda & i \\ i & i - \lambda \end{bmatrix}$$

$$\det(K - \lambda I) = 0$$

$$(i - \lambda)^2 + 1 = 0$$

$$(\lambda^2 - 2\lambda i) = 0$$

After solving following values are obtained:

$$\lambda_1 = 2i$$

$$\lambda_2 = 0$$

Step-3

To calculate Eigen vectors do the following calculations:

$$(K - \lambda I)x = 0$$

$$\begin{bmatrix} i - \lambda & i \\ i & i - \lambda \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -i & i \\ i & -i \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

On solving, values of y and z corresponding to $\lambda = 2i$ are as follows:

$$\begin{aligned} x_1 &= \begin{bmatrix} y \\ z \end{bmatrix} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} \end{aligned}$$

Step-4

Similarly, Eigen vector corresponding to Eigen value $\lambda = 0$ is as follows:

$$(K - \lambda I)x = 0$$

$$\begin{bmatrix} i - \lambda & i \\ i & i - \lambda \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} i & i \\ i & i \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

On solving values of y and z are as follows:

$$x_2 = \begin{bmatrix} y \\ z \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Step-5

Therefore, diagonalisation of a matrix can be done as below:

$$K = SAS^{-1}$$

$$= \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 2i & 0 \\ 0 & 0 \end{bmatrix} \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

Step-6

Now, compute the following:

$$e^{Kt} = Se^{At}S^{-1}$$

$$= \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} e^{2it} & 0 \\ 0 & e^{0t} \end{bmatrix} \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} e^{2it} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} e^{2it} + 1 & e^{2it} - 1 \\ e^{2it} - 1 & e^{2it} + 1 \end{bmatrix}$$

Step-7

Therefore,

$$e^{Kt} = \frac{1}{2} \begin{bmatrix} e^{2it} + 1 & e^{2it} - 1 \\ e^{2it} - 1 & e^{2it} + 1 \end{bmatrix}$$

Step-8

Recall that a matrix U is unitary if $U^H U = I$. To verify that e^{Kt} is unitary, do the following calculations:

$$\begin{aligned} (e^{Kt})^H e^{Kt} &= \frac{1}{2} \begin{bmatrix} e^{-2it} + 1 & e^{-2it} - 1 \\ e^{-2it} - 1 & e^{-2it} + 1 \end{bmatrix} \frac{1}{2} \begin{bmatrix} e^{2it} + 1 & e^{2it} - 1 \\ e^{2it} - 1 & e^{2it} + 1 \end{bmatrix} \\ &= \frac{1}{4} \begin{bmatrix} e^{-2it} + 1 & e^{-2it} - 1 \\ e^{-2it} - 1 & e^{-2it} + 1 \end{bmatrix} \begin{bmatrix} e^{2it} + 1 & e^{2it} - 1 \\ e^{2it} - 1 & e^{2it} + 1 \end{bmatrix} \\ &= \frac{1}{4} \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \\ &= I \end{aligned}$$

Therefore, e^{Kt} is unitary.

Step-9

Derivative of e^{Kt} at $t = 0$ will be:

$$\begin{aligned} e^{Kt} &= \frac{1}{2} \begin{bmatrix} e^{2it} + 1 & e^{2it} - 1 \\ e^{2it} - 1 & e^{2it} + 1 \end{bmatrix} \\ \frac{de^{Kt}}{dt} &= \frac{1}{2} \begin{bmatrix} 2ie^{2it} & 2ie^{2it} \\ 2ie^{2it} & 2ie^{2it} \end{bmatrix} \\ \left(\frac{de^{Kt}}{dt} \right)_{t=0} &= \begin{bmatrix} i & i \\ i & i \end{bmatrix} \\ &= K \end{aligned}$$

Therefore, derivative of e^{Kt} at $t = 0$ is K matrix.