

Step-1

Consider the matrices,

$$A = \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 2 & 0 & 6 \end{bmatrix}, \text{ and } b = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

The objective is to determine the echelon form U , the free variables, and special solutions.

The system $Ax = b$ is consistent when b satisfies $b_2 = \text{-----}$, and finds the complete solution.

Step-2

Consider the system,

$$Ax = b$$

$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 2 & 0 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

Apply row operation, $R_2 \rightarrow R_2 - 2R_1$.

$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 - 2b_1 \end{bmatrix} \quad \text{.....(1)}$$

Hence, the echelon form of matrix A is,

$$U = \begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Step-3

Consider the system,

$$Ax = 0$$
$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 2 & 0 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Apply row operation,

Apply row operation, $R_2 \rightarrow R_2 - 2R_1$.

$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The equations form is,

$$x_2 + 3x_4 = 0$$

Put, $x_4 = k_1, x_3 = k_2, x_1 = k_3$.

$$x_2 = -3k_1$$

Hence, the free variables are, x_1, x_3 , and x_4 .

Step-4

So, the solution is,

$$\begin{aligned} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} &= \begin{bmatrix} k_3 \\ -3k_1 \\ k_2 \\ k_1 \end{bmatrix} \\ &= \begin{bmatrix} 0+0+k_3+0 \\ -3k_1+0+0+0 \\ 0+k_2+0+0 \\ k_1+0+0+0 \end{bmatrix} \\ &= k_1 \begin{bmatrix} 0 \\ -3 \\ 0 \\ 1 \end{bmatrix} + k_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + k_3 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \end{aligned}$$

Therefore, the special solutions are, $(0, -3, 0, 1), (0, 0, 1, 0), (1, 0, 0, 0)$.

Step-5

The system is consistent if,

$$b_2 - 2b_1 = 0.$$

From system (1).

Therefore,

$$b_2 = 2b_1$$

Step-6

Consider the non-homogeneous system,

$$Ax = \begin{bmatrix} b_1 \\ 2b_1 \end{bmatrix} \text{ Since, } b_2 = 2b_1.$$

$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 2 & 0 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ 2b_1 \end{bmatrix}$$

Step-7

Apply row operation, $R_2 \rightarrow R_2 - 2R_1$.

$$\begin{bmatrix} 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ 0 \end{bmatrix}$$

$$x_2 + 3x_4 = b_1$$

$$x_2 = b_1 - 3x_4$$

Put, $x_4 = k_1, x_3 = k_2, x_1 = k_3$.

$$x_2 = 1 - 3k_1$$

Therefore, the solution is,

$$\begin{aligned}
\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} &= \begin{bmatrix} k_3 \\ b_1 - 3k_1 \\ k_2 \\ k_1 \end{bmatrix} \\
&= \begin{bmatrix} 0 \\ b_1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} k_3 \\ -3k_1 \\ k_2 \\ k_1 \end{bmatrix} \\
&= \begin{bmatrix} 0 \\ b_1 \\ 0 \\ 0 \end{bmatrix} + k_1 \begin{bmatrix} 0 \\ -3 \\ 0 \\ 1 \end{bmatrix} + k_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + k_3 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}
\end{aligned}$$

Step-8

Therefore, the complete solution is,

$$x_{\text{complete}} = x_{\text{particular}} + x_{\text{nullspace}}$$

$$x = x_p + x_n$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \left[\begin{bmatrix} 0 \\ b_1 \\ 0 \\ 0 \end{bmatrix} + k_1 \begin{bmatrix} 0 \\ -3 \\ 0 \\ 1 \end{bmatrix} + k_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + k_3 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right]$$