

# Regression with Randomized Experiments

Ryan T. Moore

2024-08-21

The Linear Regression Estimator

Models with Covariates

Heterogeneous Treatment Effects

Inference for Experiments using Regression Adjustment

“Controlling for Blocks”

Nonlinear Terms

How Causal Inference is Different

# The Linear Regression Estimator

## Estimating the ATE

In a policy experiment in India, Chattopadhyay and Duflo (2004) randomly assign reserved village council headship for women to villages.

## Estimating the ATE

In a policy experiment in India, Chattopadhyay and Duflo (2004) randomly assign reserved village council headship for women to villages.

```
library(readr)
seats <- read_csv("http://j.mp/2YfZdgv")
```

## Estimating the ATE

In a policy experiment in India, Chattopadhyay and Duflo (2004) randomly assign reserved village council headship for women to villages.

```
library(readr)
seats <- read_csv("http://j.mp/2YfZdgv")
```

```
head(seats)
```

```
## # A tibble: 6 x 6
##       GP village reserved female irrigation water
##   <dbl>   <dbl>   <dbl>  <dbl>      <dbl> <dbl>
## 1     1     2     1     1         0    10
## 2     1     1     1     1         5     0
## 3     2     2     1     1         2     2
## 4     2     1     1     1         4    31
## 5     3     2     0     0         0     0
## 6     3     1     0     0         0     0
```

## Estimating the ATE

Explore assignment:

```
table(seats$reserved)
```

```
##
```

```
##    0    1
```

```
## 214 108
```

## Estimating the ATE

Estimate treatment effect with difference in means:

```
mean(seats$water[seats$reserved == 1]) -  
  mean(seats$water[seats$reserved == 0])
```

```
## [1] 9.252423
```



## Estimating the ATE

Estimate treatment effect with difference in means:

```
mean(seats$water[seats$reserved == 1]) -  
  mean(seats$water[seats$reserved == 0])
```

```
## [1] 9.252423
```

In areas with reservations, about 9.3 more drinking water projects were undertaken.

With binary treatment as predictor, we can recover same result using least squares regression.

With binary treatment as predictor, we can recover same result using least squares regression.

Model:

$$\text{Water Projects}_i = \beta_0 + \beta_1 (\text{Women's Seats}_i) + \epsilon_i$$

With binary treatment as predictor, we can recover same result using least squares regression.

Model:

$$\text{Water Projects}_i = \beta_0 + \beta_1 (\text{Women's Seats}_i) + \epsilon_i$$

Estimate coefficients

```
lm_out <- lm(water ~ reserved, data = seats)
lm_out
```

```
##
## Call:
## lm(formula = water ~ reserved, data = seats)
##
## Coefficients:
## (Intercept)      reserved
##      14.738         9.252
```

Estimated model

$$\widehat{\text{Water Projects}}_i = 14.74 + 9.25 (\text{Women's Seats}_i)$$

## Estimated model

$$\widehat{\text{Water Projects}}_i = 14.74 + 9.25 (\text{Women's Seats}_i)$$

- ▶ Intercept: predicted value when Women's Seats are **not** reserved (**reserved** == 0)

## Estimated model

$$\widehat{\text{Water Projects}}_i = 14.74 + 9.25 (\text{Women's Seats}_i)$$

- ▶ Intercept: predicted value when Women's Seats are **not** reserved (**reserved** == 0)
- ▶ Slope: estimated average treatment effect (difference between **reserved** == 0 and **reserved** == 1).

## Estimated model

$$\widehat{\text{Water Projects}}_i = 14.74 + 9.25 (\text{Women's Seats}_i)$$

- ▶ Intercept: predicted value when Women's Seats are **not** reserved (**reserved** == 0)
- ▶ Slope: estimated average treatment effect (difference between **reserved** == 0 and **reserved** == 1).



## Estimated model

$$\widehat{\text{Water Projects}}_i = 14.74 + 9.25 (\text{Women's Seats}_i)$$

- ▶ Intercept: predicted value when Women's Seats are **not** reserved (**reserved** == 0)
- ▶ Slope: estimated average treatment effect (difference between **reserved** == 0 and **reserved** == 1).

*Causal* interpretation of coefficient comes from the science of the policy experiment, **not** from regression.

## Social Pressure Mail Experiment

Gerber, Green, and Larimer (2008): Social pressure mailer experiment; households randomly receive one of four messages

# Social Pressure Mail Experiment

Gerber, Green, and Larimer (2008): Social pressure mailer experiment; households randomly receive one of four messages

Dear Registered Voter:

DO YOUR CIVIC DUTY AND VOTE!

Why do so many people fail to vote? We've been talking about this problem for years, but it only seems to get worse.

The whole point of democracy is that citizens are active participants in government; that we have a voice in government. Your voice starts with your vote. On August 8, remember your rights and responsibilities as a citizen. Remember to vote.

DO YOUR CIVIC DUTY — VOTE!

# Social Pressure Mail Experiment

Gerber, Green, and Larimer (2008): Social pressure mailer experiment; households randomly receive one of four messages

Dear Registered Voter:

YOU ARE BEING STUDIED!

Why do so many people fail to vote? We've been talking about this problem for years, but it only seems to get worse.

This year, we're trying to figure out why people do or do not vote. We'll be studying voter turnout in the August 8 primary election.

Our analysis will be based on public records, so you will not be contacted again or disturbed in any way. Anything we learn about your voting or not voting will remain confidential and will not be disclosed to anyone else.

DO YOUR CIVIC DUTY — VOTE!

# Social Pressure Mail Experiment

Gerber, Green, and Larimer (2008): Social pressure mailer experiment; households randomly receive one of four messages

Dear Registered Voter:

## WHAT IF YOUR NEIGHBORS KNEW WHETHER YOU VOTED?

Why do so many people fail to vote? We've been talking about the problem for years, but it only seems to get worse. This year, we're taking a new approach. We're sending this mailing to you and your neighbors to publicize who does and does not vote.

The chart shows the names of some of your neighbors, showing which have voted in the past. After the August 8 election, we intend to mail an updated chart. You and your neighbors will all know who voted and who did not.

## DO YOUR CIVIC DUTY — VOTE!

MAPLE DR	Aug 04	Nov 04	Aug 06
9995 JOSEPH JAMES SMITH	Voted	Voted	_____
9995 JENNIFER KAY SMITH		Voted	_____
9997 RICHARD B JACKSON		Voted	_____
9999 KATHY MARIE JACKSON		Voted	_____
9999 BRIAN JOSEPH JACKSON		Voted	_____
9991 JENNIFER KAY THOMPSON		Voted	_____
9991 BOB R THOMPSON		Voted	_____
9993 BILL S SMITH			_____
9989 WILLIAM LUKE CASPER		Voted	_____
9989 JENNIFER SUE CASPER		Voted	_____
9987 MARIA S JOHNSON	Voted	Voted	_____
9987 TOM JACK JOHNSON	Voted	Voted	_____
9987 RICHARD TOM JOHNSON		Voted	_____

## Multiple Regression: Many Predictors

- ▶ Gerber, Green, and Larimer (2008): Social pressure mailing experiment; households randomly receive one of four messages

## Multiple Regression: Many Predictors

- ▶ Gerber, Green, and Larimer (2008): Social pressure mailing experiment; households randomly receive one of four messages
- ▶ *Four* predictors of turnout: assigned mailing with control, civic duty, Hawthorne, or social pressure message

## Multiple Regression: Many Predictors

- ▶ Gerber, Green, and Larimer (2008): Social pressure mailing experiment; households randomly receive one of four messages
- ▶ *Four* predictors of turnout: assigned mailing with control, civic duty, Hawthorne, or social pressure message
- ▶ Use all four to predict turnout in 2006 primary



## Multiple Regression: Many Predictors

- ▶ Gerber, Green, and Larimer (2008): Social pressure mailing experiment; households randomly receive one of four messages
- ▶ *Four* predictors of turnout: assigned mailing with control, civic duty, Hawthorne, or social pressure message
- ▶ Use all four to predict turnout in 2006 primary

## Multiple Regression: Many Predictors

- ▶ Gerber, Green, and Larimer (2008): Social pressure mailing experiment; households randomly receive one of four messages
- ▶ *Four* predictors of turnout: assigned mailing with control, civic duty, Hawthorne, or social pressure message
- ▶ Use all four to predict turnout in 2006 primary

Model:

$$\text{Turnout } 2006_i = \beta_0 + \beta_1 \cdot \text{Control}_i + \beta_2 \cdot \text{Hawthorne}_i + \beta_3 \cdot \text{Neighbors}_i + \epsilon_i$$

```
social <- read_csv("http://j.mp/2YenEuU")  
  
lm_out <- lm(primary2006 ~ messages, data = social)
```

```
social <- read_csv("http://j.mp/2YenEuU")  
  
lm_out <- lm(primary2006 ~ messages, data = social)
```

##	Estimate	Std. Error	t value	Pr(> t )
## (Intercept)	0.315	0.002	132.909	0.000
## messagesControl	-0.018	0.003	-6.905	0.000
## messagesHawthorne	0.008	0.003	2.341	0.019
## messagesNeighbors	0.063	0.003	18.944	0.000

Estimated model

$$\widehat{\text{Turnout } 2006}_i = 0.31 + -0.02 (\text{Control}_i) + 0.01 (\text{Hawthorne}_i) + 0.06 (\text{Neighbors}_i)$$

## Estimated model

$$\widehat{\text{Turnout } 2006}_i = 0.31 + -0.02 (\text{Control}_i) + 0.01 (\text{Hawthorne}_i) + 0.06 (\text{Neighbors}_i)$$

1. What is the predicted turnout probability under **Control**?
2. What is the predicted turnout probability under **Hawthorne**?
3. What is the predicted turnout probability under **Neighbors**?
4. What is the predicted turnout probability under **Civic Duty**?

Alternatively, estimate

$$\text{Turnout } 06_i = \beta_0 \cdot \text{CivicDuty}_i + \beta_1 \cdot \text{Contrl}_i + \beta_2 \cdot \text{Hawthorne}_i + \beta_3 \cdot \text{Neighbors}_i + \epsilon_i$$

```
lm_out <- lm(primary2006 ~ -1 + messages, data = social)
summary(lm_out)$coefficients
```

##		Estimate	Std. Error	t value	Pr(> t )
##	messagesCivic Duty	0.3145377	0.002366570	132.9087	
##	messagesControl	0.2966383	0.001057939	280.3927	
##	messagesHawthorne	0.3223746	0.002367004	136.1952	
##	messagesNeighbors	0.3779482	0.002367097	159.6674	

Estimated model:

$$\widehat{\text{Turnout } 2006}_i = 0.31 (\text{Civic Duty}_i) + 0.3 (\text{Control}_i) + \\ 0.32 (\text{Hawthorne}_i) + 0.38 (\text{Neighbors}_i)$$



Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.31 (\text{Civic Duty}_i) + 0.3 (\text{Control}_i) + 0.32 (\text{Hawthorne}_i) + 0.38 (\text{Neighbors}_i)$$

1. What is the predicted turnout probability under **Control**?
2. What is the predicted turnout probability under **Hawthorne**?
3. What is the predicted turnout probability under **Neighbors**?
4. What is the predicted turnout probability under **Civic Duty**?

## Models with Covariates

## Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$
- ▶ estimate effects for subgroups

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$
- ▶ estimate effects for subgroups
- ▶ account for design features

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$
- ▶ estimate effects for subgroups
- ▶ account for design features
- ▶ improve predictions of outcome (beyond causal inf)



# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$
- ▶ estimate effects for subgroups
- ▶ account for design features
- ▶ improve predictions of outcome (beyond causal inf)

# Why add covariates to model?

We consider adding covariates to our linear model to

- ▶ adjust for chance differences
  - ▶ reduce estimation error
- ▶ improve precision of  $\hat{\beta}_1$
- ▶ estimate effects for subgroups
- ▶ account for design features
- ▶ improve predictions of outcome (beyond causal inf)
- ▶ (not to reduce bias in estimator, if simple design)

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

- ▶ Freedman (2008): it can.



## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

- ▶ Freedman (2008): it can.
- ▶ Green and Aronow (2011): Actually, not in practice.

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

- ▶ Freedman (2008): it can.
- ▶ Green and Aronow (2011): Actually, not in practice.

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

- ▶ Freedman (2008): it can.
- ▶ Green and Aronow (2011): Actually, not in practice. And, small relative to  $SE(\widehat{TE})$ .

## Does adjusting for covariates risk bias?

$$Y_i = \beta_0 + \beta_1 T_i + \epsilon_i$$

- ▶ The bivariate regression  $\hat{\beta}_1$  is unbiased for the ATE
- ▶  $E(\hat{\beta}_1) = \beta_1$  (diff in means, with indep from randomization)

Does adding covariates introduce bias?

- ▶ Freedman (2008): it can.
- ▶ Green and Aronow (2011): Actually, not in practice. And, small relative to  $SE(\widehat{TE})$ .
- ▶ “Included variable bias” (see <https://t.ly/ttYpD>)

# Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates

# Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates
- ▶ Covariates are *pre-treatment*!

# Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates
- ▶ Covariates are *pre-treatment*!
- ▶ E.g., experiment not quite balanced on age, prior turnout; adjust diff-in-means.

## Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates
- ▶ Covariates are *pre-treatment!*
- ▶ E.g., experiment not quite balanced on age, prior turnout; adjust diff-in-means.
- ▶ E.g., make better turnout predictions by including registrant's age and prior turnout, in addition to whether received **Neighbors** postcard



## Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates
- ▶ Covariates are *pre-treatment*!
- ▶ E.g., experiment not quite balanced on age, prior turnout; adjust diff-in-means.
- ▶ E.g., make better turnout predictions by including registrant's age and prior turnout, in addition to whether received **Neighbors** postcard
- ▶ Age and prior turnout are *causally prior* to treatment

## Models with Covariates

- ▶ May want to adjust for or improve predictions based on several covariates
- ▶ Covariates are *pre-treatment*!
- ▶ E.g., experiment not quite balanced on age, prior turnout; adjust diff-in-means.
- ▶ E.g., make better turnout predictions by including registrant's age and prior turnout, in addition to whether received **Neighbors** postcard
- ▶ Age and prior turnout are *causally prior* to treatment
- ▶ Let's just compare **Neighbors** to **Control**.

## Models with Covariates

```
# Subset to two treatment conditions:  
social.neighbor <- subset(social,  
                           (messages == "Control") |  
                           (messages == "Neighbors"))  
  
# Calculate age:  
social.neighbor <- social.neighbor |>  
  mutate(age = 2006 - yearofbirth)
```

```
# Multiple regression:
```

```
lm_out <- lm(primary2006 ~ age + primary2004 +  
              messages, data = social.neighbor)
```

```
summary(lm_out)$coefficients |> round(3)
```

##	Estimate	Std. Error	t value	Pr(> t )
## (Intercept)	0.046	0.003	13.274	
## age	0.004	0.000	58.873	
## primary2004	0.147	0.002	76.464	
## messagesNeighbors	0.080	0.003	31.670	

Now, make different prediction depending on registrant's age and prior turnout, not just treatment status.

Now, make different prediction depending on registrant's age and prior turnout, not just treatment status.

Estimated model:

$$\widehat{\text{Turnout } 2006}_i = 0.05 + 0.004 (\text{Age}_i) + 0.15 (\text{Turnout } 2004_i) + 0.08 (\text{Neighbors}_i)$$

Now, make different prediction depending on registrant's age and prior turnout, not just treatment status.

Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.05 + 0.004 (\text{Age}_i) + 0.15 (\text{Turnout } 2004_i) + 0.08 (\text{Neighbors}_i)$$

Coefficients: how much 1-unit difference in predictor affects my prediction for turnout in 2006, *assuming the other predictors do not change*. (The *ceteris paribus* assumption.) Sometimes a valid assumption, sometimes not.

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?



If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout } 2006_1} = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout } 2004_1) + 0.08 (\text{Neighbors}_1)$$

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout } 2006_1} = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout } 2004_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout } 2006_2} = 0.05 + 0.004 (\text{Age}_1 + 10) + 0.15 (\text{Turnout } 2004_1) + 0.08 (\text{Neighbors}_1)$$

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout 2006}}_1 = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout 2006}}_2 = 0.05 + 0.004 (\text{Age}_1 + 10) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout 2006}}_2 - \widehat{\text{Turnout 2006}}_1 = 0.004 (\text{Age}_1 + 10) - 0.004 (\text{Age}_1)$$

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout 2006}}_1 = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout 2006}}_2 = 0.05 + 0.004 (\text{Age}_1 + 10) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\begin{aligned}\widehat{\text{Turnout 2006}}_2 - \widehat{\text{Turnout 2006}}_1 &= 0.004 (\text{Age}_1 + 10) - 0.004 (\text{Age}_1) \\ &= 0.004 \times 10\end{aligned}$$

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout 2006}}_1 = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout 2006}}_2 = 0.05 + 0.004 (\text{Age}_1 + 10) + 0.15 (\text{Turnout 2004}_1) + 0.08 (\text{Neighbors}_1)$$

$$\begin{aligned}\widehat{\text{Turnout 2006}}_2 - \widehat{\text{Turnout 2006}}_1 &= 0.004 (\text{Age}_1 + 10) - 0.004 (\text{Age}_1) \\ &= 0.004 \times 10 \\ &\approx 0.038\end{aligned}$$

If compare two registrants, *ceteris paribus*, ages differ by 10 years, how much would turnout probability prediction differ?

$$\widehat{\text{Turnout } 2006_1} = 0.05 + 0.004 (\text{Age}_1) + 0.15 (\text{Turnout } 2004_1) + 0.08 (\text{Neighbors}_1)$$

$$\widehat{\text{Turnout } 2006_2} = 0.05 + 0.004 (\text{Age}_1 + 10) + 0.15 (\text{Turnout } 2004_1) + 0.08 (\text{Neighbors}_1)$$

$$\begin{aligned}\widehat{\text{Turnout } 2006_2} - \widehat{\text{Turnout } 2006_1} &= 0.004 (\text{Age}_1 + 10) - 0.004 (\text{Age}_1) \\ &= 0.004 \times 10 \\ &\approx 0.038\end{aligned}$$

Not causal.

## Heterogeneous Treatment Effects

## Heterogeneous Treatment Effects

Often, believe  $\tau_i$  vary systematically by some  $X$ .



## Heterogeneous Treatment Effects

Often, believe  $\tau_i$  vary systematically by some  $X$ .

- ▶ Reserved seats for women may only produce more water projects in *larger* villages, e.g.

## Heterogeneous Treatment Effects

Often, believe  $\tau_i$  vary systematically by some  $X$ .

- ▶ Reserved seats for women may only produce more water projects in *larger* villages, e.g.
- ▶ Reserved seats may produce *fewer* water projects in small villages

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0))$

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0))$

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

RI test for heterogeneity:

- ▶ Calculate observed diff-in-variances

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

RI test for heterogeneity:

- ▶ Calculate observed diff-in-variances
- ▶ Assume  $\tau_i = \hat{\tau}$



## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

RI test for heterogeneity:

- ▶ Calculate observed diff-in-variances
- ▶ Assume  $\tau_i = \hat{\tau}$
- ▶ Impute missing potential outcomes

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

RI test for heterogeneity:

- ▶ Calculate observed diff-in-variances
- ▶ Assume  $\tau_i = \hat{\tau}$
- ▶ Impute missing potential outcomes
- ▶ Create reference dist'n of 1000  
diff-in-variances

## A Test for Heterogeneity

- ▶ “Heterogeneity” is  $\text{Var}(\tau_i) \neq 0$
- ▶ If no heterogeneity, then  
 $\text{Var}(Y_i(1)) = \text{Var}(Y_i(0)) \rightsquigarrow$  null hypothesis

RI test for heterogeneity:

- ▶ Calculate observed diff-in-variances
- ▶ Assume  $\tau_i = \hat{\tau}$
- ▶ Impute missing potential outcomes
- ▶ Create reference dist'n of 1000  
diff-in-variances
- ▶  $p$ -value = prop of ref dist'n as extreme or  
more extreme than observed diff-in-variances.

## Testing for Heterogeneity

```
obs_diff_in_vars <- var(seats$water[seats$reserved == 1]) -  
  var(seats$water[seats$reserved == 0])
```

## Testing for Heterogeneity

```
obs_diff_in_vars <- var(seats$water[seats$reserved == 1]) -  
  var(seats$water[seats$reserved == 0])
```

```
obs_diff_in_vars
```

```
## [1] 2266.654
```

## Testing for Heterogeneity

```
n_perms <- 2000 # how many permutations  
store_dvs <- rep(NA, n_perms) # storage  
base_assg <- c(rep(0, 214), rep(1, 108))
```

## Testing for Heterogeneity

```
n_perms <- 2000 # how many permutations
store_dvs <- rep(NA, n_perms) # storage
base_assg <- c(rep(0, 214), rep(1, 108))
```

```
for(i in 1:n_perms){
  perm_tr <- sample(base_assg)
  diff_in_var <- var(seats$water[perm_tr == 1]) -
    var(seats$water[perm_tr == 0])
  store_dvs[i] <- diff_in_var
}
```

## Testing for Heterogeneity

```
n_perms <- 2000 # how many permutations
store_dvs <- rep(NA, n_perms) # storage
base_assg <- c(rep(0, 214), rep(1, 108))
```

```
for(i in 1:n_perms){
  perm_tr <- sample(base_assg)
  diff_in_var <- var(seats$water[perm_tr == 1]) -
    var(seats$water[perm_tr == 0])
  store_dvs[i] <- diff_in_var
}
```

```
p_val <- mean(abs(store_dvs) >= abs(obs_diff_in_vars))
```

```
p_val
```

```
## [1] 0.005
```



## Testing for Heterogeneity

```
n_perms <- 2000 # how many permutations
store_dvs <- rep(NA, n_perms) # storage
base_assg <- c(rep(0, 214), rep(1, 108))
```

```
for(i in 1:n_perms){
  perm_tr <- sample(base_assg)
  diff_in_var <- var(seats$water[perm_tr == 1]) -
    var(seats$water[perm_tr == 0])
  store_dvs[i] <- diff_in_var
}
```

```
p_val <- mean(abs(store_dvs) >= abs(obs_diff_in_vars))
```

```
p_val
```

```
## [1] 0.005
```

Since  $p < \alpha = 0.05$ , reject null of “no heterogeneity”.

# Heterogeneous Effects

If evidence or theory for heterogeneous effects, estimate them!

## Heterogeneous Effects

If evidence or theory for heterogeneous effects, estimate them!

Estimand: Conditional Average Treatment Effect (CATE)

## Heterogeneous Effects

If evidence or theory for heterogeneous effects, estimate them!

Estimand: Conditional Average Treatment Effect (CATE)

$$\begin{aligned} CATE &= E(Y_1 - Y_0 | X = x) \\ &= E(Y_1 | X = x) - E(Y_0 | X = x) \end{aligned}$$

# Estimating Heterogeneous Treatment Effects

- ▶ Subset on covariate  $X$ , estimate ATE within subsets

# Estimating Heterogeneous Treatment Effects

- ▶ Subset on covariate  $X$ , estimate ATE within subsets
- ▶ Estimate with interaction term in single model

## Interaction Terms for Heterogeneous Effects

- ▶ Add predictor variable representing subgroups in multiple regression equation

## Interaction Terms for Heterogeneous Effects

- ▶ Add predictor variable representing subgroups in multiple regression equation
- ▶ “Interact it with” (i.e., multiply it by) treatment to get us treatment effect in subgroups



## Interaction Terms for Heterogeneous Effects

- ▶ Add predictor variable representing subgroups in multiple regression equation
- ▶ “Interact it with” (i.e., multiply it by) treatment to get us treatment effect in subgroups

## Interaction Terms for Heterogeneous Effects

- ▶ Add predictor variable representing subgroups in multiple regression equation
- ▶ “Interact it with” (i.e., multiply it by) treatment to get us treatment effect in subgroups

```
lm_out <- lm(primary2006 ~ primary2004 +  
              messages + primary2004:messages,  
              data = social.neighbor)  
summary(lm_out)$coef |> round(3)
```

##	Estimate	Std. Error	t value
## (Intercept)	0.237	0.001	176.32
## primary2004	0.149	0.002	69.96
## messagesNeighbors	0.069	0.003	20.93
## primary2004:messagesNeighbors	0.027	0.005	5.23

Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.24 + 0.15 (\text{Turnout } 2004_i) + 0.07 (\text{Neighbors}_i) + 0.03 (\text{Turnout } 2004_i) (\text{Neighbors}_i)$$

Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.24 + 0.15 (\text{Turnout } 2004_i) + 0.07 (\text{Neighbors}_i) + 0.03 (\text{Turnout } 2004_i) (\text{Neighbors}_i)$$

Calculate

1. Predicted turnout prob under **Control**, if no 2004 vote?

Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.24 + 0.15 (\text{Turnout } 2004_i) + 0.07 (\text{Neighbors}_i) + 0.03 (\text{Turnout } 2004_i) (\text{Neighbors}_i)$$

Calculate

1. Predicted turnout prob under **Control**, if no 2004 vote?
2. Predicted turnout prob under **Neighbors**, if no 2004 vote?

Estimated model:

$$\widehat{\text{Turnout } 2006_i} = 0.24 + 0.15 (\text{Turnout } 2004_i) + 0.07 (\text{Neighbors}_i) + 0.03 (\text{Turnout } 2004_i) (\text{Neighbors}_i)$$

Calculate

1. Predicted turnout prob under **Control**, if no 2004 vote?
2. Predicted turnout prob under **Neighbors**, if no 2004 vote?
3. Treatment effect for those who didn't vote in 2004?

Estimated model:

$$\widehat{\text{Turnout } 2006}_i = 0.24 + 0.15 (\text{Turnout } 2004_i) + 0.07 (\text{Neighbors}_i) + 0.03 (\text{Turnout } 2004_i) (\text{Neighbors}_i)$$

Calculate

1. Predicted turnout prob under **Control**, if no 2004 vote?
2. Predicted turnout prob under **Neighbors**, if no 2004 vote?
3. Treatment effect for those who didn't vote in 2004?
4. Predicted turnout prob under **Control**, if voted in 2004?
5. Predicted turnout prob under **Neighbors**, if voted in 2004?
6. Treatment effect for those who voted in 2004?

# Interaction Terms for Heterogeneous Effects

$$\widehat{\text{Turnout } 2006}_i = \beta_0 + \beta_1 X_i + \beta_2 T_i + \beta_3 X_i T_i$$



## Interaction Terms for Heterogeneous Effects

$$\begin{aligned}\widehat{\text{Turnout } 2006}_i &= \beta_0 + \beta_1 X_i + \beta_2 T_i + \beta_3 X_i T_i \\ &= \underbrace{\beta_0}_{\bar{Y} \text{ if } (0,0)} + \underbrace{\beta_1}_{\bar{Y} \text{ if } X_i=1} X_i + \underbrace{\beta_2}_{\text{ATE if } X_i=0} T_i + \underbrace{\beta_3}_{\text{HTE}} X_i T_i\end{aligned}$$

# Interaction Terms for Heterogeneous Effects

$$\begin{aligned}
 \widehat{\text{Turnout } 2006_i} &= \beta_0 + \beta_1 X_i + \beta_2 T_i + \beta_3 X_i T_i \\
 &= \underbrace{\beta_0}_{\bar{Y} \text{ if } (0,0)} + \underbrace{\beta_1}_{\bar{Y} \text{ if } X_i=1} X_i + \underbrace{\beta_2}_{\text{ATE if } X_i=0} T_i + \underbrace{\beta_3}_{\text{HTE}} X_i T_i \\
 \widehat{\text{Turnout } 2006_i} &= \underbrace{0.24}_{\bar{Y} \text{ for } (0,0), \text{ Prior Nonvoters, Control}} + \\
 &\quad \underbrace{0.15}_{\text{Additional } \bar{Y} \text{ for Prior Voters}} (\text{Turnout } 2004_i) + \\
 &\quad \underbrace{0.07}_{\text{ATE for Prior Nonvoters; part of ATE for Prior Voters}} (\text{Neighb} \\
 &\quad \underbrace{0.03}_{\text{Additional ATE for Prior Voters}} (\text{Turnout } 2004_i) (\text{Neighb}
 \end{aligned}$$

Exercise:

Consider the data on UK MP candidates at <http://j.mp/32PHfFd>

1. Read the data into R and name it `mps`
2. Create a new variable called `winner`. Set it equal to 1 if `margin` is greater than 0; set it to 0 if `margin` is less than zero.
3. Consider `winner` to be a randomly assigned treatment. Estimate the causal effect of `winner` on `ln.net`, net worth at death of these candidates.
4. Investigate whether the effect of `winner` varies by `party`.

## Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?

# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?
- ▶  $\frac{3}{4}$  of the values below it?

# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?
- ▶  $\frac{3}{4}$  of the values below it?
- ▶  $\frac{1}{5}$  of the values below it?

# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?
- ▶  $\frac{3}{4}$  of the values below it?
- ▶  $\frac{1}{5}$  of the values below it?
- ▶ ...



# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?
- ▶  $\frac{3}{4}$  of the values below it?
- ▶  $\frac{1}{5}$  of the values below it?
- ▶ ...

# Quantile Average Treatment Effects

A *quantile* is a cut-point, or a position, in a statistical distribution.

What is the name for the value that has

- ▶ half the values below it?
- ▶  $\frac{3}{4}$  of the values below it?
- ▶  $\frac{1}{5}$  of the values below it?
- ▶ ...

These are *quantiles*.


## Quantile Average Treatment Effects (data)

```
set.seed(326370675)
n <- 1000
Y1 <- Y0 <- runif(n)
# If low baseline, negative TE:
Y1[Y0 < .5] <- Y0[Y0 < .5] - rnorm(length(Y0[Y0 < .5]))
# If high baseline, positive TE:
Y1[Y0 > .5] <- Y0[Y0 > .5] + rnorm(length(Y0[Y0 > .5]))
D <- sample((1:n) %% 2) # Assign 0/1
Y <- D * Y1 + (1 - D) * Y0 # Y_obs
samp <- tibble(D, Y)
library(quantreg)
ate <- coef(lm(Y ~ D, data = samp))[2]
qtes <- rq(Y ~ D,
            tau = seq(.05, .95, length.out = 10), # At what p
            data = samp, method = "fn")
```

## Quantile Average Treatment Effects (data)

```
set.seed(326370675)
n <- 1000
Y1 <- Y0 <- runif(n)
# If low baseline, negative TE:
Y1[Y0 < .5] <- Y0[Y0 < .5] - rnorm(length(Y0[Y0 < .5]))
# If high baseline, positive TE:
Y1[Y0 > .5] <- Y0[Y0 > .5] + rnorm(length(Y0[Y0 > .5]))
D <- sample((1:n) %% 2) # Assign 0/1
Y <- D * Y1 + (1 - D) * Y0 # Y_obs
samp <- tibble(D, Y)
library(quantreg)
ate <- coef(lm(Y ~ D, data = samp))[2]
qtes <- rq(Y ~ D,
            tau = seq(.05, .95, length.out = 10), # At what p
            data = samp, method = "fn")
```


(See <https://bit.ly/3dnNhGP>)



## Quantile Average Treatment Effects (new data)

```
set.seed(21578100)
n <- 1000
Y1 <- Y0 <- runif(n)
# If low baseline, zero TE:
Y1[Y0 < .5] <- Y0[Y0 < .5]
# If high baseline, positive TE:
Y1[Y0 > .5] <- Y0[Y0 > .5] + rnorm(length(Y0[Y0 > .5]))
D <- sample((1:n) %% 2) # Assign 0/1
Y <- D * Y1 + (1 - D) * Y0 # Y_obs
samp <- tibble(D, Y)

ate <- coef(lm(Y ~ D, data = samp))[2]
qtes <- rq(Y ~ D,
           tau = seq(.05, .95, length.out = 10),
           data = samp, method = "fn")
```



# Inference for Experiments using Regression Adjustment



# Inference for Experiments using Regression Adjustment

Lin (2013) compares several estimates of SE for linear models with experimental data, adjusting for covariates  $x$ .

Tailored for fixed-sample, variation-from-assignment situations.

# Inference for Experiments using Regression Adjustment

Lin (2013) compares several estimates of SE for linear models with experimental data, adjusting for covariates  $x$ .

Tailored for fixed-sample, variation-from-assignment situations.

“Sandwich” estimators of variance:

# Inference for Experiments using Regression Adjustment

Lin (2013) compares several estimates of SE for linear models with experimental data, adjusting for covariates  $x$ .

Tailored for fixed-sample, variation-from-assignment situations.

“Sandwich” estimators of variance:

$$\text{Var} \left( \widehat{\beta_{\text{OLS}}} \right) = (X'X)^{-1} (X' \text{diag}(\hat{\epsilon}_1, \dots, \hat{\epsilon}_n) X) (X'X)^{-1}$$

# Inference for Experiments using Regression Adjustment

Model:

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (x_i - \bar{x}) + \beta_3 T_i (x_i - \bar{x}) + \epsilon_i$$

# Inference for Experiments using Regression Adjustment

Model:

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (x_i - \bar{x}) + \beta_3 T_i (x_i - \bar{x}) + \epsilon_i$$

“Full set of treatment by (demeaned) covariate interactions”

# Inference for Experiments using Regression Adjustment

Model:

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (x_i - \bar{x}) + \beta_3 T_i (x_i - \bar{x}) + \epsilon_i$$

“Full set of treatment by (demeaned) covariate interactions”

Then, use “sandwich” HC2 standard errors.

# Inference for Experiments using Regression Adjustment

Model:

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (x_i - \bar{x}) + \beta_3 T_i (x_i - \bar{x}) + \epsilon_i$$

“Full set of treatment by (demeaned) covariate interactions”

Then, use “sandwich” HC2 standard errors.

Causal estimate:

$$\widehat{ATE}_{\text{interact}} = \hat{\beta}_1$$

# Inference for Experiments using Regression Adjustment

Result: SE's that are



# Inference for Experiments using Regression Adjustment

Result: SE's that are

- ▶ Consistent under heteroskedasticity

# Inference for Experiments using Regression Adjustment

Result: SE's that are

- ▶ Consistent under heteroskedasticity
- ▶ At least as efficient as usual  $\widehat{ATE}_{OLS}$

# Inference for Experiments using Regression Adjustment

Result: SE's that are

- ▶ Consistent under heteroskedasticity
- ▶ At least as efficient as usual  $\widehat{ATE}_{OLS}$
- ▶ More efficient if  $n_T \neq n_C$

# Inference for Experiments using Regression Adjustment

Result: SE's that are

- ▶ Consistent under heteroskedasticity
- ▶ At least as efficient as usual  $\widehat{ATE}_{OLS}$
- ▶ More efficient if  $n_T \neq n_C$
- ▶ Doesn't *hurt* precision  
( $\widehat{ATE}_{OLS}$  does if  $x$  varies more with  $\tau_i$  than  $Y_1, Y_0$ , e.g.)

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 G_i + \beta_3 T_i G_i + \epsilon_i \quad (1)$$

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 G_i + \beta_3 T_i G_i + \epsilon_i \quad (1)$$

What's  $\widehat{ATE}$ ?

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 G_i + \beta_3 T_i G_i + \epsilon_i \quad (1)$$

What's  $\widehat{ATE}$ ? Not  $\hat{\beta}_1$ .



# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 G_i + \beta_3 T_i G_i + \epsilon_i \quad (1)$$

What's  $\widehat{ATE}$ ? Not  $\hat{\beta}_1$ . Not  $\hat{\beta}_1 + \hat{\beta}_3$ .

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

Suppose categorical  $G$  with 2 levels. Let  $G_i$  be indicator for first level. Expect heterogeneous treatment effects.

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 G_i + \beta_3 T_i G_i + \epsilon_i \quad (1)$$

What's  $\widehat{ATE}$ ? Not  $\hat{\beta}_1$ . Not  $\hat{\beta}_1 + \hat{\beta}_3$ .

(Need to weight by how many of each  $G$  type we have.)

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$ ?

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$ ?

$$\hat{\beta}_1$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$ ?

$$\hat{\beta}_1$$

What's  $\widehat{ATE}$  for  $G_i = 1$  group?

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$ ?

$$\hat{\beta}_1$$

What's  $\widehat{ATE}$  for  $G_i = 1$  group?

$$\widehat{ATE}_{G1} = \hat{\beta}_1 + \hat{\beta}_3(1 - \bar{G})$$

# Inference for Experiments using Regression Adjustment

## Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$ ?

$$\hat{\beta}_1$$

What's  $\widehat{ATE}$  for  $G_i = 1$  group?

$$\begin{aligned}\widehat{ATE}_{G1} &= \hat{\beta}_1 + \hat{\beta}_3(1 - \bar{G}) \\ &= \hat{\beta}_1 + \hat{\beta}_3 \times (\text{prop. of } G = 0)\end{aligned}$$



# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$  for  $G_i = 0$  group?

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$  for  $G_i = 0$  group?

$$\widehat{ATE}_{G0} = \hat{\beta}_1 + \hat{\beta}_3(0 - \bar{G})$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$  for  $G_i = 0$  group?

$$\begin{aligned}\widehat{ATE}_{G0} &= \hat{\beta}_1 + \hat{\beta}_3(0 - \bar{G}) \\ &= \hat{\beta}_1 - \hat{\beta}_3 \bar{G}\end{aligned}$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$  for  $G_i = 0$  group?

$$\begin{aligned}\widehat{ATE}_{G0} &= \hat{\beta}_1 + \hat{\beta}_3(0 - \bar{G}) \\ &= \hat{\beta}_1 - \hat{\beta}_3 \bar{G} \\ &= \hat{\beta}_1 - \hat{\beta}_3 \times (\text{prop. of } G = 1)\end{aligned}$$

# Inference for Experiments using Regression Adjustment

Recovering the ATE

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (G_i - \bar{G}) + \beta_3 T_i (G_i - \bar{G}) + \epsilon_i$$

What's  $\widehat{ATE}$  for  $G_i = 0$  group?

$$\begin{aligned}\widehat{ATE}_{G0} &= \hat{\beta}_1 + \hat{\beta}_3(0 - \bar{G}) \\ &= \hat{\beta}_1 - \hat{\beta}_3 \bar{G} \\ &= \hat{\beta}_1 - \hat{\beta}_3 \times (\text{prop. of } G = 1)\end{aligned}$$

# Inference for Experiments using Regression Adjustment

So, to adjust for covariates in experimental data,

1. Estimate

$$Y_i = \beta_0 + \beta_1 T_i + \beta_2 (x_i - \bar{x}) + \beta_3 T_i (x_i - \bar{x}) + \epsilon_i$$

2. Use HC2 standard errors (or analogous CR2 for clustered designs)

```
library(estimatr)
lm_robust(y ~ t + I(x - mean(x)) + t * I(x - mean(x)),
          data = df)
```

# Inference for Experiments using Regression Adjustment

So, to adjust for covariates  $x$  in experimental data,

```
lm_lin(y ~ t, covariates = ~ x, data = df)
```



# Inference for Experiments using Regression Adjustment

With two levels of treatment,

$$Y_i = \beta_0 + \beta_1 T1_i + \beta_2 T2_i + \beta_3 (x_i - \bar{x}) + \beta_4 T1_i (x_i - \bar{x}) + \beta_5 T2_i (x_i - \bar{x}) + \epsilon_i$$

# Inference for Experiments using Regression Adjustment

With two levels of treatment,

$$Y_i = \beta_0 + \beta_1 T1_i + \beta_2 T2_i + \beta_3 (x_i - \bar{x}) + \beta_4 T1_i (x_i - \bar{x}) + \beta_5 T2_i (x_i - \bar{x}) + \epsilon_i$$

etc.

## “Controlling for Blocks”

## Beware LSDV: “controlling for blocks”

A first-thought for estimating ATE for blocked design ...

## Beware LSDV: “controlling for blocks”

A first-thought for estimating ATE for blocked design ...

Estimate linear model with indicators for each block:

$$Y_i = \beta_0 + \beta_1 T_i + \gamma_1 B_{i1} + \gamma_2 B_{i2} + \dots + \epsilon_i$$

## Beware LSDV: “controlling for blocks”

However, when

1. TE varies by block
2.  $P(T = 1)$  varies by block

## Beware LSDV: “controlling for blocks”

However, when

1. TE varies by block
2.  $P(T = 1)$  varies by block

Using “block fixed effects” or “controlling for block IDs” is

## Beware LSDV: “controlling for blocks”

However, when

1. TE varies by block
2.  $P(T = 1)$  varies by block

Using “block fixed effects” or “controlling for block IDs” is

- ▶ biased for the estimate (ATE)



## Beware LSDV: “controlling for blocks”

However, when

1. TE varies by block
2.  $P(T = 1)$  varies by block

Using “block fixed effects” or “controlling for block IDs” is

- ▶ biased for the estimate ( $ATE$ )
- ▶ biased for the standard error ( $SE(ATE)$ )

## Beware LSDV: “controlling for blocks”


However, when

1. TE varies by block
2.  $P(T = 1)$  varies by block

Using “block fixed effects” or “controlling for block IDs” is

- ▶ biased for the estimate ( $ATE$ )
- ▶ biased for the standard error ( $SE(ATE)$ )
- ▶ worse than most everything else!

# Beware LSDV: “controlling for blocks”



## Beware LSDV: “controlling for blocks”

The intuition: regression over-weights high-variance blocks

block	block_ATE	block_ATE_est	n_j	p_j	sample_weight	fe_weight
1	4	4.06	100	0.5	0.33	0.45
2	2	1.66	100	0.7	0.33	0.38
3	0	0.02	100	0.9	0.33	0.16

## Beware LSDV: “controlling for blocks”

- ▶ Should I adjust for continuous  $x$ , or block IDs, where blocks come from coarsened  $x$ ?

## Beware LSDV: “controlling for blocks”

- ▶ Should I adjust for continuous  $x$ , or block IDs, where blocks come from coarsened  $x$ ?
- ▶ If both TE and Tr Prob vary across blocks, use **block IDs** to remove bias  
(since those determine assignment)

## Beware LSDV: “controlling for blocks”

- ▶ Should I adjust for continuous  $x$ , or block IDs, where blocks come from coarsened  $x$ ?
- ▶ If both TE and Tr Prob vary across blocks, use **block IDs** to remove bias  
(since those determine assignment)
- ▶ In all 4 of the “probabilities vary/constant” by “treatment effects vary/constant” possibilities, using **both** the IDs and the variables themselves feels like overkill, but performs well.

## (Related note: regression weights the cases)

**FIGURE 1 Example of nominal and effective samples from Jensen (2003)**

---



- ▶ Aronow and Samii (2016) shows how multiple regression weights its cases
- ▶ Interp: for causal inf., need causal methods, not just regression + representative sample




## Nonlinear Terms

# Terms to Capture Nonlinear Relationships

Let's take a step back from causal inference, and just think about trying to model the relationship between turnout probability and age.

```
# Calculate turnout prop for every year of age:  
prop_turnout <- tapply(social.neighbor$primary2006,  
                        social.neighbor$age, mean)  
  
# Plot:  
plot(prop_turnout ~ names(prop_turnout), xlab = "Age",  
      ylab = "Prop Turnout in Primary 2006", pch = 16)
```



This looks like a nonlinear relationship. Let's include an age-squared term to try to model this nonlinearity:

```
lm_out <- lm(primary2006 ~ age + I(age^2),  
              data = social.neighbor)  
lm_out
```

```
##
```

```
## Call:
```

```
## lm(formula = primary2006 ~ age + I(age^2), data = social
```

```
##
```

```
## Coefficients:
```

```
## (Intercept)          age      I(age^2)
```

```
## -2.868e-02      9.818e-03    -5.572e-05
```

The estimated model is

$$\widehat{\text{Turnout } 2006}_i = -0.02868 + 0.00982 (\text{Age}_i) + -6 \times 10^{-5} (\text{Age}_i^2)$$

1. What is the predicted turnout probability for a 30-year-old?
2. What is the predicted turnout probability for a 60-year-old?
3. What is the predicted turnout probability for a 80-year-old?
4. What is the predicted turnout probability for a 90-year-old?

```
predict(lm_out, data.frame(age = 40))
```

```
##           1  
## 0.2748905
```

```
predict(lm_out, data.frame(age = 60))
```

```
##           1  
## 0.3598197
```

```
predict(lm_out, data.frame(age = 80))
```


```
##           1  
## 0.4001768
```

```
predict(lm_out, data.frame(age = 100))
```

```
##           1  
## 0.3959618
```

We can also look at all the predicted values:

```
plot(lm_out$fitted.values ~ social.neighbor$age,  
     xlab = "Age", ylab = "Predicted Turnout")
```



## How Causal Inference is Different



# Data Science Approaches

Three tasks of data science:

# Data Science Approaches

Three tasks of data science:

- ▶ Description

# Data Science Approaches

Three tasks of data science:

- ▶ Description
- ▶ Prediction

# Data Science Approaches

Three tasks of data science:

- ▶ Description
- ▶ Prediction
- ▶ Causal Inference

# Data Science Approaches

Three tasks of data science:

- ▶ Description
- ▶ Prediction
- ▶ Causal Inference

# Data Science Approaches

Three tasks of data science:

- ▶ Description
- ▶ Prediction
- ▶ Causal Inference

Models/algorithms central to all three.

# Data Science Approaches

Three tasks of data science:

- ▶ Description
- ▶ Prediction
- ▶ Causal Inference

Models/algorithms central to all three.

Hernán, Hsu, and Healy (2019)

# Data Science Approaches

## Description

- ▶ Identifying patterns, etc.



# Data Science Approaches

## Description

- ▶ Identifying patterns, etc.
- ▶ E.g., clustering to discover groups

# Data Science Approaches

## Prediction

- ▶ Components

# Data Science Approaches

## Prediction

- ▶ Components

- ▶ Inputs/outputs  
(predictors/outcomes, features/responses, ...)

# Data Science Approaches

## Prediction

### ► Components

- Inputs/outputs  
(predictors/outcomes, features/responses, ...)
- Mapping from inputs to outputs  
(linear model, decision tree, ...)

# Data Science Approaches

## Prediction

### ► Components

- Inputs/outputs  
(predictors/outcomes, features/responses, ...)
- Mapping from inputs to outputs  
(linear model, decision tree, ...)
- Metric for evaluating mapping

# Data Science Approaches

## Prediction

- ▶ Components
  - ▶ Inputs/outputs  
(predictors/outcomes, features/responses, ...)
  - ▶ Mapping from inputs to outputs  
(linear model, decision tree, ...)
  - ▶ Metric for evaluating mapping
- ▶ With these, model/machine learning algorithm does the work

# Data Science Approaches

## Prediction

- ▶ Components
  - ▶ Inputs/outputs  
(predictors/outcomes, features/responses, ...)
  - ▶ Mapping from inputs to outputs  
(linear model, decision tree, ...)
  - ▶ Metric for evaluating mapping
- ▶ With these, model/machine learning algorithm does the work
- ▶ E.g., regression, random forests, neural networks, ...

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective



# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure
  - ▶ Not all inputs treated same

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure
  - ▶ Not all inputs treated same
  - ▶  $T$  v.  $\mathbf{X}$  – very different!

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure
  - ▶ Not all inputs treated same
  - ▶  $T$  v.  $\mathbf{X}$  – very different!
  - ▶ (the more knowledge, the better!)

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure
  - ▶ Not all inputs treated same
  - ▶  $T$  v.  $\mathbf{X}$  – very different!
  - ▶ (the more knowledge, the better!)
  - ▶ (alternative: solve fundamental problem of causal inference!  
☺)

# Data Science Approaches

## Causal Inference

- ▶ Potential outcomes/counterfactual/interventionist perspective
- ▶ Requires *expertise* different to description/prediction
- ▶ Requires more than summary statistics, metrics, etc.
- ▶ Requires some knowledge of causal structure
  - ▶ Not all inputs treated same
  - ▶  $T$  v.  $\mathbf{X}$  – very different!
  - ▶ (the more knowledge, the better!)
  - ▶ (alternative: solve fundamental problem of causal inference!  
☺)
- ▶ E.g., experiments, observational causal designs, ...



# Causal Inference with Models

Consider two loaded datasets:

# Causal Inference with Models

Consider two loaded datasets:


```
str(df1)
```

```
## tibble [100 x 3] (S3: tbl_df/tbl/data.frame)
##  $ covariate: num [1:100] -0.622 1.137 -0.238 1.529 -0.1
##  $ exposure : num [1:100] 0.0332 0.3627 0.2422 1.4633 0
##  $ outcome  : num [1:100] -0.429 2.675 -0.647 2.238 1.04
```


```
str(df2)
```

```
## tibble [100 x 3] (S3: tbl_df/tbl/data.frame)
##  $ exposure : num [1:100] 0.4862 0.0653 -1.4021 -0.546 -
##  $ outcome  : num [1:100] 1.706 0.669 -1.597 -1.733 0.61
##  $ covariate: num [1:100] 2.24 0.924 -0.999 -2.343 0.207
```

# Causal Inference with Models



# Causal Inference with Models



# Causal Inference with Models

Model each

```
lm_df1 <- lm(outcome ~ exposure, data = df1)
lm_df2 <- lm(outcome ~ exposure, data = df2)
```

```
## # A tibble: 4 x 4
##   data term          estimate std.error
##   <chr> <chr>          <dbl>    <dbl>
## 1 df1   (Intercept) -0.00671  0.120
## 2 df1   exposure      0.996    0.0927
## 3 df2   (Intercept)  0.133    0.0890
## 4 df2   exposure      1.00    0.0841
```

# Causal Inference with Models

Model each

```
lm_df1 <- lm(outcome ~ exposure, data = df1)
lm_df2 <- lm(outcome ~ exposure, data = df2)
```

```
## # A tibble: 4 x 4
##   data term          estimate std.error
##   <chr> <chr>          <dbl>    <dbl>
## 1 df1   (Intercept) -0.00671  0.120
## 2 df1   exposure      0.996    0.0927
## 3 df2   (Intercept)  0.133    0.0890
## 4 df2   exposure      1.00    0.0841
```

- Both cases: effect of exposure  $\approx 1$ .

# Causal Inference with Models

Model each

```
lm_df1 <- lm(outcome ~ exposure, data = df1)
lm_df2 <- lm(outcome ~ exposure, data = df2)
```

```
## # A tibble: 4 x 4
##   data term          estimate std.error
##   <chr> <chr>          <dbl>    <dbl>
## 1 df1   (Intercept) -0.00671  0.120
## 2 df1   exposure      0.996    0.0927
## 3 df2   (Intercept)  0.133    0.0890
## 4 df2   exposure      1.00    0.0841
```

- ▶ Both cases: effect of exposure  $\approx 1$ .
- ▶ Is this good? Is it correct?

# Causal Inference with Models

Model each

```
lm_df1 <- lm(outcome ~ exposure, data = df1)
lm_df2 <- lm(outcome ~ exposure, data = df2)
```

```
## # A tibble: 4 x 4
##   data term          estimate std.error
##   <chr> <chr>          <dbl>    <dbl>
## 1 df1   (Intercept) -0.00671  0.120
## 2 df1   exposure      0.996    0.0927
## 3 df2   (Intercept)  0.133    0.0890
## 4 df2   exposure      1.00    0.0841
```

- ▶ Both cases: effect of exposure  $\approx 1$ .
- ▶ Is this good? Is it correct?
- ▶ What if we adjust for covariate?



## Causal Inference with Models

```
lm_df1_adj <- lm(outcome ~ exposure + covariate, data = df1)
lm_df2_adj <- lm(outcome ~ exposure + covariate, data = df2)
```

```
## # A tibble: 4 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
## 2 df1  covariate    0.970    0.147
## 3 df2  exposure    0.554    0.0990
## 4 df2  covariate    0.385    0.0598
```

- Both cases: effect of exposure  $\approx 0.5$ .

## Causal Inference with Models

```
lm_df1_adj <- lm(outcome ~ exposure + covariate, data = df1)
lm_df2_adj <- lm(outcome ~ exposure + covariate, data = df2)
```

```
## # A tibble: 4 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
## 2 df1  covariate    0.970    0.147
## 3 df2  exposure    0.554    0.0990
## 4 df2  covariate    0.385    0.0598
```

- ▶ Both cases: effect of exposure  $\approx 0.5$ .
- ▶ Is this good? Is it correct?

## Causal Inference with Models

```
lm_df1_adj <- lm(outcome ~ exposure + covariate, data = df1)
lm_df2_adj <- lm(outcome ~ exposure + covariate, data = df2)
```

```
## # A tibble: 4 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
## 2 df1  covariate    0.970    0.147
## 3 df2  exposure    0.554    0.0990
## 4 df2  covariate    0.385    0.0598
```

- ▶ Both cases: effect of exposure  $\approx 0.5$ .
- ▶ Is this good? Is it correct?
- ▶ Which is correct?  $\beta = 1$ ?  $\beta = 0.5$ ?

## Causal Inference with Models

```
lm_df1_adj <- lm(outcome ~ exposure + covariate, data = df1)
lm_df2_adj <- lm(outcome ~ exposure + covariate, data = df2)
```

```
## # A tibble: 4 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
## 2 df1  covariate    0.970    0.147
## 3 df2  exposure    0.554    0.0990
## 4 df2  covariate    0.385    0.0598
```

- ▶ Both cases: effect of exposure  $\approx 0.5$ .
- ▶ Is this good? Is it correct?
- ▶ Which is correct?  $\beta = 1$ ?  $\beta = 0.5$ ?
- ▶ *Should* we adjust for covariate?

## Causal Inference with Models

There is nothing in the data that tells us.


## Causal Inference with Models

There is nothing in the data that tells us. ☹

## Causal Inference with Models

There is nothing in the data that tells us. ☹


Here are the true structures: First



## Causal Inference with Models

There is nothing in the data that tells us. ☹

Here are the true structures: Second





## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**

# Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**

## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**

## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**

```
g_conf <- dagitty("dag{ x -> y ; x <- c -> y }")  
g_coll <- dagitty("dag{ x -> y ; x -> c <- y }")
```

## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**

```
g_conf <- dagitty("dag{ x -> y ; x <- c -> y }")  
g_coll <- dagitty("dag{ x -> y ; x -> c <- y }")
```

```
adjustmentSets(g_conf, "x", "y")
```

```
## { c }
```

## Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**

```
g_conf <- dagitty("dag{ x -> y ; x <- c -> y }")  
g_coll <- dagitty("dag{ x -> y ; x -> c <- y }")
```

```
adjustmentSets(g_conf, "x", "y")
```

```
## { c }
```


```
adjustmentSets(g_coll, "x", "y")
```

```
## {}
```

# Causal Inference with Models

When know structures, adjustment sets for unbiasedness differ:

- ▶ df1: confounding  $\Rightarrow$  **adjust for  $X$**
- ▶ df2: collider  $\Rightarrow$  **do not adjust for  $X$**





## Causal Inference with Models

So, correct adjustments to reveal causal effect of  $T \rightsquigarrow Y$ :

## Causal Inference with Models

So, correct adjustments to reveal causal effect of  $T \rightsquigarrow Y$ :

df1, **adjust** for  $X$ ,  $\beta = 0.5$ :

```
## # A tibble: 1 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
```

df2, **do not adjust** for  $X$ ,  $\beta = 1$ :

```
## # A tibble: 1 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df2  exposure    1.00    0.0841
```

## Causal Inference with Models

So, correct adjustments to reveal causal effect of  $T \rightsquigarrow Y$ :

df1, **adjust** for  $X$ ,  $\beta = 0.5$ :

```
## # A tibble: 1 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df1  exposure    0.501    0.108
```

df2, **do not adjust** for  $X$ ,  $\beta = 1$ :

```
## # A tibble: 1 x 4
##   data term      estimate std.error
##   <chr> <chr>      <dbl>    <dbl>
## 1 df2  exposure    1.00    0.0841
```

(Data from D'Agostino McGowan (2023))

# Causal Inference with Models

- Importance of identifying “pre-treatment covariates”, “proper covariates”; doing “design before analysis”

# Causal Inference with Models

- ▶ Importance of identifying “pre-treatment covariates”, “proper covariates”; doing “design before analysis”
- ▶ Importance of experiments: strong knowledge about (part of) causal structure – assign mechanism

# Causal Inference with Models

- ▶ Importance of identifying “pre-treatment covariates”, “proper covariates”; doing “design before analysis”
- ▶ Importance of experiments: strong knowledge about (part of) causal structure – assign mechanism
- ▶ Causal inference is critical to scientific questions, and separate from prediction

# Causal Inference with Models

- ▶ Importance of identifying “pre-treatment covariates”, “proper covariates”; doing “design before analysis”
- ▶ Importance of experiments: strong knowledge about (part of) causal structure – assign mechanism
- ▶ Causal inference is critical to scientific questions, and separate from prediction
- ▶ Though, methods from prediction can aid causal inference (see, especially, *causal forests*)

# Causal Inference with Models

- ▶ Importance of identifying “pre-treatment covariates”, “proper covariates”; doing “design before analysis”
- ▶ Importance of experiments: strong knowledge about (part of) causal structure – assign mechanism
- ▶ Causal inference is critical to scientific questions, and separate from prediction
- ▶ Though, methods from prediction can aid causal inference (see, especially, *causal forests*)
- ▶ “Causal euphemisms” don’t help (Hernán 2018)



# References I

- Aronow, Peter M., and Cyrus Samii. 2016. “Does Regression Produce Representative Estimates of Causal Effects?” *American Journal of Political Science* 60 (1): 250–67.
- Chattopadhyay, Raghabendra, and Esther Duflo. 2004. “Women as Policy Makers: Evidence from a Randomized Policy Experiment in India.” *Econometrica* 72 (5): 1409–43.
- D’Agostino McGowan, Lucy. 2023. *quartets: Datasets to Help Teach Statistics*. <https://r-causal.github.io/quartets/>.
- Freedman, David A. 2008. “On Regression Adjustments to Experimental Data.” *Advances in Applied Mathematics* 40: 180–93.
- Gerber, Alan S., Donald P. Green, and Christopher W. Larimer. 2008. “Social Pressure and Voter Turnout: Evidence from a Large-Scale Field Experiment.” *American Political Science Review* 102 (1): 33–48.
- Green, Donald P., and Peter M. Aronow. 2011. “Analyzing Experimental Data Using Regression: When Is Bias a Practical Concern?” *SSRN Working Paper*. [https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=1466886](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=1466886).

# References II

- Hernán, Miguel A. 2018. “The c-Word: Scientific Euphemisms Do Not Improve Causal Inference from Observational Data.” *American Journal of Public Health* 108 (5): 616–19.
- Hernán, Miguel A., John Hsu, and Brian Healy. 2019. “A Second Chance to Get Causal Inference Right: A Classification of Data Science Tasks.” *CHANCE* 32 (1): 42–49. <https://doi.org/10.1080/09332480.2019.1579578>.
- Lin, Winston. 2013. “Agnostic Notes on Regression Adjustments to Experimental Data: Reexamining Freedman’s Critique.” *Annals of Applied Statistics*, 295–318. <https://doi.org/10.1214/12-AOAS583>.