

Randomized Experiments with Clusters

Ryan T. Moore

American University

2024-08-21

Table of contents I

Clusters in Experiments

Variance

Bias

An Example

Clusters in Experiments

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum
 - ▶ Schools (not teachers or students) to new curriculum

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum
 - ▶ Schools (not teachers or students) to new curriculum
 - ▶ Media markets (not viewers) to ad campaign

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum
 - ▶ Schools (not teachers or students) to new curriculum
 - ▶ Media markets (not viewers) to ad campaign
 - ▶ Clinic catchment areas (not patients) to health infrastructure spending

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum
 - ▶ Schools (not teachers or students) to new curriculum
 - ▶ Media markets (not viewers) to ad campaign
 - ▶ Clinic catchment areas (not patients) to health infrastructure spending
- ▶ These groups of individuals are *clusters*

Clusters in Experiments

- ▶ So far, *individual units* assigned to treatment
- ▶ But, sometimes, *groups* of units assigned to treatment.
- ▶ Consider assigning ...
 - ▶ Classrooms (not students) to new curriculum
 - ▶ Schools (not teachers or students) to new curriculum
 - ▶ Media markets (not viewers) to ad campaign
 - ▶ Clinic catchment areas (not patients) to health infrastructure spending
- ▶ These groups of individuals are *clusters*
- ▶ (Let's start with easy case: all clusters same size ...)
(Gerber and Green 2012)

Variance

Variance

- ▶ CLT SE for diff in means for clustered design:

Variance

► CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE large?

- ▶ Dividing by $k \equiv$ number of *clusters*

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE large?

- ▶ Dividing by $k \equiv$ number of *clusters*
- ▶ So, small number of clusters \leadsto larger SE

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE large?

- ▶ Dividing by $k \equiv$ number of *clusters*
- ▶ So, small number of clusters \leadsto larger SE
- ▶ Small clusters (more variability in means)

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE large?

- ▶ Dividing by $k \equiv$ number of *clusters*
- ▶ So, small number of clusters \leadsto larger SE
- ▶ Small clusters (more variability in means)
- ▶ If clusters differ in potential outcomes \leadsto larger SE

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE large?

- ▶ Dividing by $k \equiv$ number of *clusters*
- ▶ So, small number of clusters \leadsto larger SE
- ▶ Small clusters (more variability in means)
- ▶ If clusters differ in potential outcomes \leadsto larger SE
- ▶ I.e., if clusters meaningful \leadsto larger SE

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE small?

- ▶ Many clusters

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE small?

- ▶ Many clusters
- ▶ If clusters look like each other, \leadsto smaller SE

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE small?

- ▶ Many clusters
- ▶ If clusters look like each other, \leadsto smaller SE
- ▶ $\text{Var}(\overline{Y}_j(1))$: variance in cluster averages of Y_1

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE small?

- ▶ Many clusters
- ▶ If clusters look like each other, \leadsto smaller SE
- ▶ $\text{Var}(\overline{Y}_j(1))$: variance in cluster averages of Y_1
- ▶ $\text{Var}(\overline{Y}_j(0))$: variance in cluster averages of Y_0

Variance

- ▶ CLT SE for diff in means for clustered design:

$$SE(\widehat{ATE}) = \sqrt{\frac{1}{k-1} \left[\frac{m \text{Var}(\overline{Y}_j(0))}{N-m} + \frac{(N-m) \text{Var}(\overline{Y}_j(1))}{m} + 2 \text{Cov}(\overline{Y}_j(0), \overline{Y}_j(1)) \right]}$$

What might make this SE small?

- ▶ Many clusters
- ▶ If clusters look like each other, \leadsto smaller SE
- ▶ $\text{Var}(\overline{Y}_j(1))$: variance in cluster averages of Y_1
- ▶ $\text{Var}(\overline{Y}_j(0))$: variance in cluster averages of Y_0
- ▶ If $\text{Var}(\overline{Y}_j(1))$, $\text{Var}(\overline{Y}_j(0))$ are small, SE is small

Bias

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1, Y_0

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1, Y_0
 - ▶ Classrooms differ in number of students, and larger classes have lower test scores

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1 , Y_0
 - ▶ Classrooms differ in number of students, and larger classes have lower test scores
 - ▶ Media markets vary in number of viewers, and voters in cities less likely to turn out

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1, Y_0
 - ▶ Classrooms differ in number of students, and larger classes have lower test scores
 - ▶ Media markets vary in number of viewers, and voters in cities less likely to turn out
 - ▶ Health clusters vary in patient population, and larger clusters are healthier

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1, Y_0
 - ▶ Classrooms differ in number of students, and larger classes have lower test scores
 - ▶ Media markets vary in number of viewers, and voters in cities less likely to turn out
 - ▶ Health clusters vary in patient population, and larger clusters are healthier
- ▶ In this case, individual difference in means estimator is *biased* for true ATE

Bias

- ▶ Consider case where clusters vary in size, and size correlates with Y_1 , Y_0
 - ▶ Classrooms differ in number of students, and larger classes have lower test scores
 - ▶ Media markets vary in number of viewers, and voters in cities less likely to turn out
 - ▶ Health clusters vary in patient population, and larger clusters are healthier
- ▶ In this case, individual difference in means estimator is *biased* for true ATE
 - ▶ (Different assignments of clusters will produce different counts of treated units!)

Bias

Alternative estimator: Difference in totals (not means)

$$\widehat{ATE} = \frac{k_T + k_C}{N} \left(\frac{\sum Y_i(1)|T_i = 1}{k_T} - \frac{\sum Y_i(0)|T_i = 0}{k_C} \right)$$

Alternative estimator: Difference in totals (not means)

$$\widehat{ATE} = \frac{k_T + k_C}{N} \left(\frac{\sum Y_i(1) | T_i = 1}{k_T} - \frac{\sum Y_i(0) | T_i = 0}{k_C} \right)$$

- ▶ But, doesn't include cluster sizes, so may be high variance

Alternative estimator: Difference in totals (not means)

$$\widehat{ATE} = \frac{k_T + k_C}{N} \left(\frac{\sum Y_i(1) | T_i = 1}{k_T} - \frac{\sum Y_i(0) | T_i = 0}{k_C} \right)$$

- ▶ But, doesn't include cluster sizes, so may be high variance
- ▶ I.e., more students per classroom doesn't help precision here (unlike above)

An Example

An Example

- ▶ 1000 voters, split across 10 cities

An Example

- ▶ 1000 voters, split across 10 cities
- ▶ Measure ideology, which varies by city and voter

An Example

- ▶ 1000 voters, split across 10 cities
- ▶ Measure ideology, which varies by city and voter
- ▶ Do 2 hypothetical assignments

An Example

- ▶ 1000 voters, split across 10 cities
- ▶ Measure ideology, which varies by city and voter
- ▶ Do 2 hypothetical assignments
 - ▶ Individual-level assignment to news

An Example

- ▶ 1000 voters, split across 10 cities
- ▶ Measure ideology, which varies by city and voter
- ▶ Do 2 hypothetical assignments
 - ▶ Individual-level assignment to news
 - ▶ City-level cluster assignment to news

An Example

- ▶ 1000 voters, split across 10 cities
- ▶ Measure ideology, which varies by city and voter
- ▶ Do 2 hypothetical assignments
 - ▶ Individual-level assignment to news
 - ▶ City-level cluster assignment to news
- ▶ Use RI to find SE in each case

An Example

```
library(fabricatr)
library(randomizr)
library(tidyverse)

# Make data and visualise

set.seed(95852894)

voters <- fabricate(
  N = 1000,
  city_id = rep(1:10, 100),
  ideology = draw_normal_icc(mean = 0, N = N,
                             clusters = city_id, ICC = 0.7)
  city_id_fac = as.factor(city_id),
  city_id_fac = fct_reorder(city_id_fac, ideology)
)
```

An Example

```
table(voters$city_id)
```

1	2	3	4	5	6	7	8	9	10
100	100	100	100	100	100	100	100	100	100

An Example


```
table(voters$city_id)
```

```
 1    2    3    4    5    6    7    8    9   10  
100 100 100 100 100 100 100 100 100 100
```

```
voters |> group_by(city_id) |>  
  summarise(mean_ideo = mean(ideology)) |>  
  arrange(mean_ideo)
```

```
# A tibble: 10 x 2  
  city_id mean_ideo  
    <int>    <dbl>  
1       4    -2.90  
2       7    -1.94  
3       1    -1.68  
4       6    -1.10  
5       5    -0.903  
6       8    -0.686  
7      10    -0.161  
8       3     0.267  
9       2     1.46  
10      9     2.60
```

An Example



An Example

Assignments

```
voters <- voters |>
  mutate(
    tr_ind = sample(rep(0:1, nrow(voters) / 2)),
    tr_cl = cluster_ra(clusters = voters$city_id, m = 5))
```

An Example

Assignments

```
voters <- voters |>  
  mutate(  
    tr_ind = sample(rep(0:1, nrow(voters) / 2)),  
    tr_cl = cluster_ra(clusters = voters$city_id, m = 5))
```

```
voters |> count(tr_ind)
```

	tr_ind	n
1	0	500
2	1	500

```
voters |> count(tr_cl)
```

	tr_cl	n
1	0	500
2	1	500

An Example

Conditions by city, individual-level assignment:

```
table(voters$city_id, voters$tr_ind)
```

	0	1
1	48	52
2	47	53
3	52	48
4	47	53
5	51	49
6	49	51
7	49	51
8	53	47
9	48	52
10	56	44

An Example

Conditions by city, cluster-level assignment:

```
table(voters$city_id, voters$tr_cl)
```

	0	1
1	0	100
2	100	0
3	100	0
4	0	100
5	100	0
6	0	100
7	0	100
8	0	100
9	100	0
10	100	0

An Example


Draw responses under each assignment:

```
voters <- voters |>
  mutate(
    response_ind = draw_likert(x = ideology + tr_ind, m
    response_cl = draw_likert(x = ideology + tr_cl, m
```

An Example

Draw responses under each assignment:

```
voters <- voters |>
  mutate(
    response_ind = draw_likert(x = ideology + tr_ind, m = 10),
    response_cl = draw_likert(x = ideology + tr_cl, m = 10)
```



An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments

An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments
- ▶ Do 1000 cluster-level hypothetical assignments

An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments
- ▶ Do 1000 cluster-level hypothetical assignments
- ▶ Use sharp null $H_0 : \tau_i = 0$ to impute unobserved potential outcomes

An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments
- ▶ Do 1000 cluster-level hypothetical assignments
- ▶ Use sharp null $H_0 : \tau_i = 0$ to impute unobserved potential outcomes
- ▶ Calculate \widehat{ATE} in each case

An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments
- ▶ Do 1000 cluster-level hypothetical assignments
- ▶ Use sharp null $H_0 : \tau_i = 0$ to impute unobserved potential outcomes
- ▶ Calculate \widehat{ATE} in each case
- ▶ Compare distribution of $\widehat{ATE}_{\text{Individual}}$ to distribution of $\widehat{ATE}_{\text{Cluster}}$

An Example

Analyse precision with RI:

- ▶ Do 1000 individual-level hypothetical assignments
- ▶ Do 1000 cluster-level hypothetical assignments
- ▶ Use sharp null $H_0 : \tau_i = 0$ to impute unobserved potential outcomes
- ▶ Calculate \widehat{ATE} in each case
- ▶ Compare distribution of $\widehat{ATE}_{\text{Individual}}$ to distribution of $\widehat{ATE}_{\text{Cluster}}$
- ▶ With partner, sketch the two randomization distributions!

An Example

Analyse precision with RI:

```
n_sims <- 1000
df_ates <- tibble(ate_ind = NA,
                  ate_cl = NA)

for(idx in 1:n_sims){

  voters <- voters |>
    mutate(
      hyp_tr_individ = sample(rep(0:1, nrow(voters) / 2)),
      hyp_tr_cl = cluster_ra(clusters = voters$city_id, m = 5))

  ate_ind <- mean(voters$response_ind[voters$hyp_tr_ind == 1]) -
    mean(voters$response_ind[voters$hyp_tr_ind == 0])

  ate_cl <- mean(voters$response_cl[voters$hyp_tr_cl == 1]) -
    mean(voters$response_cl[voters$hyp_tr_cl == 0])


  df_ates[idx, "ate_ind"] <- ate_ind
  df_ates[idx, "ate_cl"] <- ate_cl
}
```

An Example

Analyse precision with RI:


An Example

Analyse precision with RI:



An Example

Analyse precision with RI:



```
# A tibble: 1 x 2  
  se_ind se_cl  
    <dbl> <dbl>  
1 0.0614 0.408
```

Next:

Regression and Experiments

`rtm@american.edu`
`www.ryantmoore.org`

References I

Gerber, Alan S., and Donald P. Green. 2012. *Field Experiments: Design, Analysis, and Interpretation*. New York, NY: WW Norton.