

# Randomization (Design-based) Inference for Experiments

Ryan T. Moore

American University

2024-08-20

# Table of contents I

An Exercise

Overview of Randomization Inference

Counting Principles

Examples

## An Exercise

A volunteer?

# A volunteer?

The task: select the 2 folders with messages

# A volunteer?

The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?

# A volunteer?

The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?
  - ▶ “No x-ray vision. No ESP. Effect of messages on choice = 0.”

# A volunteer?

The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?
  - ▶ “No x-ray vision. No ESP. Effect of messages on choice = 0.”
- ▶ What is an alternative?



# A volunteer?

The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?
  - ▶ “No x-ray vision. No ESP. Effect of messages on choice = 0.”
- ▶ What is an alternative?
  - ▶ “Some way to detect messages. Message location  $\rightarrow$  choice.”

# A volunteer?

The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?
  - ▶ “No x-ray vision. No ESP. Effect of messages on choice = 0.”
- ▶ What is an alternative?
  - ▶ “Some way to detect messages. Message location  $\rightarrow$  choice.”

# A volunteer?


The task: select the 2 folders with messages

- ▶ What is our baseline expectation/model for this process?
  - ▶ “No x-ray vision. No ESP. Effect of messages on choice = 0.”
- ▶ What is an alternative?
  - ▶ “Some way to detect messages. Message location  $\rightarrow$  choice.”

Select!


# Randomization Inference

The possible choices:



# Randomization Inference


The possible choices:



► You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.

# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?

# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$

# Randomization Inference

The possible choices:




- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?



# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?  $\frac{7}{10} = 0.7$

# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?  $\frac{7}{10} = 0.7$
- ▶ What is “prob result at least this extreme, given model of no effect”?

# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?  $\frac{7}{10} = 0.7$
- ▶ What is “prob result at least this extreme, given model of no effect”?
- ▶ Definition of  $p$ -value!

# Randomization Inference


The possible choices:



- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?  $\frac{7}{10} = 0.7$
- ▶ What is “prob result at least this extreme, given model of no effect”?
- ▶ Definition of  $p$ -value!
- ▶ Valid, exact, with no distributional assumption, no large  $n$ .


# Randomization Inference

The possible choices:




- ▶ You chose \_\_\_\_ and \_\_\_\_\_. Let  $X$  = number found.
- ▶ What was  $P(X \geq 2 | \text{no ESP})$ ?  $\frac{1}{10} = 0.1$
- ▶ What was  $P(X \geq 1 | \text{no ESP})$ ?  $\frac{7}{10} = 0.7$
- ▶ What is “prob result at least this extreme, given model of no effect”?
- ▶ Definition of  $p$ -value!
- ▶ Valid, exact, with no distributional assumption, no large  $n$ .
- ▶ *Randomization* creates dist’n of possible numbers correct


# The Randomization Distribution of $X$



# The Randomization Distribution of $X$



# The Randomization Distribution of $X$





# Parametric Null Hypothesis Significance Testing

- ▶ Specify and assume  $H_0$
- ▶ Define  $H_A$
- ▶ Examine reference dist'n ( $t$ ,  $\chi^2$ , ...) under  $H_0$
- ▶ Calculate  $p$ -value
- ▶ Compare to some  $\alpha$ ; reject  $H_0$  if  $p < \alpha$

# Overview of Randomization Inference

## Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)

# Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)
- ▶ Define  $H_A$

## Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)
- ▶ Define  $H_A$
- ▶ Create reference dist'n from all possible values of  $X$  under  $H_0$   
(or at least a big sample of them)

## Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)
- ▶ Define  $H_A$
- ▶ Create reference dist'n from all possible values of  $X$  under  $H_0$   
(or at least a big sample of them)
- ▶ What prop. of possible values are “at least as extreme as” observed?  
 $\leadsto$   $p$ -value!

## Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)
- ▶ Define  $H_A$
- ▶ Create reference dist'n from all possible values of  $X$  under  $H_0$   
(or at least a big sample of them)
- ▶ What prop. of possible values are “at least as extreme as” observed?  
 $\leadsto$   $p$ -value!
- ▶ Compare to some  $\alpha$ ; reject  $H_0$  if  $p < \alpha$

## Randomization Inference

- ▶ Specify and assume  $H_0$   
(sharp null of no treatment effect)
- ▶ Define  $H_A$
- ▶ Create reference dist'n from all possible values of  $X$  under  $H_0$   
(or at least a big sample of them)
- ▶ What prop. of possible values are “at least as extreme as” observed?  
 $\leadsto$   $p$ -value!
- ▶ Compare to some  $\alpha$ ; reject  $H_0$  if  $p < \alpha$
- ▶ CA ballot ordering effects (JASA 2006)



# Randomization Inference

The RI  $p$ -value is

$$p = \frac{\# \text{ outcomes } \geq \text{as extreme as obs}}{\text{total } \# \text{ outcomes}}$$

# Randomization Inference

The RI  $p$ -value is

$$p = \frac{\# \text{ outcomes } \geq \text{as extreme as obs}}{\text{total } \# \text{ outcomes}}$$

or

$$p = \frac{\# \text{ randomizations producing extreme } \widehat{ATE}}{\text{total } \# \text{ randomizations}}$$

# Randomization Inference

The RI  $p$ -value is

$$p = \frac{\# \text{ outcomes } \geq \text{as extreme as obs}}{\text{total } \# \text{ outcomes}}$$

or

$$p = \frac{\# \text{ randomizations producing extreme } \widehat{ATE}}{\text{total } \# \text{ randomizations}}$$

How many randomizations are there?

# Counting Principles

## Combinations: Counting selected sets

How many ways to **select**  $k$  things from a set of  $n$  things?

$${}_nC_k = \binom{n}{k} = \frac{{}_nP_k}{k!} = \frac{n!}{k!(n-k)!}$$

## Combinations: Counting selected sets

How many ways to **select**  $k$  things from a set of  $n$  things?

$${}_nC_k = \binom{n}{k} = \frac{{}_nP_k}{k!} = \frac{n!}{k!(n-k)!}$$

Suppose 5 units,  $A, B, C, D, E$ .

- How many ways to order?  ${}_nP_k$ :  $ABCDE, ABCED,$   
... =  $5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5! = 120$

## Combinations: Counting selected sets

How many ways to **select**  $k$  things from a set of  $n$  things?

$${}_nC_k = \binom{n}{k} = \frac{{}_nP_k}{k!} = \frac{n!}{k!(n-k)!}$$

Suppose 5 units,  $A, B, C, D, E$ .

- ▶ How many ways to order?  ${}_nP_k$ :  $ABCDE, ABCED, \dots = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5! = 120$
- ▶ If only want 3 of 5? Divide by  $2! = (n - k)!$   
(removing permutations from last 2 slots)

## Combinations: Counting selected sets

How many ways to **select**  $k$  things from a set of  $n$  things?

$${}_nC_k = \binom{n}{k} = \frac{{}_nP_k}{k!} = \frac{n!}{k!(n-k)!}$$

Suppose 5 units,  $A, B, C, D, E$ .

- ▶ How many ways to order?  ${}_nP_k$ :  $ABCDE, ABCED, \dots = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5! = 120$
- ▶ If only want 3 of 5? Divide by  $2! = (n - k)!$   
(removing permutations from last 2 slots)
- ▶ What if order doesn't matter? Divide by  $3! = k!$   
(6 permutations for  $ABC$ , but only one combination)



## Combinations: Counting selected sets

How many ways to choose 5 villages of 10 for treatment?

## Combinations: Counting selected sets

How many ways to choose 5 villages of 10 for treatment?

$${}_{10}C_5 = \binom{10}{5} = \frac{10!}{5!(10-5)!}$$

## Combinations: Counting selected sets

How many ways to choose 5 villages of 10 for treatment?

$${}_{10}C_5 = \binom{10}{5} = \frac{10!}{5!(10-5)!}$$

$$\frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = 252$$

# Common Assumptions, Null Hypotheses

- ▶ Constant effect:

$$\tau_i = Y_{i1} - Y_{i0} = \tau \quad \forall i$$

- ▶ Null hypothesis of no average effect:

$$ATE = \bar{\tau} = 0$$

- ▶ Sharp null hypothesis of no effect:

$$\tau_i = 0$$

## Examples

## An Assignment Mechanism: Perfect Doctor

Calculate RI  $p$ -value for Perfect Doctor, under sharp null.

Patient	Y(0)	Y(1)	$\tau$	T
1	(1)	6	(5)	1
2	(3)	12	(9)	1
3	9	(8)	(-1)	0
4	11	(10)	(-1)	0
Mean	10	9	(3)	

## An Assignment Mechanism: Perfect Doctor

Calculate RI  $p$ -value for Perfect Doctor, under sharp null.

Patient	Y(0)	Y(1)	$\tau$	T
1	(1)	6	(5)	1
2	(3)	12	(9)	1
3	9	(8)	(-1)	0
4	11	(10)	(-1)	0
Mean	10	9	(3)	

(See 01-ri-perfect-dr.R)

## RI versus the $t$ -test

Perfect Doctor:

- ▶ RI:  $p = 1$
- ▶ `t.test()`:  $p \approx 0.8$
- ▶ “If no tr effect, then this result typical”



## RI versus the $t$ -test

Perfect Doctor:

- ▶ RI:  $p = 1$
- ▶ `t.test()`:  $p \approx 0.8$
- ▶ “If no tr effect, then this result typical”

(Odd logic of NHST: “assume false thing, how strange is data?”)

## Randomization Inference, Example 2

Assumed table of potential outcomes:

Village	$T$	% if Female Head, $Y(1)$	% if Male Head, $Y(0)$	$\tau_i$
1	—	15	10	5
2	—	15	15	0
3	—	30	20	10
4	—	15	20	-5
5	—	20	10	10
6	—	15	15	0
7	—	30	15	15
Average		20	15	5

## Randomization Inference, Example 2

Suppose we randomly select 2 villages to have female-headed councils, and observe

Village	$T$	% if Female Head, $Y(1)$	% if Male Head, $Y(0)$	$\tau_i$
1	F	15		
2	M		15	
3	M		20	
4	M		20	
5	M		10	
6	M		15	
7	F	30		
Average		22.5	16	6.5

## Randomization Inference, Example 2

We assume the *sharp null* hypothesis (assumed values in red):

Village	$T$	% if Female Head, $Y(1)$	% if Male Head, $Y(0)$	$\tau_i$
1	F	15	15	0
2	M	15	15	0
3	M	20	20	0
4	M	20	20	0
5	M	10	10	0
6	M	15	15	0
7	F	30	30	0
Average				0

## Randomization Inference, Example 2

Then we estimate what the observed ATE would be for all the possible random assignments.

First,

Village	$T$	% if Female Head, $Y(1)$	% if Male Head, $Y(0)$	$\tau_i$
1	F	15		
2	M		15	
3	M		20	
4	M		20	
5	M		10	
6	F	15		
7	M		30	
Average		15	19	-4

## Randomization Inference, Example 2

Second,

Village	$T$	% if Female Head, $Y(1)$	% if Male Head, $Y(0)$	$\tau_i$
1	F	15		
2	M		15	
3	M		20	
4	M		20	
5	F	10		
6	M		15	
7	M		30	
Average		12.5	20	-7.5

## Randomization Inference, Example 2

...

## Randomization Inference, Example 2

..., and all the others. The full set of  $\frac{7!}{2!5!} = 21$  differences in means:

Estimate	Frequency
-7.5	3
-4	5
-0.5	6
3	2
6.5	3
10	2
Total	21



## Randomization Inference, Example 2

..., and all the others. The full set of  $\frac{7!}{2!5!} = 21$  differences in means:

Estimate	Frequency
-7.5	3
-4	5
-0.5	6
3	2
6.5	3
10	2
Total	21

How many are at least as extreme as my 6.5?

## Randomization Inference, Example 2

..., and all the others. The full set of  $\frac{7!}{2!5!} = 21$  differences in means:

Estimate	Frequency
-7.5	3
-4	5
-0.5	6
3	2
6.5	3
10	2
Total	21

How many are at least as extreme as my 6.5?

Two-sided (“women’s %  $\neq$  men’s”):  $p = \frac{8}{21} \approx 0.38$

## Randomization Inference, Example 2

..., and all the others. The full set of  $\frac{7!}{2!5!} = 21$  differences in means:

Estimate	Frequency
-7.5	3
-4	5
-0.5	6
3	2
6.5	3
10	2
Total	21

How many are at least as extreme as my 6.5?

Two-sided (“women’s %  $\neq$  men’s”):  $p = \frac{8}{21} \approx 0.38$

One-sided (“women’s %  $>$  men’s”):  $p = \frac{5}{21} \approx 0.24$

# Randomization Inference

These  $p$ -values

▶ are exact, not approximate

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )
- ▶ do not require calculating standard error (SE)

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )
- ▶ do not require calculating standard error (SE)
- ▶ *Randomization* creates distribution of possible numbers correct/treatment effects



## Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )
- ▶ do not require calculating standard error (SE)
- ▶ *Randomization* creates distribution of possible numbers correct/treatment effects
- ▶ can be approximated by sampling

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )
- ▶ do not require calculating standard error (SE)
- ▶ *Randomization* creates distribution of possible numbers correct/treatment effects
- ▶ can be approximated by sampling
  - ▶ Assigning 500 of 1000 respondents to treatment? Randomly generate 10,000 assignments from  $2.7 \times 10^{299}$  possible ...

# Randomization Inference

These  $p$ -values

- ▶ are exact, not approximate
- ▶ do not require asymptotic large  $N \rightarrow \infty$
- ▶ need no distrib assump (normal as  $N \rightarrow \infty$ )
- ▶ do not require calculating standard error (SE)
- ▶ *Randomization* creates distribution of possible numbers correct/treatment effects
- ▶ can be approximated by sampling
  - ▶ Assigning 500 of 1000 respondents to treatment? Randomly generate 10,000 assignments from  $2.7 \times 10^{299}$  possible ...
- ▶ can be computationally intensive

## Randomization Inference

- ▶ Resume audit study, Bertrand and Mullainathan (2004)

	0	1
black	2278	157
white	2200	235

## Randomization Inference

- ▶ Resume audit study, Bertrand and Mullainathan (2004)

	0	1
black	2278	157
white	2200	235

- ▶ Only possible values:  $\tau_i \in \{-1, 0, 1\}$

## Randomization Inference

- ▶ Resume audit study, Bertrand and Mullainathan (2004)

	0	1
black	2278	157
white	2200	235

- ▶ Only possible values:  $\tau_i \in \{-1, 0, 1\}$

```
# A tibble: 2 x 2
  race  call_rate
<chr>    <dbl>
1 black    0.0645
2 white    0.0965
```

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$



- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

$${}_{4870}C_{2435} = \binom{4870}{2435} = \frac{4870 \cdot 4869 \cdot \dots \cdot 2436}{2435!}$$

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

$${}_{4870}C_{2435} = \binom{4870}{2435} = \frac{4870 \cdot 4869 \cdot \dots \cdot 2436}{2435!}$$

$$\approx 1.1 \times 10^{1464}$$

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

$${}_{4870}C_{2435} = \binom{4870}{2435} = \frac{4870 \cdot 4869 \cdot \dots \cdot 2436}{2435!}$$

$$\approx 1.1 \times 10^{1464}$$

(There are  $\approx 10^{86}$  fundamental particles in the universe.)

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments

$${}_{4870}C_{2435} = \binom{4870}{2435} = \frac{4870 \cdot 4869 \cdot \dots \cdot 2436}{2435!}$$

$$\approx 1.1 \times 10^{1464}$$

(There are  $\approx 10^{86}$  fundamental particles in the universe.)

- ▶ Let's do 1000, or 100,000 – something reasonable

- ▶ Assume the sharp null  $\tau_i = 0$  for every employer.
- ▶  $H_0 : \mu_{\text{black name}} = \mu_{\text{white name}}$
- ▶  $H_A : \mu_{\text{black name}} \neq \mu_{\text{white name}}$
- ▶ Create reference dist'n of all possible assignments


$${}_{4870}C_{2435} = \binom{4870}{2435} = \frac{4870 \cdot 4869 \cdot \dots \cdot 2436}{2435!}$$


$$\approx 1.1 \times 10^{1464}$$

(There are  $\approx 10^{86}$  fundamental particles in the universe.)

- ▶ Let's do 1000, or 100,000 – something reasonable
- ▶ See `01-ri-resume-donate.R`









## Randomization Inference

- ▶ Gerber and Green (2012) donations example, p. 65
- ▶ Possible values  $\tau_i \in (-\infty, \infty)$
- ▶  $Y_1, Y_0, \tau$  likely very skewed
- ▶ See `01-ri-resume-donate.R`





# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

Create RI confidence ints (“invert the randomization test”):

► Posit  $H_0 : \tau = \tau^* \in \{\dots, -2, -1, 0, 1, 2, \dots\}$

# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

Create RI confidence ints (“invert the randomization test”):

- ▶ Posit  $H_0 : \tau = \tau^* \in \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ RI test whether to reject  $H_0$



# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

Create RI confidence ints (“invert the randomization test”):

- ▶ Posit  $H_0 : \tau = \tau^* \in \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ RI test whether to reject  $H_0$
- ▶ If not, then  $\tau^*$  is in CI

# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

Create RI confidence ints (“invert the randomization test”):

- ▶ Posit  $H_0 : \tau = \tau^* \in \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ RI test whether to reject  $H_0$
- ▶ If not, then  $\tau^*$  is in CI
- ▶ CI consists of set of  $\tau^*$  not unusual, given data

# The RI Confidence Interval

Recall that

“reject  $H_0$  at  $\alpha = 0.05$ ”  $\equiv$  “ $H_0$  falls outside 95% CI”

Create RI confidence ints (“invert the randomization test”):

- ▶ Posit  $H_0 : \tau = \tau^* \in \{\dots, -2, -1, 0, 1, 2, \dots\}$
- ▶ RI test whether to reject  $H_0$
- ▶ If not, then  $\tau^*$  is in CI
- ▶ CI consists of set of  $\tau^*$  not unusual, given data
- ▶ See `01-ri-resume-donate.R`

We can adapt this testing procedure for any quantity we want to test!

We can adapt this testing procedure for any quantity we want to test!

► Difference in variances

We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient

We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient
- ▶ Sum of two regression coefficients

We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient
- ▶ Sum of two regression coefficients
- ▶ Ratio of skewnesses



We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient
- ▶ Sum of two regression coefficients
- ▶ Ratio of skewnesses
- ▶ ...(any crazy thing!)

We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient
- ▶ Sum of two regression coefficients
- ▶ Ratio of skewnesses
- ▶ ...(any crazy thing!)

We can adapt this testing procedure for any quantity we want to test!

- ▶ Difference in variances
- ▶ Regression coefficient
- ▶ Sum of two regression coefficients
- ▶ Ratio of skewnesses
- ▶ ...(any crazy thing!)

No need to derive the correct asymptotic standard error (SE).

“There is only one test.”

“There is only one test.”

- Allen Downey, posit::conf(2024)

Next:

# Covariates in Experiments

- Bertrand, Marianne, and Sendhil Mullainathan. 2004. “Are Emily and Greg More Employable Than Lakisha and Jamal? A Field Experiment on Labor Market Discrimination.” *American Economic Review* 94 (4): 991–1013.
- Gerber, Alan S., and Donald P. Green. 2012. *Field Experiments: Design, Analysis, and Interpretation*. New York, NY: WW Norton.