

Potential Outcomes and Randomised Experiments

Ryan T. Moore

American University

2026-07-21

Table of contents I

Preliminaries

Statistical Refreshment

Assignment of Treatment

Preliminaries

Preliminaries

- ▶ Accept GitHub invitation (click link in email)
- ▶ Upload `.txt` or `.R` or `.qmd` or `.Rmd` of solutions
(`.pdf`, but comments)
- ▶ How `.qmd/.Rmd` works (more to come?)

Preliminaries

- ▶ Accept GitHub invitation (click link in email)
- ▶ Upload `.txt` or `.R` or `.qmd` or `.Rmd` of solutions
(`.pdf`, but comments)
- ▶ How `.qmd`/`.Rmd` works (more to come?)
- ▶ PS1 for today

Preliminaries

- ▶ Accept GitHub invitation (click link in email)
- ▶ Upload `.txt` or `.R` or `.qmd` or `.Rmd` of solutions
(`.pdf`, but comments)
- ▶ How `.qmd`/`.Rmd` works (more to come?)
- ▶ PS1 for today
- ▶ PS2 for Thursday

Preliminaries

- ▶ Accept GitHub invitation (click link in email)
- ▶ Upload `.txt` or `.R` or `.qmd` or `.Rmd` of solutions
(`.pdf`, but comments)
- ▶ How `.qmd/.Rmd` works (more to come?)
- ▶ PS1 for today
- ▶ PS2 for Thursday
- ▶ Work together, but every keystroke your own

Preliminaries

- ▶ Accept GitHub invitation (click link in email)
- ▶ Upload `.txt` or `.R` or `.qmd` or `.Rmd` of solutions
(`.pdf`, but comments)
- ▶ How `.qmd/.Rmd` works (more to come?)
- ▶ PS1 for today
- ▶ PS2 for Thursday
- ▶ Work together, but every keystroke your own
- ▶ Ask questions in the Slack

Review Questions

1. When can we observe both $Y_i(1)$ and $Y_i(0)$?
2. What is this fact called?
3. When does the empirical difference-in-means estimator exactly equal the true, underlying ATE?
4. What are the two parts of SUTVA?

Exercise in Potential Outcomes

Statistical Refreshment

Expectation

$$E(Y) = \sum_{i=1}^n [y_i \cdot p(y_i)]$$

Expectation

$$E(Y) = \sum_{i=1}^n [y_i \cdot p(y_i)]$$

Calculate $E(\text{Ideology})$.

Respondent	Ideology
1	3
2	-2
3	3
4	1
5	1

Expectation

$$E(Y) = \sum_{i=1}^n [y_i \cdot p(y_i)]$$

Calculate $E(\text{Ideology})$.

Respondent	Ideology
1	3
2	-2
3	3
4	1
5	1

$$1 \cdot \frac{2}{5} + 3 \cdot \frac{2}{5} + (-2) \cdot \frac{1}{5} = \frac{6}{5}$$

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$.

Respondent	Ideology	Dem
1	3	0
2	-2	1
3	3	0
4	1	0
5	1	1

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$.

Respondent	Ideology	Dem
1	3	0
2	-2	1
3	3	0
4	1	0
5	1	1

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$.

Respondent	Ideology	Dem
1	3	0
2	-2	1
3	3	0
4	1	0
5	1	1

$$1 \cdot \frac{1}{3} + 3 \cdot \frac{2}{3} = \frac{7}{3}$$

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$

		Dem?	
		0	1
Ideology	3	2	0
	1	1	1
	-2	0	1

(tabled data, cells: counts of units with row/column scores)

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$

		Dem?	
		0	1
Ideology	3	0.4	0.0
	1	0.2	0.2
	-2	0.0	0.2

(tabled data, cells: probability unit has row/column scores)

Conditional Expectation

$$E(Y|X = x) = \sum_{i=1}^n [y_i \cdot p(y_i|X = x)]$$

Calculate $E(\text{Ideology}|\text{Dem} = 0)$

		Dem?	
		0	1
Ideology	3	0.4	0.0
	1	0.2	0.2
	-2	0.0	0.2

(tabled data, cells: probability unit has row/column scores)

$$3 \cdot \frac{0.4}{0.6} + 1 \cdot \frac{0.2}{0.6} = \frac{7}{3}$$

Unbiasedness

Does estimator return the truth, on average?

Unbiasedness

Does estimator return the truth, on average?

$$E(\hat{\theta}) = \theta$$

Unbiasedness

Does estimator return the truth, on average?

$$E(\hat{\theta}) = \theta$$

$$\blacktriangleright E \left[\widehat{\overline{Y_1 - Y_0}} \right] = \overline{Y_1 - Y_0}$$

Unbiasedness

Does estimator return the truth, on average?

$$E(\hat{\theta}) = \theta$$

$$\blacktriangleright E \left[\widehat{\overline{Y_1 - Y_0}} \right] = \overline{Y_1 - Y_0}$$

$$\blacktriangleright E \left[\widehat{\beta_1} \right] = \beta_1$$

Unbiasedness

What estimator used to estimate $\overline{Y_1 - Y_0}$?

Unbiasedness

What estimator used to estimate $\overline{Y_1 - Y_0}$?

The difference-in-means estimator.

Unbiasedness

What estimator used to estimate $\overline{Y_1 - Y_0}$?

The difference-in-means estimator.

$$\widehat{\overline{Y_1 - Y_0}} = (Y_1|T_i = 1) - (Y_0|T_i = 0)$$

Unbiasedness

What estimator used to estimate $\overline{Y_1 - Y_0}$?

The difference-in-means estimator.

$$\widehat{\overline{Y_1 - Y_0}} = (Y_1|T_i = 1) - (Y_0|T_i = 0)$$

Is it a “good” estimator?

Unbiasedness

The difference-in-means estimator is *unbiased* for the true average treatment effect.

Unbiasedness

The difference-in-means estimator is *unbiased* for the true average treatment effect.

(See 02-unbiased.R)

Probability

The Three Axioms

1. $P(A) \geq 0$
2. $P(\Omega) = 1$
3. If events *mutually exclusive* (or, sets *disjoint*), then

$$P(A \text{ or } B) = P(A) + P(B)$$

Conditional Probability

For events A and B ,

$$\begin{aligned}P(A \text{ and } B) &= P(A)P(B|A) \\ &= P(B)P(A|B)\end{aligned}$$

Divide both sides by marginal probability $P(B)$ yields

$$P(A|B) = \frac{P(A \text{ and } B)}{P(B)}$$

Statistical Independence

A and B are independent iff both

$$P(A|B) = P(A)$$

and

$$P(B|A) = P(B)$$

Conditional Independence

A and B can be independent *conditional on* C .

Conditional Independence

A and B can be independent *conditional on* C .

Many times, *only* independent given C .

Conditional Independence

A and B can be independent *conditional on* C .

Many times, *only* independent given C .

E.g., Y_1, Y_0 indep of T , given X .

Conditional Independence

A and B can be independent *conditional on* C .

Many times, *only* independent given C .

E.g., Y_1, Y_0 indep of T , given X .

$$P(A \text{ and } B|C) = P(A|C)P(B|C)$$

Potential Outcomes Model: Estimands

- ▶ Individual TE

$$\tau_i = Y_i(1) - Y_i(0)$$

- ▶ Average treatment effect

$$ATE = E(Y_1 - Y_0) = E(Y_1) - E(Y_0)$$

- ▶ Average treatment effect for the treated

$$ATT = E(Y_1|T = 1) - E(Y_0|T = 1)$$

Potential Outcomes Model: Estimands

The average treatment effect (ATE):

$$\begin{aligned}E(Y_1 - Y_0) &= \frac{1}{2n} \sum_{i=1}^{2n} (Y_{i1} - Y_{i0}) \\&= \frac{1}{2n} \sum_{i=1}^{2n} (Y_{i1}) - \frac{1}{2n} \sum_{i=1}^{2n} (Y_{i0}) \\&= E(Y_1) - E(Y_0)\end{aligned}$$

Potential Outcomes Model: Estimands

- ▶ If we know (Y_1, Y_0) indep of T
- ▶ Then,

$$\begin{aligned}E(Y_1) &= E(Y_1|T = 1) \\E(Y_0) &= E(Y_0|T = 0)\end{aligned}$$

- ▶ Then, can substitute

$$\begin{aligned}ATE &= E(Y_1) - E(Y_0) \\&= E(Y_1|T = 1) - E(Y_0|T = 0)\end{aligned}$$

Potential Outcomes Model: Estimands

- ▶ If we know (Y_1, Y_0) indep of T
- ▶ Then,

$$\begin{aligned}E(Y_1) &= E(Y_1|T = 1) \\E(Y_0) &= E(Y_0|T = 0)\end{aligned}$$

- ▶ Then, can substitute

$$\begin{aligned}ATE &= E(Y_1) - E(Y_0) \\&= E(Y_1|T = 1) - E(Y_0|T = 0)\end{aligned}$$

Observed diff in Tr and Co group means gives ATE!

Potential Outcomes Model: Estimands

Holland (1986): “*prima facie effect*”:

$$E(Y_t|S = t) - E(Y_c|S = c)$$

Potential Outcomes Model: Estimands

Holland (1986): “*prima facie effect*”:

$$E(Y_t|S = t) - E(Y_c|S = c)$$

“It is important to recognize that $E(Y_t)$ and $E(Y_t|S = t)$ are *not* the same thing ...”

Potential Outcomes Model: Estimands, Interpretation

Gerber and Green (2012):

When $Y(1)$ and $Y(0)$ indep of T ,

$$ATE = E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 0)$$

Potential Outcomes Model: Estimands, Interpretation

Gerber and Green (2012):

When $Y(1)$ and $Y(0)$ indep of T ,

$$\begin{aligned}ATE &= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 0) \\ &= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1) + \\ &\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)\end{aligned}$$

Potential Outcomes Model: Estimands, Interpretation

Gerber and Green (2012):

When $Y(1)$ and $Y(0)$ indep of T ,

$$\begin{aligned}ATE &= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1) + \\&\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= [E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1)] + \\&\quad [E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)]\end{aligned}$$

Potential Outcomes Model: Estimands, Interpretation

Gerber and Green (2012):

When $Y(1)$ and $Y(0)$ indep of T ,

$$\begin{aligned}ATE &= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1) + \\&\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= [E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1)] + \\&\quad [E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)] \\&= E(Y_i(1) - Y_i(0)|T_i = 1) + \\&\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)\end{aligned}$$

Potential Outcomes Model: Estimands, Interpretation

Gerber and Green (2012):

When $Y(1)$ and $Y(0)$ indep of T ,

$$\begin{aligned}ATE &= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1) + \\&\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0) \\&= [E(Y_i(1)|T_i = 1) - E(Y_i(0)|T_i = 1)] + \\&\quad [E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)] \\&= E(Y_i(1) - Y_i(0)|T_i = 1) + \\&\quad E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)\end{aligned}$$

$$\underbrace{E(Y_i(1) - Y_i(0)|T_i = 1)}_{\text{ATT}} + \underbrace{E(Y_i(0)|T_i = 1) - E(Y_i(0)|T_i = 0)}_{\text{Selection Bias}}$$

Assignment of Treatment

What can be a Treatment?

- ▶ Timing clearly defined

What can be a Treatment?

- ▶ Timing clearly defined
- ▶ Covariates: causally prior to treatment

What can be a Treatment?

- ▶ Timing clearly defined
- ▶ Covariates: causally prior to treatment
- ▶ “Attributes”, “immutable characteristics”:
difficult to isolate

What can be a Treatment?

- ▶ Timing clearly defined
- ▶ Covariates: causally prior to treatment
- ▶ “Attributes”, “immutable characteristics”:
difficult to isolate

What can be a Treatment?

- ▶ Timing clearly defined
- ▶ Covariates: causally prior to treatment
- ▶ “Attributes”, “immutable characteristics”:
difficult to isolate

Sen and Wasow (2016) : “Race as a Bundle of Sticks: Designs that Estimate Effects of Seemingly Immutable Characteristics” (*elements* of attributes varyingly manipulable)

Attributes

“Causal effect of race”?

```
data(resume, package = "qss")  
dim(resume)
```

```
[1] 4870    4
```


Attributes

“Causal effect of race”?

```
data(resume, package = "qss")  
dim(resume)
```

```
[1] 4870    4
```

```
kable(table(resume$race, resume$call))
```

	0	1
black	2278	157
white	2200	235

Attributes

“Causal effect of race”?

```
data(resume, package = "qss")  
dim(resume)
```

```
[1] 4870    4
```

```
kable(table(resume$race, resume$call))
```

	0	1
black	2278	157
white	2200	235

Bertrand and Mullainathan (2004): “Are Emily and Greg More Employable Than Lakisha and Jamal? A Field Experiment on Labor Market Discrimination”

Random Assignment Mechanisms

- ▶ Simple/complete randomisation
(Bernoulli trial, prob π)

Random Assignment Mechanisms

- ▶ Simple/complete randomisation
(Bernoulli trial, prob π)
- ▶ Complete randomisation / random allocation
(fixed proportion to tr)

Random Assignment Mechanisms

- ▶ Simple/complete randomisation
(Bernoulli trial, prob π)
- ▶ Complete randomisation / random allocation
(fixed proportion to tr)
- ▶ Blocked randomisations
(fixed proportion to tr, w/in group)

Random Assignment Mechanisms

- ▶ Simple/complete randomisation
(Bernoulli trial, prob π)
- ▶ Complete randomisation / random allocation
(fixed proportion to tr)
- ▶ Blocked randomisations
(fixed proportion to tr, w/in group)
- ▶ Cluster randomisations
(assignment at higher level)

Random Assignment Mechanisms

- ▶ Simple/complete randomisation
(Bernoulli trial, prob π)
- ▶ Complete randomisation / random allocation
(fixed proportion to tr)
- ▶ Blocked randomisations
(fixed proportion to tr, w/in group)
- ▶ Cluster randomisations
(assignment at higher level)
- ▶ Waitlists, stepped wedges, etc.

The Potential Outcomes Model: Assignment

Observed outcome: $Y_i = Y_i(1) \cdot T_i + Y_i(0) \cdot (1 - T_i)$

The assignment mechanism *selects* which potential outcome we observe.

The Potential Outcomes Model: Assignment

Observed outcome: $Y_i = Y_i(1) \cdot T_i + Y_i(0) \cdot (1 - T_i)$

The assignment mechanism *selects* which potential outcome we observe.

(Gerber and Green (2012) use $Y_i = Y_i(1) \cdot d_i + Y_i(0) \cdot (1 - d_i)$ to highlight that we observe pot outcome from treatment **actually** taken, not hypothetical or assigned treatment.)

The Potential Outcomes Model: Assignment

“Assignment mechanisms” are really missing-data-generating procedures.

Ignorability

Assignment mechanism is *ignorable* if Y_{mis} conditionally indep of T

$$P(T|X, Y_{obs}, Y_{mis}) = P(T|X, Y_{obs})$$

Ignorability

Assignment mechanism is *ignorable* if Y_{mis} conditionally indep of T

$$P(T|X, Y_{obs}, Y_{mis}) = P(T|X, Y_{obs})$$

Nothing in unobserved Y_{mis} informs relationship betw'n Y_{obs} , T .

Unconfoundedness

Some ignorable mechanisms are *unconfounded*, too.

$$P(T|X, Y_{obs}, Y_{mis}) = P(T|X)$$

Unconfoundedness

Some ignorable mechanisms are *unconfounded*, too.

$$P(T|X, Y_{obs}, Y_{mis}) = P(T|X)$$

Nothing in Y informs T .

Unconfoundedness

Some ignorable mechanisms are *unconfounded*, too.

$$P(T|X, Y_{obs}, Y_{mis}) = P(T|X)$$

Nothing in Y informs T .

These are special (key!) cases of conditional independence.

An Assignment Mechanism

Little & Rubin (2000):

Patient	Y	T
1	6	1
2	12	1
3	9	0
4	11	0

An Assignment Mechanism

Little & Rubin (2000):

Patient	Y	T
1	6	1
2	12	1
3	9	0
4	11	0

Clearly, treatment is harmful. $\overline{Y(1)} - \overline{Y(0)} = 9 - 10 = -1$.

An Assignment Mechanism

Little & Rubin (2000):

Patient	Y(0)	Y(1)	τ	T
1		6		1
2		12		1
3	9			0
4	11			0
Mean	10	9		

Clearly, treatment is harmful.

$$\overline{Y(1)|T=1} - \overline{Y(0)|T=0} = 9 - 10 = -1$$

An Assignment Mechanism: Perfect Doctor

Little & Rubin (2000):

Patient	Y(0)	Y(1)	τ	T
1	(1)	6		1
2	(3)	12		1
3	9	(8)		0
4	11	(10)		0
Mean	10	9		

An Assignment Mechanism: Perfect Doctor

Little & Rubin (2000):

Patient	Y(0)	Y(1)	τ	T
1	(1)	6	(5)	1
2	(3)	12	(9)	1
3	9	(8)	(-1)	0
4	11	(10)	(-1)	0
Mean	10	9	(3)	

An Assignment Mechanism: Perfect Doctor

Little & Rubin (2000):

Patient	Y(0)	Y(1)	τ	T
1	(1)	6	(5)	1
2	(3)	12	(9)	1
3	9	(8)	(-1)	0
4	11	(10)	(-1)	0
Mean	10	9	(3)	

Clearly, treatment is beneficial:

$$\overline{Y(1)} - \overline{Y(0)} = 9 - 6 = 3$$

An Assignment Mechanism: Perfect Doctor

Little & Rubin (2000):

Patient	Y(0)	Y(1)	τ	T
1	(1)	6	(5)	1
2	(3)	12	(9)	1
3	9	(8)	(-1)	0
4	11	(10)	(-1)	0
Mean	10	9	(3)	

Clearly, treatment is beneficial:

$$\overline{Y(1)} - \overline{Y(0)} = 9 - 6 = 3$$

This assg mechanism is non-ignorable, confounded.

Next:
Inference and Testing
PS2 due

References I

- Bertrand, Marianne, and Sendhil Mullainathan. 2004. “Are Emily and Greg More Employable Than Lakisha and Jamal? A Field Experiment on Labor Market Discrimination.” *American Economic Review* 94 (4): 991–1013.
- Gerber, Alan S., and Donald P. Green. 2012. *Field Experiments: Design, Analysis, and Interpretation*. New York, NY: WW Norton.
- Holland, Paul. 1986. “Statistics and Causal Inference.” *Journal of the American Statistical Association* 81 (396): 945–60.
- Sen, Maya, and Omar Wasow. 2016. “Race as a Bundle of Sticks: Designs That Estimate Effects of Seemingly Immutable Characteristics.” *Annual Reviews of Political Science* 19: 499–522.