TRƯỜNG ĐẠI HỌC TÔN ĐỨC THẮNG
KHOA TOÁN – THỐNG KÊ

[Chèn Logo TDTU tại đây]

TIỂU LUẬN CUỐI KỲ

Năm học: 2024 – 2025 Học kỳ: 2

Môn học: Soạn thảo tài liệu khoa học bằng LaTeX
Mã môn học: ...
Nhóm thực hiện: Nhóm 1
Mã đề: ...
Giảng viên giảng dạy: ...

TP. HCM, ngày ... tháng ... năm 2024

LỜI CAM KẾT

- Tiểu luận này được nhóm 1 biên soạn lại bằng phần mềm LaTeX với mục đích hoàn thành môn học thực hành Soạn thảo tài liệu khoa học bằng LaTeX, ngoài ra không còn mục đích nào khác.
- Tiểu luận này được biên soạn lại bằng LaTeX dựa trên tài liệu sau:
  + Tiêu đề: Error estimates for the finite element method of the Navier-Stokes-Poisson-Nernst-Planck equations
  + Tác giả: Minghao Li, Zhenzhen Li

BẢNG PHÂN CÔNG CÔNG VIỆC

STT	Họ và tên	Nội dung phân công	Mức độ hoàn thành
1	...	...	...
2	...	...	...
3	...	...	...
4	...	...	...

Appl. Numer. Math. 197 (2024) 186–209

Contents lists available at ScienceDirect

Applied Numerical Mathematics

journal homepage: www.elsevier.com/locate/apnum

ELSEVIER

Error estimates for the finite element method of the Navier-Stokes-Poisson-Nernst-Planck equations ⋆

Minghao Li[a,*,1], Zhenzhen Li[b]

[a] College of Science, Henan University of Technology, Zhengzhou, 450001, China

[b] College of Mathematics and Information Science, Zhengzhou University of Light Industry, Zhengzhou, 450002, China

ARTICLE INFO

Keywords:
NSPNP equations
Unconditional stability
Unconditional error estimates

Abstract

In this paper, we consider the linearized backward Euler finite element scheme for the Navier-Stokes-Poisson-Nernst-Planck (NSPNP) equations. We aim to derive the unconditional error estimates of all variables in $l^\infty(L^2)$ and $l^\infty(H^1)$ norms. Compared with some previous results, we remove the mesh restriction. In addition, we derive two important physical properties for the proposed numerical scheme. Finally, three numerical examples are implemented to verify the theoretical analysis.

1. Introduction

The NSPNP equations describe the dynamics of electrically charged fluids, the motions of ionized particles or molecules and their interactions with electric fields and the surrounding fluid, which are widely used in colloidal science, biophysics, semiconductors, micro- and nanofluidics, and electrochemistry [42, 31, 46, 20]. The NSPNP equations are given as follows

$$\begin{cases} u_t - \Delta u + (u\cdot\nabla)u + \nabla p = -(c_1 - c_2)\nabla\psi, \ \text{in } \Omega\times(0,T], \\ \operatorname{div} u = 0, \ \text{in } \Omega\times(0,T], \\ c_{1t} - \Delta c_1 - \operatorname{div}(c_1\nabla\psi) + (u\cdot\nabla)c_1 = 0, \ \text{in } \Omega\times(0,T], \\ c_{2t} - \Delta c_2 + \operatorname{div}(c_2\nabla\psi) + (u\cdot\nabla)c_2 = 0, \ \text{in } \Omega\times(0,T], \\ -\Delta\psi = c_1 - c_2, \ \text{in } \Omega\times(0,T], \\ u = 0, \ \dfrac{\partial c_1}{\partial n} = \dfrac{\partial c_2}{\partial n} = \dfrac{\partial \psi}{\partial n} = 0, \ \text{on } \partial\Omega\times(0,T], \\ c_1(x,0) = c_{10}(x), \ c_2(x,0) = c_{20}(x), \ u(x,0) = u_0(x), \end{cases} \tag{1}$$

⋆ This work was supported by Cultivation Programme for Young Backbone Teachers in Henan University of Technology, Natural Science Foundation of Henan Province (222300420143, 232300420117, 232300420118), and the Innovative Funds Plan of Henan University of Technology (2021ZKCJ11).

https://doi.org/10.1016/j.apnum.2023.11.012

Received 14 February 2023; Received in revised form 27 August 2023; Accepted 16 November 2023

Available online 20 November 2023
0168-9274/© 2023 IMACS. Published by Elsevier B.V. All rights reserved.
*Corresponding author.
lyminghao@126.com (M. Li)
ORCID(s): 0000-0000-0000-0000 (M. Li)
[1] *E-mail address:* lyminghao@126.com (M. Li).

where $\Omega \in \mathcal{R}^d$, $d = 2, 3$ is a bounded domain with a Lipschitz continuous boundary $\partial\Omega$, n is the outward unit-normal vector to $\partial\Omega$, c_1 and c_2 are the concentrations of positive and negative charges, u and p are the velocity and pressure of the fluid, and ψ is the quasi-electrostatic potential. For the qualitative analysis of the solutions, such as the existence, uniqueness, stability and asymptotic behavior, we refer to [47, 29, 12, 1, 6, 5, 19].

Many numerical methods have been developed for the PNP and NSPNP equations. It is crucial to preserve some important physical properties at the discrete level, such as the mass conservation, concentration positivity and energy stability, and the related work can be referred to [13, 22, 39, 37, 11, 43, 14, 44, 45, 52, 40, 48, 28, 18, 7, 17]. In addition, some numerical techniques and implementation strategies were proposed to improve the performance efficiency, such as the adaptive finite element methods [21, 58, 60], the two grid methods [49], the stabilized techniques [55, 32], the proper orthogonal decomposition method [9], and some improved iteration techniques [59, 56, 38, 2].

There are some studies about the error estimates of the finite element method for the PNP and NSPNP equations. Sun and He analyzed an implicit Crank-Nicolson finite element scheme for the PNP equations in [53], and obtained the optimal error estimates in $l^\infty(H^1)$ norm for all variables, however, the convergence order of $l^\infty(L^2)$ norm was suboptimal. They also proposed the mixed finite element method for the PNP equations [23], and obtained optimal error estimates for $l^\infty(H^1)$ and $l^\infty(L^2)$ norms, and later they discussed the PNP-Stokes equations and modified NSPNP equations [24, 25]. Shi and Yang et al. investigated the conforming and nonconforming finite element methods on rectangular meshes [50, 51], and obtained the superconvergence results by integral identities techniques. Yang et al. researched the virtual element method on general polygonal meshes [57], and derived the error estimates in $l^\infty(H^1)$ and $l^\infty(L^2)$ norms. However, due to the use of inverse inequality and mathematical inductive argument, a mesh restriction is needed in the above references [53, 23, 24, 25, 50, 51, 57]. Gao and He [15, 16] presented linearized conservative finite element and mixed finite element methods for the PNP equations, and established unconditionally optimal error estimates for all variables in $l^2(H^1)$ and $l^\infty(L^2)$ norms, but they do not derive the error estimates in $l^\infty(H^1)$ norm. For the convergence analysis and error estimates of some other gradient flow model coupled with fluid motion, such as the Cahn-Hilliard-Hele-Shaw equations and the Cahn-Hilliard-Navier-Stokes system, one can refer to [4, 3, 41, 10].

In this paper, we focus on the error estimates of the linearized backward Euler finite element scheme for the NSPNP equations, and use low order conforming finite element pairs. Firstly, under an inductive hypothesis, we derive the existence and uniqueness of solutions, and the unconditional error estimates of all variables in $l^\infty(L^2)$ norms. Then, we obtain the $l^\infty(H^1)$ and $l^2(H^2)$ norm stability of the finite element solutions. Next, based on the stability, we deduce the unconditional error estimates in $l^\infty(H^1)$ norm. Moreover, the global mass conservation and the energy stability are also derived for the proposed numerical scheme. Finally, several numerical examples are implemented to verify the theoretical analysis. Compared with the results in [53, 23, 24, 25, 50, 51, 57], we remove the mesh restriction.

Throughout the article, the notations of the standard Sobolev spaces are used. The generic positive constant C with or without subscript is independent of the mesh size h and the time step τ.

2. Preliminary

3. The linearized backward Euler scheme

4. The error estimates

4.1. The $l^\infty(L^2)$-norm error estimates

4.2. The $l^\infty(H^1)$-norm error estimate

4.3. Improved error estimate for the $l^\infty(L^2)$-norm

5. Two physical properties

6. Numerical implementation

7. Conclusion and future work

References

[1] Bothe, D., Fischer, A., Saal, J., 2014. Global well-posedness and stability of electrokinetic flows. SIAM J. Math. Anal. 46, 1263–1316.

[2] Bousquet, A., Hu, X., Metti, M.S., et al., 2018. Newton solvers for drift-diffusion and electrokinetic equations. SIAM J. Sci. Comput. 40, B982–B1006.

[3] Chen, W., Feng, W., Liu, Y., et al., 2019. A second order energy stable scheme for the Cahn-Hilliard-Hele-Shaw equations. Discrete Contin. Dyn. B 24, 149–182.

[4] Chen, W., Liu, Y., Wang, C., et al., 2016. Convergence analysis of a fully discrete finite difference scheme for the Cahn-Hilliard-Hele-Shaw equation. Math. Comput. 85, 2231–2257.

[5] Constantin, P., Ignatova, M., 2019. On the Nernst-Planck-Navier-Stokes system. Arch. Ration. Mech. Anal. 232, 1379–1428.

[6] Constantin, P., Ignatova, M., Lee, F.N., 2021. Nernst-Planck-Navier-Stokes systems far from equilibrium. Arch. Ration. Mech. Anal. 240, 1147–1168.

[7] Dehghan, M., Gharibi, Z., 2022. An analysis of weak Galerkin finite element method for a steady state Boussinesq problem. J. Comput. Appl. Math. 406, 114029.

[8] Dehghan, M., Gharibi, Z., Ruiz-Baier, R., 2023. Optimal error estimates of coupled and divergence-free virtual element methods for the Poisson-Nernst-Planck/Navier-Stokes equations and applications in electrochemical systems. J. Sci. Comput. 94, 72.

[9] Dehghan, M., Hooshyarfarzin, B., Abbaszadeh, M., 2022. Numerical simulation based on a combination of finite element method and proper orthogonal decomposition to prevent the groundwater contamination. Eng. Comput. 38, 3445–3461.

[10] Diegel, A., Wang, C., Wang, X., et al., 2017. Convergence analysis and error estimates for a second order accurate finite element method for the Cahn-Hilliard-Navier-Stokes system. Numer. Math. 137, 495–534.

[11] Ding, J., Wang, Z., Zhou, S., 2020. Structure-preserving and efficient numerical methods for ion transport. J. Comput. Phys. 418, 109597.

[12] Fischer, A., Saal, J., 2017. Global weak solutions in three space dimensions for electrokinetic flow processes. J. Evol. Equ. 17, 309–333.

[13] Flavell, A., Machen, M., Eisenberg, B., et al., 2014. A conservative finite difference scheme for Poisson-Nernst-Planck equations. J. Comput. Electron. 13, 235–249.

[14] Fu, G., Xu, Z., 2022. High-order space-time finite element methods for the Poisson-Nernst-Planck equations: positivity and unconditional energy stability. Comput. Methods Appl. Mech. Eng. 395, 115031.

[15] Gao, H., He, D., 2017. Linearized conservative finite element methods for the Nernst-Planck-Poisson equations. J. Sci. Comput. 72, 1269–1289.

[16] Gao, H., Sun, P., 2018. A linearized local conservative mixed finite element method for Poisson-Nernst-Planck equations. J. Sci. Comput. 77, 793–817.

[17] Gharibi, Z., Dehghan, M., 2021. Convergence analysis of weak Galerkin flux-based mixed finite element method for solving singularly perturbed convection-diffusion-reaction problem. Appl. Numer. Math. 163, 303–316.

[18] Gharibi, Z., Dehghan, M., Abbaszadeh, M., 2021. Numerical analysis of locally conservative weak Galerkin dual-mixed finite element method for the time-dependent Poisson-Nernst-Planck system. Comput. Math. Appl. 92, 88–108.

[19] Gong, H., Wang, C., Zhang, X., 2021. Partial regularity of suitable weak solutions of the Navier-Stokes-Planck-Nernst-Poisson equation. SIAM J. Math. Anal. 53, 3306–3337.

[20] Gubbiotti, A., Baldelli, M., Muccio, G.D., et al., 2022. Electroosmosis in nanopores: computational methods and technological applications. Adv. Phys. X 7, 2036638.

[21] Hao, T., Ma, M., Xu, X., 2022. Adaptive finite element approximation for steady-state Poisson-Nernst-Planck equations. Adv. Comput. Math. 48, 1–27.

[22] He, D., Pan, K., Yue, X., 2019. A positivity preserving and free energy dissipative difference scheme for the Poisson-Nernst-Planck system. J. Sci. Comput. 81, 436–458.

[23] He, M., Sun, P., 2017. Error analysis of mixed finite element method for Poisson-Nernst-Planck system. Numer. Methods Partial Differ. Equ. 33, 1924–1948.

[24] He, M., Sun, P., 2018. Mixed finite element analysis for the Poisson-Nernst-Planck/Stokes coupling. J. Comput. Appl. Math. 341, 61–79.

[25] He, M., Sun, P., 2021. Mixed finite element method for modified Poisson-Nernst-Planck/Navier-Stokes equations. J. Sci. Comput. 87, 1–33.

[26] Heywood, J.G., Rannacher, R., 1982. Finite element approximation of the nonstationary Navier-Stokes problem. I. Regularity of solutions and second-order error estimates for spatial discretization. SIAM J. Numer. Anal. 19, 275–311.

[27] Heywood, J.G., Rannacher, R., 1990. Finite-element approximation of the nonstationary Navier-Stokes problem. Part IV: Error analysis for second-order time discretization. SIAM J. Numer. Anal. 27, 353–384.

[28] Huang, F., Shen, J., 2021. Bound/positivity preserving and energy stable scalar auxiliary variable schemes for dissipative systems: applications to Keller-Segel and Poisson-Nernst-Planck equations. SIAM J. Sci. Comput. 43, A1832–A1857.

[29] Jerome, J.W., Sacco, R., 2009. Global weak solutions for an incompressible charged fluid with multi-scale couplings: initial-boundary-value problem. Nonlinear Anal. 71, e2487–e2497.

[30] John, V., 2016. Finite Element Methods for Incompressible Flow Problems. Springer-Verlag, Berlin.

[31] Kirby, B.J., 2010. Micro- and Nanoscale Fluid Mechanics: Transport in Microfluidic Devices. Cambridge University Press, New York.

[32] Li, J., Ying, J., 2022. A stabilized finite volume element method for solving Poisson-Nernst-Planck equations. Int. J. Numer. Methods Biomed. Eng. 38, e3543.

[33] Li, M., Xiao, L., Li, Z., 2023a. The supercloseness property of the Stoke projection for the transient Navier-Stokes equations and global superconvergence analysis. J. Appl. Math. Comput. 69, 2041–2057.

[34] Li, Z., Li, M., Shi, D., 2021. Unconditional convergence and superconvergence analysis for the transient Stokes equations with damping. Appl. Math. Comput. 389, 125572.

[35] Li, Z., Xiao, L., Li, M., et al., 2023b. Error estimates for the finite element method of the chemotaxis-Navier-Stokes equations. J. Appl. Math. Comput. 69, 1–27.

[36] Lin, Q., Lin, J., 2006. Finite Element Methods: Accuracy and Improvement. Science Press, Beijing.

[37] Liu, C., Wang, C., Wise, S., et al., 2021. A positivity-preserving, energy stable and convergent numerical scheme for the Poisson-Nernst-Planck system. Math. Comput. 90, 2071–2106.

[38] Liu, C., Wang, C., Wise, S.M., et al., 2022. An iteration solver for the Poisson-Nernst-Planck system and its convergence analysis. J. Comput. Appl. Math. 406, 114017.

[39] Liu, H., Maimaitiyiming, W., 2021. Efficient, positive, and energy stable schemes for multi-d Poisson-Nernst-Planck systems. J. Sci. Comput. 87, 1–36.

[40] Liu, X., Xu, C., 2017. Efficient time-stepping/spectral methods for the Navier-Stokes-Nernst-Planck-Poisson equations. Commun. Comput. Phys. 21, 1408–1428.

[41] Liu, Y., Chen, W., Wang, C., et al., 2017. Error analysis of a mixed finite element method for a Cahn-Hilliard-Hele-Shaw system. Numer. Math. 135, 679–709.

[42] Markowich, P.A., Ringhofer, C.A., Schmeiser, C., 1990. Semiconductor Equations. Springer-Verlag, New York.

[43] Metti, M.S., Xu, J., Liu, C., 2016. Energetically stable discretizations for charge transport and electrokinetic models. J. Comput. Phys. 306, 1–18.

[44] Prohl, A., Schmuck, M., 2009. Convergent discretizations for the Nernst-Planck-Poisson system. Numer. Math. 111, 591–630.

[45] Prohl, A., Schmuck, M., 2010. Convergent finite element discretizations of the Navier-Stokes-Nernst-Planck-Poisson system. ESAIM: Math. Model. Numer. Anal. 44, 531–571.

[46] Ramos, A., 2011. Electrokinetics and Electrohydrodynamics in Microsystems. Springer, New York.

[47] Schmuck, M., 2009. Analysis of the Navier-Stokes-Nernst-Planck-Poisson system. Math. Models Methods Appl. Sci. 19, 993–1014.

[48] Shen, J., Xu, J., 2021. Unconditionally positivity preserving and energy dissipative schemes for Poisson-Nernst-Planck equations. Numer. Math. 148, 671–697.
[49] Shen, R., Shu, S., Yang, Y., et al., 2020. A decoupling two-grid method for the time-dependent Poisson-Nernst-Planck equations. Numer. Algorithms 83, 1613–1651.
[50] Shi, D., Yang, H., 2019. Superconvergence analysis of finite element method for Poisson-Nernst-Planck equations. Numer. Methods Partial Differ. Equ. 35, 1206–1223.
[51] Shi, X., Lu, L., 2021. Nonconforming finite element method for coupled Poisson-Nernst-Planck equations. Numer. Methods Partial Differ. Equ. 37, 2714–2729.
[52] Su, S., Tang, H., 2022. A positivity-preserving and free energy dissipative hybrid scheme for the Poisson-Nernst-Planck equations on polygonal and polyhedral meshes. Comput. Math. Appl. 108, 33–48.
[53] Sun, Y., Sun, P., Zheng, B., et al., 2016. Error analysis of finite element method for Poisson-Nernst-Planck equations. J. Comput. Appl. Math. 301, 28–43.
[54] Wang, C., Bao, J., Pan, W., et al., 2017. Modeling electrokinetics in ionic liquids. Electrophoresis , 1–13.
[55] Wang, Q., Li, H., Zhang, L., et al., 2021. A stabilized finite element method for the Poisson-Nernst-Planck equations in three-dimensional ion channel simulations. Appl. Math. Lett. 111, 106652.
[56] Xie, D., Lu, B., 2020. An effective finite element iterative solver for a Poisson-Nernst-Planck ion channel model with periodic boundary conditions. SIAM J. Sci. Comput. 42, B1490–B1516.
[57] Yang, Y., Liu, Y., Shu, S., 2022a. Error analysis of virtual element methods for the time-dependent Poisson-Nernst-Planck equations. arXiv preprint arXiv:2207.07231 .
[58] Yang, Y., Shen, R., Fang, M., et al., 2022b. Local averaging type a posteriori error estimates for the nonlinear steady-state Poisson-Nernst-Planck equations. J. Comput. Appl. Math. 404, 113874.
[59] Ying, J., Fan, R., Li, J., et al., 2021. A new block preconditioner and improved finite element solver of Poisson-Nernst-Planck equation. J. Comput. Phys. 430, 110098.
[60] Zhu, W., Yang, Y., Ji, G., et al., 2022. Residual type a posteriori error estimates for the time-dependent Poisson-Nernst-Planck equations. J. Sci. Comput. 90, 1–35.