

SMP

11873803

MATH3401

work submission coversheet

Submit via Blackboard Course Link

 School of Mathematics and Physics  
 Faculty of Science

This cover sheet should be attached to the front of your submission (not behind plastic). It is important that each barcode on this page is read. Make sure that the barcode sections are devoid of ink or stains etc.

Last Name **White**Student Number **44990392**First Name **Ryan**Submission Topic **Assignment 3**Due date of submission **9/05/2022**Due time of submission **11:50 AM**Group **05**Actual submission date **8/05/2022**Name of tutor  
(if appropriate)Session attended  
(if appropriate)

day

time

**Marked****Yes / No****IMPORTANT POINTS:**

- Place the submission through the slot with the barcode **FACE UP**.
- It is your responsibility to ensure that you keep a copy of any work submitted.
- Please read the Statement of Original Authorship at the bottom of the page.
- Sign this coversheet in the space provided below.


**Statement of Original Authorship**

I, **Ryan White** hereby state that the work contained in this assignment has not previously been submitted for assessment, either in whole or in part, by either myself or any other student at either The University of Queensland or at any other tertiary institution except where explicitly acknowledged. To the best of my knowledge and belief, the assignment contains no material that has been previously published or written by another person except where due reference is made. I make this Statement in full knowledge of an understanding that, should it be found to be false, I will be subject to disciplinary action under Student Integrity and Misconduct Policy 3.60.04 of the University of Queensland.

The University of Queensland's policy on plagiarism can be found at  
<http://ppl.app.uq.edu.au/content/3.60.04-student-integrity-and-misconduct>

(Reference 3.60.04)

Signed .....

Date **08/05/2022** .....

# MATH3401 Assignment 3

Ryan White      s4499039

9th of May, 2022

## Question 1

a. Let

$$f(z) = \frac{7z^3}{z^3 - 13z}$$

If  $\lim_{z \rightarrow \infty} f(z) = 7$ , then  $\lim_{z \rightarrow 0} f(1/z) = 7$ .

$$\begin{aligned}\lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) &= \lim_{z \rightarrow 0} \frac{7\left(\frac{1}{z}\right)^3}{\left(\frac{1}{z}\right)^3 - 13\left(\frac{1}{z}\right)} \\ &= \lim_{z \rightarrow 0} \frac{7}{z^3\left(\frac{1}{z^3} - \frac{13}{z}\right)} \\ &= \lim_{z \rightarrow 0} \frac{7}{1 - 13z^2} \\ &= \frac{7}{1 - 13 \cdot 0} = 7\end{aligned}$$

Therefore,  $\lim_{z \rightarrow \infty} f(z) = 7$ .

b. Let

$$f(z) = \frac{7z^3}{z^2 + 13z}$$

If  $\lim_{z \rightarrow \infty} f(z) = \infty$ , then  $\lim_{z \rightarrow 0} 1/f(1/z) = 0$ .

$$\begin{aligned}\lim_{z \rightarrow 0} \frac{1}{f\left(\frac{1}{z}\right)} &= \lim_{z \rightarrow 0} \frac{\left(\frac{1}{z}\right)^2 + 13\left(\frac{1}{z}\right)}{\left(\frac{1}{z}\right)^3} \\ &= \lim_{z \rightarrow 0} z^3 \left(\frac{1}{z^2} + \frac{13}{z}\right) \\ &= \lim_{z \rightarrow 0} z + 13z^2 \\ &= 0\end{aligned}$$

Therefore,  $\lim_{z \rightarrow \infty} f(z) = \infty$ .

c. Let

$$f(z) = \frac{(az + b)^2}{(cz + d)^2} = \frac{a^2z^2 + 2abz + b^2}{c^2z^2 + 2cdz + d^2}$$

If  $\lim_{z \rightarrow \infty} f(z) = a^2/c^2$ , then  $\lim_{z \rightarrow 0} f(1/z) = a^2/c^2$ .

$$\begin{aligned}\lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) &= \lim_{z \rightarrow 0} \frac{a^2\left(\frac{1}{z}\right)^2 + 2ab\left(\frac{1}{z}\right) + b^2}{c^2\left(\frac{1}{z}\right)^2 + 2cd\left(\frac{1}{z}\right) + d^2} \\ &= \lim_{z \rightarrow 0} \frac{a^2\left(\frac{1}{z}\right)^2 + 2ab\left(\frac{1}{z}\right) + b^2}{c^2\left(\frac{1}{z}\right)^2 + 2cd\left(\frac{1}{z}\right) + d^2} \cdot \frac{z^2}{z^2} \\ &= \lim_{z \rightarrow 0} \frac{a^2 + 2abz + b^2z^2}{c^2 + 2cdz + d^2z^2} = \frac{a^2}{c^2}\end{aligned}$$

Therefore, provided  $c \neq 0$ ,  $\lim_{z \rightarrow \infty} f(z) = a^2/c^2$ .

## Question 2

- a. If  $f(z) = 2xy + i(x^2 - y^2)$  is analytic, the Cauchy-Riemann equations must hold. Since  $f(z)$  is of the form of a bivariate polynomial, it is defined on all  $\mathbb{C}$  (since polynomials are defined everywhere).  
 $f(z)$  is of the form  $f(x, y) = u(x, y) + iv(x, y)$ , with

$$\begin{aligned} u(x, y) &= 2xy & v(x, y) &= x^2 - y^2 \\ \Rightarrow u_x &= 2y & \Rightarrow v_x &= 2x \\ u_y &= 2x & v_y &= -2y \end{aligned}$$

For C/R to be satisfied,  $u_x = v_y$ , but  $2y \neq -2y$  anywhere except  $y = 0$ . Since  $f(z)$  is not  $\mathbb{C}$ -differentiable in the neighbourhood of  $y = 0$  (and hence nowhere else), it is nowhere analytic.

- b. We have

$$\begin{aligned} f(z) &= \sin(\bar{z}) \\ &= \sin(x + i(-y)) \\ &= \sin(x) \cos(-iy) + \cos(x) \sin(-iy) \\ &= \sin(x) \cosh(y) - i \cos(x) \sinh(y) \end{aligned}$$

The function  $\sin(\bar{z})$  is defined on all  $\mathbb{C}$  by the result in lectures, and  $f(z)$  is of the form  $f(x, y) = u(x, y) + iv(x, y)$  with

$$\begin{aligned} u(x, y) &= \sin(x) \cosh(y) & v(x, y) &= -\cos(x) \sinh(y) \\ \Rightarrow u_x &= \cos(x) \cosh(y) & \Rightarrow v_x &= \sin(x) \sinh(y) \\ u_y &= \sin(x) \sinh(y) & v_y &= -\cos(x) \cosh(y) \end{aligned}$$

Clearly,  $u_x \neq v_y$  and  $-v_x \neq u_y$  in general. Only at points where  $u_x = v_y = 0$  and  $u_y = -v_x = 0$  will  $f(z)$  be complex differentiable.

Recall that  $C/R \Leftrightarrow df/d\bar{z} = 0$ .

$$\Rightarrow \frac{df}{d\bar{z}} = \cos(\bar{z})$$

which is only 0 at  $\bar{z} = n\pi + \pi/2$ , where  $n \in \mathbb{Z}$ .

Therefore,  $f(z) = \sin(\bar{z})$  is differentiable at some points in  $\mathbb{C}$ , but not on their neighbourhoods and so  $f(z)$  is analytic nowhere.

## Question 3

The function

$$f(z) = x^4 + i(1 - y)^4$$

is of the form  $f(x, y) = u(x, y) + iv(x, y)$ , with

$$\begin{aligned} u(x, y) &= x^4 & v(x, y) &= (1 - y)^4 \\ \Rightarrow u_x &= 4x^3 & \Rightarrow v_x &= 0 \\ u_y &= 0 & v_y &= -4(1 - y)^3 \end{aligned}$$

As before,  $f(z)$  is defined on all  $z \in \mathbb{C}$  by polynomial existence.

Clearly,  $-v_x = u_y$  everywhere.  $u_x$  equals  $v_y$  only when

$$\begin{aligned} x = -(1 - y) &\Rightarrow x^3 = -(1 - y)^3 \\ 4x^3 &= -4(1 - y)^3 \\ \Rightarrow u_x &= v_y \end{aligned}$$

And so  $f(z)$  is only differentiable along the line  $x = y - 1$ , or equivalently  $y = x + 1$ . However,  $f(z)$  is not differentiable in any neighbourhood of  $y = x + 1$  and so is nowhere analytic.

## Question 4

- a. Since  $x$  and  $y$  are real numbers,  $f(z) = \sqrt{|xy|}$  is a real-valued function (due to the absolute value being taken). As such,  $f(z) = \sqrt{|xy|} + i \cdot 0$ . Therefore,  $u(x, y) = \sqrt{|xy|}$  and  $v(x, y) = 0 \Rightarrow v_x = v_y = 0$ .  $u_x$  and  $u_y$  can be calculated using the definition of the derivative. Firstly, consider approaching the origin along the  $x$ -axis (since the derivative is independent of approach path)

$$\begin{aligned} u_x(0, 0) &= \lim_{(\Delta x, 0) \rightarrow (0, 0)} \frac{u(x_0 + \Delta x, y_0) - u(x_0, y_0)}{\Delta x} \\ &= \lim_{(\Delta x, 0) \rightarrow (0, 0)} \frac{\sqrt{|0 \cdot 0 + \Delta x \cdot 0|} - \sqrt{|0 \cdot 0|}}{\Delta x} \\ &= \lim_{(\Delta x, 0) \rightarrow (0, 0)} \frac{0}{\Delta x} = 0 \end{aligned}$$

Now, approaching the origin along the  $y$ -axis:

$$\begin{aligned} u_y(0, 0) &= \lim_{(0, \Delta y) \rightarrow (0, 0)} \frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{\Delta y} \\ &= \lim_{(0, \Delta y) \rightarrow (0, 0)} \frac{\sqrt{|0 \cdot 0 + 0 \cdot \Delta y|} - \sqrt{|0 \cdot 0|}}{\Delta y} \\ &= \lim_{(0, \Delta y) \rightarrow (0, 0)} \frac{0}{\Delta y} = 0 \end{aligned}$$

Therefore,  $u_x = 0 = v_y$  and  $-v_x = 0 = u_y$  at the origin, and so C/R is satisfied at the origin.

- b. Since  $f$  differentiability is independent of approach path, choose the path  $y = x$  to approach the origin.

$$\Rightarrow f(z) = \sqrt{|x \cdot x|} = \sqrt{|x^2|}$$

Since  $x$  is real-valued,  $x^2 \geq 0$  and so

$$f(z) = \sqrt{x^2} = |x|$$

but  $|x|$  is not differentiable at the origin and so neither is  $f(z)$ .

- c. The definitions from lectures state that C/R is necessary for differentiability but not sufficient. As such, if C/R doesn't hold,  $f(z)$  is definitely not differentiable but if C/R does hold,  $f(z)$  is **not** necessarily differentiable and other tests are needed to properly determine the differentiability of  $f$ .

## Question 5

- a. Boisgerault defines a rectifiable path as  $C : [0, 1] \rightarrow \mathbb{C}$  if the curve  $C$  is piecewise continuous differentiable, with finite arc length. This holds if there are consecutive continuously differentiable paths over a partition over the domain such that the joining of all of the piecewise paths results in the path  $C$ .

Source: Boisgerault, S, 2017. *Complex Analysis and Applications*. Accessed at:

[https://diren.mines-paristech.fr/Sites/Complex-analysis/Complex%20Analysis%20and%20Applications%20\(a4\).pdf](https://diren.mines-paristech.fr/Sites/Complex-analysis/Complex%20Analysis%20and%20Applications%20(a4).pdf)

- b. An example of a non-rectifiable curve in  $\mathbb{C}$  is the curve  $f(z = x + iy) = x \sin(1/x)$ . As  $x$  approaches 0, the curve oscillates more frequently and the path length increases towards infinity.