


Magnitude & Delay Sweeps with Heat Maps


Subject: AB01_Jimin

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -19.5%)




Bicep Femoris (NoAssi = -40.5%)



Gastrocnemius (NoAssi = -12.6%)




Rectus Femoris (NoAssi = -20.2%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris







Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = 32.8%)




Bicep Femoris (NoAssi = -43.1%)







Gastrocnemius (NoAssi = -2.7%)



Rectus Femoris (NoAssi = 24.8%)







Peak EMG Activation Contour Map - RA Walking







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)



Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = 34.4%)



Bicep Femoris (NoAssi = 44.1%)











Gastrocnemius (NoAssi = -0.9%)



Rectus Femoris (NoAssi = 136.9%)




Peak EMG Activation Contour Map - RD Walking







Subject: AB02_Rajiv

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -7.5%)



Bicep Femoris (NoAssi = -4.8%)











Gastrocnemius (NoAssi = -4.7%)



Rectus Femoris (NoAssi = 60.3%)




Peak EMG Activation Contour Map - LG Walking























Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)



EMG Delay Sweeps for RA Walking (Normalized)



Gait Cycle (%)


Gait Cycle (%)

Gait Cycle (%)


Gait Cycle (%)

Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = -2.6%)




Bicep Femoris (NoAssi = 22.1%)







Gastrocnemius (NoAssi = -6.9%)



Rectus Femoris (NoAssi = -7.7%)


























Peak EMG Activation Contour Map - RA Walking







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = 2.8%)




Bicep Femoris (NoAssi = 6.1%)











Gastrocnemius (NoAssi = -8.6%)



Rectus Femoris (NoAssi = 23.9%)




Peak EMG Activation Contour Map - RD Walking




Subject: AB03_Amy

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -5.6%)




Bicep Femoris (NoAssi = 24.0%)



Gastrocnemius (NoAssi = -15.4%)




Rectus Femoris (NoAssi = -36.1%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris







Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = -18.2%)



Bicep Femoris (NoAssi = -6.8%)



Gastrocnemius (NoAssi = -3.7%)




Rectus Femoris (NoAssi = -1.9%)




Peak EMG Activation Contour Map - RA Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris




Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)

The figure displays a 6x4 grid of line plots showing normalized EMG magnitude for four muscles: Tibialis Anterior, Bicep Femoris, Gastrocnemius, and Rectus Femoris. The rows represent different delay values: 100 ms, 150 ms, 200 ms, 250 ms, and 300 ms. The columns represent the muscles. The x-axis for all plots is Gait Cycle (%) from 0 to 100. The y-axis is EMG Magnitude, with scales varying by muscle and delay. The legend indicates two conditions: NoAssi (dotted lines) and NoExo (solid lines). The plots show how muscle activity changes with delay and condition.




Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)

Tibialis Anterior (NoAssi = -10.8%)




Bicep Femoris (NoAssi = 7.4%)



Gastrocnemius (NoAssi = -60.5%)




Rectus Femoris (NoAssi = 91.6%)




Peak EMG Activation Contour Map - RD Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris







Subject: AB05_Maria

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -35.6%)




Bicep Femoris (NoAssi = -63.5%)



Gastrocnemius (NoAssi = -29.2%)




Rectus Femoris (NoAssi = 18.9%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris























Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = 17.8%)




Bicep Femoris (NoAssi = 55.3%)











Gastrocnemius (NoAssi = -7.1%)



Rectus Femoris (NoAssi = 12.4%)




Peak EMG Activation Contour Map - RA Walking























Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = -26.2%)




Bicep Femoris (NoAssi = 42.3%)



Gastrocnemius (NoAssi = 60.1%)




Rectus Femoris (NoAssi = 164.5%)




Peak EMG Activation Contour Map - RD Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris























Subject: AB08_Adrian

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)



EMG Delay Sweeps for LG Walking (Normalized)



Gait Cycle (%)

Gait Cycle (%)

Gait Cycle (%)


Gait Cycle (%)

Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -27.4%)




Bicep Femoris (NoAssi = -61.2%)



Gastrocnemius (NoAssi = -23.6%)




Rectus Femoris (NoAssi = -59.1%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris























Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = 49.5%)



Bicep Femoris (NoAssi = 40.9%)



Gastrocnemius (NoAssi = 4.4%)




Rectus Femoris (NoAssi = 41.5%)




Peak EMG Activation Contour Map - RA Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)



Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = -24.1%)




Bicep Femoris (NoAssi = -15.8%)







Gastrocnemius (NoAssi = 1.1%)



Rectus Femoris (NoAssi = 90.0%)




Peak EMG Activation Contour Map - RD Walking







Subject: AB11_Ryan

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)



EMG Delay Sweeps for LG Walking (Normalized)



100 ms
150 ms
200 ms
250 ms
300 ms
NoAssi
NoExo

100 ms
150 ms
200 ms
250 ms
300 ms
NoAssi
NoExo


100 ms
150 ms
200 ms
250 ms
300 ms
NoAssi
NoExo

100 ms
150 ms
200 ms
250 ms
300 ms
NoAssi
NoExo


100 ms
150 ms
200 ms
250 ms
300 ms
NoAssi
NoExo

Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -16.2%)




Bicep Femoris (NoAssi = -29.5%)



Gastrocnemius (NoAssi = -28.4%)




Rectus Femoris (NoAssi = 126.5%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris







Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = -5.6%)



Bicep Femoris (NoAssi = 162.6%)



Gastrocnemius (NoAssi = -17.8%)




Rectus Femoris (NoAssi = 104.7%)




Peak EMG Activation Contour Map - RA Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = -8.5%)




Bicep Femoris (NoAssi = -20.0%)











Gastrocnemius (NoAssi = 8.4%)



Rectus Femoris (NoAssi = 34.8%)




Peak EMG Activation Contour Map - RD Walking




Subject: AB15_Daniel

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = 3.7%)



Bicep Femoris (NoAssi = 8.5%)



Gastrocnemius (NoAssi = 13.9%)




Rectus Femoris (NoAssi = 114.3%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius







Rectus Femoris







Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)



Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = 1.8%)



Bicep Femoris (NoAssi = 24.4%)











Gastrocnemius (NoAssi = -3.0%)



Rectus Femoris (NoAssi = 121.9%)







Peak EMG Activation Contour Map - RA Walking







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = 7.6%)




Biceps Femoris (NoAssi = 28.8%)



Gastrocnemius (NoAssi = 36.5%)




Rectus Femoris (NoAssi = 49.3%)




Peak EMG Activation Contour Map - RD Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris




Subject: AB16_Ilseung

Level-Ground

EMG Magnitude Sweeps for LG Walking (Normalized)




EMG Delay Sweeps for LG Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - LG Walking (Phased)


Tibialis Anterior (NoAssi = -1.2%)




Bicep Femoris (NoAssi = -13.4%)



Gastrocnemius (NoAssi = -21.4%)




Rectus Femoris (NoAssi = -20.5%)




Peak EMG Activation Contour Map - LG Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris







Ramp-Ascent

EMG Magnitude Sweeps for RA Walking (Normalized)




EMG Delay Sweeps for RA Walking (Normalized)




Peak EMG Activation (% Change from NoExo) - RA Walking (Phased)


Tibialis Anterior (NoAssi = 19.1%)




Bicep Femoris (NoAssi = -8.7%)







Gastrocnemius (NoAssi = -5.9%)



Rectus Femoris (NoAssi = 52.9%)







Peak EMG Activation Contour Map - RA Walking







Ramp-Descent

EMG Magnitude Sweeps for RD Walking (Normalized)




EMG Delay Sweeps for RD Walking (Normalized)



Peak EMG Activation (% Change from NoExo) - RD Walking (Phased)


Tibialis Anterior (NoAssi = -2.0%)




Bicep Femoris (NoAssi = 37.8%)



Gastrocnemius (NoAssi = 80.2%)




Rectus Femoris (NoAssi = 21.6%)




Peak EMG Activation Contour Map - RD Walking


Tibialis Anterior




Bicep Femoris



Gastrocnemius




Rectus Femoris



N=6 Average Heat Map

Average Peak EMG (% Change from NoExo) - LG (N=6, Phased)


Tibialis Anterior (NoAssi: -12.3%)




Bicep Femoris (NoAssi: -18.4%)



Gastrocnemius (NoAssi: -13.4%)




Rectus Femoris (NoAssi: +13.0%)




Average Peak EMG (% Change from NoExo) - RA (N=6, Phased)


Tibialis Anterior (NoAssi: +11.2%)



Bicep Femoris (NoAssi: +21.2%)



Gastrocnemius (NoAssi: -3.7%)




Rectus Femoris (NoAssi: +36.5%)




Average Peak EMG (% Change from NoExo) - RD (N=6, Phased)

Tibialis Anterior (NoAssi: -8.8%)




Bicep Femoris (NoAssi: +17.8%)



Gastrocnemius (NoAssi: +18.1%)







Rectus Femoris (NoAssi: +73.5%)







N=6 Average Contour Map





Average Peak EMG Contour Map - LG Walking (N=6)



Average Peak EMG Contour Map - RA Walking (N=6)







Average Peak EMG Contour Map - RD Walking (N=6)







N=8 Average Contour Map

Average Peak EMG Contour Map - LG Walking (N=8)



Average Peak EMG Contour Map - RA Walking (N=8)



Average Peak EMG Contour Map - RD Walking (N=8)

