

Systems of many forms with differing degrees

Simon L. Rydin Myerson

2 September 2024

Warwick Mathematics Institute

Download these slides:

maths.fan/ICTS.pdf

Systems of polynomial equations to be solved in integers. Ubiquitous in mathematics; relatively few general results.

Notation (density of solutions)

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ will be a system of R **homogenous forms** of **degrees d_i** in $s > \sum d_i$ variables with integer coefficients.
 - We count solutions of $\vec{f} = \vec{0}$ in **integers of size B** , where B is big.
-
- $\vec{f}$ takes about $B^{\sum d_i}$ values; maybe it is zero about $\frac{1}{B^{\sum d_i}}$ of the time.
 - That would mean about $B^{s - \sum d_i}$ solutions.
 - Also need to consider the number of solutions modulo m for $m \in \mathbb{N}$.

Systems of polynomial equations to be solved in integers. Ubiquitous in mathematics; relatively few general results. **No completely general result possible**: Matiyasevich (1970), unsolvability of Hilbert's tenth problem.

Notation (density of solutions)

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ will be a system of R **homogenous forms** of **degrees d_i** in $s > \sum d_i$ variables with integer coefficients.
 - We count solutions of $\vec{f} = \vec{0}$ in **integers of size B** , where B is big.
 - $\vec{\alpha} \cdot \vec{f} = \sum^R \alpha_i f_i$ is nonzero and indefinite for all $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$.
 - We study $N_{\vec{f}}(B) = \#\{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\} : \vec{f}(\vec{x}) = \vec{0}\}$.
-
- $\vec{f}$ takes about $B^{\sum d_i}$ values; maybe it is zero about $\frac{1}{B^{\sum d_i}}$ of the time.
 - That would mean about $B^{s - \sum d_i}$ solutions.
 - Also need to consider the number of solutions modulo m for $m \in \mathbb{N}$.
 - Let's make this rigorous.

Systems of polynomial equations to be solved in integers. Ubiquitous in mathematics; relatively few general results. No completely general result possible: Matiyasevich (1970), unsolvability of Hilbert's tenth problem.

Notation (density of solutions)

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ will be a system of R homogenous forms of degrees d_i in $s > \sum d_i$ variables with integer coefficients.
 - We count solutions of $\vec{f} = \vec{0}$ in integers of size B , where B is big.
 - $\vec{\alpha} \cdot \vec{f} = \sum \alpha_i f_i$ is nonzero and indefinite for all $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$.
 - We study $N_{\vec{f}}(B) = \#\{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\} : \vec{f}(\vec{x}) = \vec{0}\}$.
-
- $\vec{f}$ takes about $B^{\sum d_i}$ values; maybe it is zero about $\frac{1}{B^{\sum d_i}}$ of the time.
 - That would mean about $B^{s - \sum d_i}$ solutions.
 - Also need to consider the number of solutions modulo m for $m \in \mathbb{N}$.
 - Let's make this rigorous.

Notation (density of solutions)

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ will be a system of R **homogenous forms** of **degrees d_i** in $s > \sum d_i$ variables with integer coefficients.
- We count solutions of $\vec{f} = \vec{0}$ in **integers of size B** , where B is big.
- $\vec{\alpha} \cdot \vec{f} = \sum^R \alpha_i f_i$ is nonzero and indefinite for all $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$.
- We study $N_{\vec{f}}(B) = \#\{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\} : \vec{f} = \vec{0}\} \leq (2B + 1)^s$.

Heuristics

- Model $\vec{x}$ by a random real vector $\vec{X}$, and model $f_i(\vec{x})$ by $\lfloor f_i(\vec{X}) \rfloor$.
- That is, let $\vec{X}$ be a **uniform random variable on $[-B, B]^s$** . Maybe $N_{\vec{f}}(B) \asymp (2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1)^R]$, which is **typically $\sim \nu_{\vec{f}} B^{s - \sum d_i}$** .

Notation (density of solutions)

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ will be a system of R **homogenous forms** of **degrees d_i** in $s > \sum d_i$ variables with integer coefficients.
- We count solutions of $\vec{f} = \vec{0}$ in **integers of size B** , where B is big.
- $\vec{\alpha} \cdot \vec{f} = \sum \alpha_i f_i$ is nonzero and indefinite for all $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$.
- We study $N_{\vec{f}}(B) = \#\{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\} : \vec{f} = \vec{0}\} \leq (2B + 1)^s$.

Heuristics

- Model $\vec{x}$ by a random real vector $\vec{X}$, and model $f_i(\vec{x})$ by $\lfloor f_i(\vec{X}) \rfloor$.
- That is, let $\vec{X}$ be a **uniform random variable on $[-B, B]^s$** . Maybe $N_{\vec{f}}(B) \asymp (2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1)^R]$, which is **typically $\sim \nu_{\vec{f}} B^{s - \sum d_i}$** .
- But: if $f(\vec{x}) = x_1^2 + x_2^2 - 3x_3^2$, then $N_f(B) = 1$ as $\vec{x} = \vec{0}$ “mod 2^∞ ”.
- Fix: let $\vec{X}_p$ be uniformly distributed on $\mathbb{Z}_p^s$. Predict

$$N_{\vec{f}}(B) = (1 + o(1))(2B)^s \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1)^R]$$

$$\cdot \prod_p \lim_{N \rightarrow \infty} p^{NR} \mathbb{P}[p^N \mid \vec{f}(\vec{X}_p)].$$

- Let $\vec{X}$ be a uniform random variable on $[-B, B]^s$. Maybe $N_{\vec{f}}(B) \asymp (2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R]$, which is typically $\sim \nu_{\vec{f}} B^{s - \sum d_i}$.
- But: if $R = 1$, $f(\vec{x}) = x_1^2 + x_2^2 - 3x_3^2$, then $N_{\vec{f}}(B) = 0$.
- Fix: let $\vec{X}_p$ be uniformly distributed on $\mathbb{Z}_p^s$. Perhaps

$$N_{\vec{f}}(B) = (1 + o(1))(2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R] \prod_p \lim_{N \rightarrow \infty} p^{NR} \mathbb{P}[p^N \mid \vec{f}(\vec{X}_p)].$$

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.
- We say $a = O(b)$ iff $a \ll b$ iff $|a| < Cb$ for some constant C . Also write $a \asymp b$ iff $a \ll b \ll a$. And put $a \sim b$ iff $a/b \rightarrow 1$ as $B \rightarrow \infty$. And put $a = o(b)$ iff $a/b \rightarrow 0$ as $B \rightarrow \infty$.

- Let $\vec{X}$ be a uniform random variable on $[-B, B]^s$. Maybe $N_{\vec{f}}(B) \asymp (2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R]$, which is typically $\sim \nu_{\vec{f}} B^{s - \sum d_i}$.
- But: if $R = 1$, $f(\vec{x}) = x_1^2 + x_2^2 - 3x_3^2$, then $N_f(B) = 0$.
- Fix: let $\vec{X}_p$ be uniformly distributed on $\mathbb{Z}_p^s$. Perhaps

$$N_{\vec{f}}(B) = (1 + o(1))(2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R] \prod_p \lim_{N \rightarrow \infty} p^{NR} \mathbb{P}[p^N \mid \vec{f}(\vec{X}_p)].$$

- This is the *analytic Hasse principle*; the *Manin-Peyre conjecture* is a more sophisticated version needed for $s \leq 2 \sum d_i$ or $\vec{f}$ singular.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.
- We say $a = O(b)$ iff $a \ll b$ iff $|a| < Cb$ for some constant C . Also write $a \asymp b$ iff $a \ll b \ll a$. And put $a \sim b$ iff $a/b \rightarrow 1$ as $B \rightarrow \infty$. And put $a = o(b)$ iff $a/b \rightarrow 0$ as $B \rightarrow \infty$.

- Let $\vec{X}$ be a uniform random variable on $[-B, B]^s$. Maybe $N_{\vec{f}}(B) \asymp (2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R]$, which is typically $\sim \nu_{\vec{f}} B^{s - \sum d_i}$.
- But: if $R = 1$, $f(\vec{x}) = x_1^2 + x_2^2 - 3x_3^2$, then $N_{\vec{f}}(B) = 0$.
- Fix: let $\vec{X}_p$ be uniformly distributed on $\mathbb{Z}_p^s$. Perhaps

$$N_{\vec{f}}(B) = (1 + o(1))(2B)^s \cdot \mathbb{P}[\vec{f}(\vec{X}) \in [0, 1]^R] \prod_p \lim_{N \rightarrow \infty} p^{NR} \mathbb{P}[p^N \mid \vec{f}(\vec{X}_p)].$$

- This is the *analytic Hasse principle*; the *Manin-Peyre conjecture* is a more sophisticated version needed for $s \leq 2 \sum d_i$ or $\vec{f}$ singular.

Notation


- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.
- We say $a = O(b)$ iff $a \ll b$ iff $|a| < Cb$ for some constant C . Also write $a \asymp b$ iff $a \ll b \ll a$. And put $a \sim b$ iff $a/b \rightarrow 1$ as $B \rightarrow \infty$. And put $a = o(b)$ iff $a/b \rightarrow 0$ as $B \rightarrow \infty$.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- Set $N_{\vec{f}}(B) := \#\{\vec{x} \in \mathbb{Z}^s : \vec{f}(\vec{x}) = \vec{0}, \|\vec{x}\|_{\infty} \leq B\}$, $\vec{\alpha} \cdot \vec{f} = \sum^R \alpha_i f_i$.

$$N_{\vec{f}}(B) = \int_0^1 \cdots \int_0^1 \sum_{\substack{\vec{x} \in \mathbb{Z}^s \\ \|\vec{x}\|_{\infty} \leq B}} e^{2\pi i \vec{t} \cdot \vec{f}(\vec{x})} d\vec{t}$$

$$|\sum_{x=1}^{50} e(tx^2/500)|$$



Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- We set $N_{\vec{f}}(B) := \#\{\vec{x} \in \mathbb{Z}^s : \vec{f}(\vec{x}) = \vec{0}, \|\vec{x}\|_{\infty} \leq B\}$.
- $\vec{f}$ is **smooth** if $\vec{f} = 0$ defines a smooth $s - R$ dimensional complex manifold in $\mathbb{C}^s \setminus \{\vec{0}\}$ (away from the origin).

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ **smooth**, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- We set $N_{\vec{f}}(B) := \#\{\vec{x} \in \mathbb{Z}^s : \vec{f}(\vec{x}) = \vec{0}, \|\vec{x}\|_{\infty} \leq B\}$.
- $\vec{f}$ is **smooth** if $\vec{f} = 0$ defines a smooth $s - R$ dimensional complex manifold in $\mathbb{C}^s \setminus \{\vec{0}\}$ (away from the origin).

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ **smooth**, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Hope for $s > 2dR$. Much work on the range for s if $R = 1$. For $R \geq 2$:

- $(d, R, s) = (2, 2, 11)$ by Munshi (2015) - $s = 10$, Li-RM-Vishe, soon!
- $d = 2, s \geq 9R$, RM (2018);
- $d = 3, s \geq 25R$, RM (2019);
- $(d, R, s) = (3, 2, 39)$ by Northey and Vishe (2024).

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- We set $N_{\vec{f}}(B) := \#\{\vec{x} \in \mathbb{Z}^s : \vec{f}(\vec{x}) = \vec{0}, \|\vec{x}\|_{\infty} \leq B\}$.
- **smooth** if $\vec{f} = 0$ defines an $s - R$ dimensional manifold in $\mathbb{C}^s \setminus \{\vec{0}\}$.

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ **smooth**, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Hope for $s > 2dR$. Much work on the range for s if $R = 1$. For $R \geq 2$:

- $(d, R, s) = (2, 2, 11)$ by Munshi (2015) - $s = 10$, Li-RM-Vishe, soon!
- $d = 2, s \geq 9R$, RM (2018);
- $d = 3, s \geq 25R$, RM (2019);
- $(d, R, s) = (3, 2, 39)$ by Northey and Vishe (2024).

For certain $\vec{f}$: $(d, R, s) = (2, 2, 10)$ Heath-Brown–Pierce 2015; $(2, 3, 20)$ Pierce-Schindler-Wood 2016; $(2, R, 6R)$ Browning-Pierce-Schindler 2024.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ smooth, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Generalisations to unequal d_i .

- Browning–Dietmann–Heath-Brown (2014), $d_1 = 2, d_2 = 3, s \geq 29$.
- If $d_1, \dots, d_R \leq d$, Browning–Heath-Brown (2017) handle around $d^3 R^2 2^d$ variables, improving on Schmidt (1985).

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ smooth, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Generalisations to unequal d_i .

- Browning–Dietmann–Heath-Brown (2014), $d_1 = 2, d_2 = 3, s \geq 29$.
- If $d_1, \dots, d_R \leq d$, Browning–Heath-Brown (2017) handle around $d^3 R^2 2^d$ variables, improving on Schmidt (1985).

Theorem (RM 2024)

$$s \geq R + \sum_{i=1}^R d_i 2^{d_i} 3^{2d(d-d_i)}.$$

Typically this is about $3^{2d^2} R$ variables; when $d_1 = \dots = d_R = d$ it is $(1 + d2^d)R$ variables. Improve both Birch's result and, in big R , BHB.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ smooth, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

Generalisations to unequal d_i .

- Browning–Dietmann–Heath-Brown (2014), $d_1 = 2, d_2 = 3, s \geq 29$.
- If $d_1, \dots, d_R \leq d$, Browning–Heath-Brown (2017) handle around $d^3 R^2 2^d$ variables, improving on Schmidt (1985).

Theorem (RM 2024)

$$s \geq R + \sum_{i=1}^R d_i 2^{d_i} 3^{2d(d-d_i)}.$$

Typically this is about $3^{2d^2} R$ variables; when $d_1 = \dots = d_R = d$ it is $(1 + d2^d)R$ variables. Improve both Birch's result and, in big R , BHB.

Notation

- $\vec{f}(\vec{x}) \in \mathbb{Z}[x_1, \dots, x_s]^R$ is a system of R forms in $s > \sum d_i$ variables with integer coefficients, and $N_{\vec{f}}(B) = \sum_{\vec{x} \in \mathbb{Z}^s \cap [-B, B]^s \setminus \{\vec{0}\}, \vec{f}(\vec{x}) = \vec{0}} 1$.

Theorem (Birch 1962)

We have $N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s-dR}$ as above if $\vec{f}$ smooth, $d_1 = \dots = d_R = d$,
 $s \geq (d-1)2^{d-1}R(R+1) + R$.

- If $d_1, \dots, d_R \leq d$, Browning–Heath-Brown (2017) handle around $d^3 R^2 2^d$ variables, improving on Schmidt (1985).

Theorem (RM 2024)

$$s \geq R + \sum_{i=1}^R d_i 2^{d_i} 3^{2d(d-d_i)}.$$

Typically this is about $3^{2d^2} R$ variables; when $d_1 = \dots = d_R = d$ it is $(1 + d2^d)R$ variables. Improve both Birch's result and, in big R , BHB.

$d_1 = 2, d_3 = 3, s \geq 5858$, worse than BDHB $s \geq 29$! Key ideas: p -adic repulsion and a way to extract lower-order terms in exponential sums.

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
 - The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
 - $\vec{g}$ is nonsingular if the $R \times s$ Jacobian matrix $(\partial g_i(\vec{x})/\partial x_j)_{ij}$ has rank R at every nontrivial complex solution $\vec{x}$ to $\vec{g}(\vec{x}) = \vec{0}$.
-
- If $\vec{g}$ is nonsingular and the number of variables s is very large, can we estimate $M(B)$?
 - In the case $R = 1, d = 2$ the breakthrough work of Margulis, which introduced ideas from ergodic theory, led to:

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
- $\vec{g}$ is nonsingular if the $R \times s$ Jacobian matrix $(\partial g_i(\vec{x})/\partial x_j)_{ij}$ has rank R at every nontrivial complex solution $\vec{x}$ to $\vec{g}(\vec{x}) = \vec{0}$.
- If $\vec{g}$ is nonsingular and the number of variables s is very large, can we estimate $M(B)$?
- In the case $R = 1$, $d = 2$ the breakthrough work of Margulis, which introduced ideas from ergodic theory, led to:

Theorem (Eskin, Margulis and Mozes 1998)

Let g be a single quadratic form. Suppose g is nonsingular, and not a multiple of a form with rational coefficients.

If $s \geq 5$, then $M(B) \sim \nu_g B^{s-2}$ as $B \rightarrow \infty$ for some $\nu \geq 0$.

If $\vec{X}$ is a uniform RV on $[-B, B]^s$, then $\nu_g B^{s-2} \sim B^s \mathbb{P}[|g(\vec{X})| \leq 1]$.

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
- $\vec{g}$ is nonsingular if the $R \times s$ Jacobian matrix $(\partial g_i(\vec{x})/\partial x_j)_{ij}$ has rank R at every nontrivial complex solution $\vec{x}$ to $\vec{g}(\vec{x}) = \vec{0}$.
- If $\vec{g}$ is nonsingular and the number of variables s is very large, can we estimate $M(B)$?
- In the case $R = 1$, $d = 2$ the breakthrough work of Margulis, which introduced ideas from ergodic theory, led to:

Theorem (Eskin, Margulis and Mozes 1998)

Let g be a single quadratic form. Suppose g is nonsingular, and not a multiple of a form with rational coefficients.

If $s \geq 5$, then $M(B) \sim \nu_g B^{s-2}$ as $B \rightarrow \infty$ for some $\nu \geq 0$.

If $\vec{X}$ is a uniform RV on $[-B, B]^s$, then $\nu_g B^{s-2} \sim B^s \mathbb{P}[|g(\vec{X})| \leq 1]$.

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
 - The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
 - $\vec{g}$ is nonsingular if the $R \times s$ Jacobian matrix $(\partial g_i(\vec{x})/\partial x_j)_{ij}$ has rank R at every nontrivial complex solution $\vec{x}$ to $\vec{g}(\vec{x}) = \vec{0}$.
 - $\vec{g}(\vec{x})$ is **irrational** if no $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$ satisfies $\vec{\alpha} \cdot \vec{g}(\vec{x}) \in \mathbb{Z}[\vec{x}]$.
-
- Bentkus and Götze (1999) used the circle method to give a new proof of EMM's result $M(B) \sim \nu_g B^{s-2}$ when **s** $\geq$ **9**.

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
- $\vec{g}$ is nonsingular if the $R \times s$ Jacobian matrix $(\partial g_i(\vec{x})/\partial x_j)_{ij}$ has rank R at every nontrivial complex solution $\vec{x}$ to $\vec{g}(\vec{x}) = \vec{0}$.
- $\vec{g}(\vec{x})$ is **irrational** if no $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$ satisfies $\vec{\alpha} \cdot \vec{g}(\vec{x}) \in \mathbb{Z}[\vec{x}]$.

- Bentkus and Götze (1999) used the circle method to give a new proof of EMM's result $M(B) \sim \nu_g B^{s-2}$ when $s \geq 9$.

Theorem (Müller, 2008, slightly rephrased)

If $d_1 = \dots = d_R = 2$, $\vec{g}$ is nonsingular and irrational, and $s \geq 9R$, then we have $M(B) \sim \nu_{\vec{g}} B^{s-2}$ as $B \rightarrow \infty$ for some $\nu \geq 0$.

- Freeman (2000, 2001): Standardised this form of the circle method.
- Buterus-Götze-Hille-Margulis (2022): EMM's $R = 1$, $s \geq 5$, explicit, by circle method! Exp sums $\ll$ nice functions on 1-param groups

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
 - The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
 - $\vec{g}(\vec{x})$ is irrational if no $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$ satisfies $\vec{\alpha} \cdot \vec{g}(\vec{x}) \in \mathbb{Z}[\vec{x}]$.
-
- Diagonal case, $d_i = d$, indefinite: $M(B) > 0$ for $R = 1$ and $s \gg cd \log d$ by Davenport-Heilbronn-Roth-...; $M(B) > 0$ for $s \geq R \lceil Rd^2 \log 3Rd \rceil$ by Nadesalingam-Pitman (1989).
 - Schmidt (1980) gave a lower bound for $M(B)$ when s is (very) large.
 - Cubic case: Freeman (2004) got $M(B) > 0$ for $s > (10R)^{(10R)^5}$.
Chow (2014) took $R = 1$ and $s \geq 358\,823\,708$.
 - Asymptotics? Diagonal case: Freeman ('03), Wooley ('03).
 - Ergodic (EMM-like) methods? Special cases only, see overview of Yukie (arxiv:9710214).

- Diagonal case, $d_i = d$, indefinite: $M(B) > 0$ for $R = 1$ and $s \gg cd \log d$ by Davenport-Heilbronn-Roth-...; $M(B) > 0$ for $s \geq R \lceil Rd^2 \log 3Rd \rceil$ by Nadesalingam-Pitman (1989).
- Schmidt (1980) gave a lower bound for $M(B)$ when s is (very) large.
- Cubic case: Freeman (2004) got $M(B) > 0$ for $s > (10R)^{(10R)^5}$.
Chow (2014) took $R = 1$ and $s \geq 358\,823\,708$.
- Asymptotics? Diagonal case: Freeman ('03), Wooley ('03).
- Ergodic (EMM-like) methods? Special cases only, see overview of Yukie (arxiv:9710214).

Theorem (RM 2024)

Let $d_i \leq d$. Suppose that $\vec{g}$ is nonsingular and irrational, and that

$$s \geq R + \sum_{i=1}^R d_i \max\{d_i - 2, 1\} 2^{d_i} 3^{2d(d-d_i)}.$$

Then $M(B) \sim \nu_{\vec{g}} B^{n - \sum d_i}$ where $\nu_{\vec{g}} = \lim B^{\sum d_i} \mathbb{P}[\vec{g}(\vec{X}) \in [0, 1)^R]$.

Notation

- $\vec{g}(\vec{x}) \in \mathbb{R}[\vec{x}]^R$ is a system of R forms in s variables of degrees $d_i \geq 2$.
- The function $M(B) := \#\{\vec{x} \in \mathbb{Z}^s \setminus \{\vec{0}\} : \|\vec{g}(\vec{x})\|_\infty \leq 1, \|\vec{x}\|_\infty \leq B\}$.
- $\vec{g}(\vec{x})$ is irrational if no $\vec{\alpha} \in \mathbb{R}^R \setminus \{\vec{0}\}$ satisfies $\vec{\alpha} \cdot \vec{g}(\vec{x}) \in \mathbb{Z}[\vec{x}]$.

Theorem (RM 2024)

Let $d_i \leq d$. Suppose that $\vec{g}$ is nonsingular and irrational, and that

$$s \geq R + \sum_{i=1}^R d_i \max\{d_i - 2, 1\} 2^{d_i} 3^{2d(d-d_i)}.$$

Then $M(B) \sim \nu_{\vec{g}} B^{n - \sum d_i}$ where $\nu_{\vec{g}} = \lim B^{\sum d_i} \mathbb{P}[\vec{g}(\vec{X}) \in [0, 1)^R]$.

Setup for repulsion

- $e(t) = 2^{2\pi it}$, $\Delta_{\vec{h}} f(\vec{x}) = f(\vec{x} + \vec{h}) - f(\vec{x})$, $\Delta_{\vec{h}_1, \dots, \vec{h}_r} = \Delta_{\vec{h}_1} \cdots \Delta_{\vec{h}_r}$
- $f^{[k]}$ is the degree k part of f , with $|f^{[k]}|_\infty = |\text{biggest coefficient}|$.
- NB $\Delta_{\vec{h}_1, \dots, \vec{h}_k} f^{[k]}$ is a multilinear form in the $\vec{h}_i$, independent of $\vec{x}$
- $\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}$ is the vector of coefficients of $\Delta_{\vec{h}_1, \dots, \vec{h}_{k-1}, (\cdot)} f^{[k]}$

For $w \in C_c^\infty(\mathbb{R}^s)$, $f, g \in \mathbb{R}[\vec{x}]$ with degree $\leq k$, and $1 \leq P \leq B$ we have

$$\begin{aligned}
 & \left| \sum_{\vec{x} \in \mathbb{Z}^s} B^{-s} w(\vec{x}/B) e(g(\vec{x})) \sum_{\vec{x} \in \mathbb{Z}^s} B^{-s} w(\vec{x}/B) e(g(\vec{x}) + f(\vec{x})) \right|^{2^{k-1}} \\
 & \leq \left| \sum_{\vec{h} \in \mathbb{Z}^s} \sum_{\vec{x} \in \mathbb{Z}^s} B^{-2s} w(\vec{x} + \vec{h}/B) \overline{w}(\vec{x}/B) e(f(\vec{x}) - \Delta_{\vec{h}} g(\vec{x})) \right|^{2^{k-1}} \\
 & \ll \frac{1}{B^{(k-1)s}} \# \{(\vec{h}_1, \dots, \vec{h}_{k-1}, \vec{u}) \in \mathbb{Z}^{ks} : |\vec{h}_i| \leq B, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k} - \vec{u}| \leq B^{-1}\} \\
 & \ll \frac{1}{P^{(k-1)s}} \# \{(\vec{h}_1, \dots, \vec{h}_{k-1}, \vec{u}) \in \mathbb{Z}^{ks} : |\vec{h}_i| \leq P, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k} - \vec{u}| \leq \frac{P^{k-1}}{B^k}\}
 \end{aligned}$$

If $PB^{-k} \ll |f^{[k]}| \ll P^{1-k}$, this is

$$< P^{-(k-1)s} \# \{(\vec{h}_1, \dots, \vec{h}_{k-1}) \in \mathbb{Z}^{(k-1)s} : |\vec{h}_i| \leq P, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}| \leq |f^{[k]}| P^{k-2}\}_{10}$$

Setup for repulsion

- $e(t) = 2^{2\pi it}$, $\Delta_{\vec{h}} f(\vec{x}) = f(\vec{x} + \vec{h}) - f(\vec{x})$, $\Delta_{\vec{h}_1, \dots, \vec{h}_r} = \Delta_{\vec{h}_1} \cdots \Delta_{\vec{h}_r}$
- $f^{[k]}$ is the degree k part of f , with $|f^{[k]}|_\infty = |\text{biggest coefficient}|$.
- $\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}$ is the vector of coefficients of $\Delta_{\vec{h}_1, \dots, \vec{h}_{k-1}, (\cdot)} f^{[k]}$
- The *normalised* sum $S_w(B; f) = \sum_{\vec{x} \in \mathbb{Z}^s} B^{-s} w(\vec{x}/B) e(f(\vec{x}))$

For $f, g \in \mathbb{R}[\vec{x}]$ with degree $\leq k$, and $PB^{-k} \ll |f^{[k]}| \ll P^{1-k}$ we have

$$|S_w(B; f) S_w(B; g + f)|^{2^{k-1}} \ll P^{-(k-1)s} \\ \cdot \#\{(\vec{h}_1, \dots, \vec{h}_{k-1}) \in \mathbb{Z}^{(k-1)s} : |\vec{h}_i| \leq P, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}| \leq |f^{[k]}| P^{k-2}\}.$$

If $k = 3$, $f^{[3]}$ smooth get $\ll P^{-s}$ by ideas of Davenport; else, pigeonhole.

Suppose that, whenever $PB^{-k} \ll |\vec{\beta} \cdot \vec{f}^{[k]}| \ll P^{1-k}$, we have

$$S_w(B; \vec{\alpha} \cdot \vec{f}) S_w\left(B; (\vec{\alpha} + \vec{\beta}) \cdot \vec{f}\right) \ll P^{-E(k)}.$$

Then $\text{meas}\{\vec{\alpha} \in [0, 1]^R : |S_w(B; \vec{\alpha} \cdot \vec{f})| > B^{s-K}\} \ll B^{\sum^R d_i (K/E(d_i) - 1)}$.

If $\sum^R d_i/E(d_i) < 1$ then $\begin{cases} N_{\vec{f}}(B) \sim c_{\vec{f}} B^{s - \sum^R d_i} & \text{if } \vec{f} \in \mathbb{Z}[\vec{x}]^R, \text{ or} \\ M(B) \sim \nu_{\vec{f}} B^{s - \sum^R d_i} & \text{if } \vec{f} \text{ is irrational.} \end{cases}$

Setup for p -adic repulsion

- $e(t) = 2^{2\pi it}$, $\Delta_{\vec{h}} f(\vec{x}) = f(\vec{x} + \vec{h}) - f(\vec{x})$, $\Delta_{\vec{h}_1, \dots, \vec{h}_r} = \Delta_{\vec{h}_1} \cdots \Delta_{\vec{h}_r}$
- $f^{[k]}$ is the degree k part of f , with $|f^{[k]}|_\infty = |\text{biggest coefficient}|$.
- $\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}$ is the vector of coefficients of $\Delta_{\vec{h}_1, \dots, \vec{h}_{k-1}, (\cdot)} f^{[k]}$
- The *normalised* sum $S_w(B; f) = \sum_{\vec{x} \in \mathbb{Z}^s} B^{-s} w(\vec{x}/B) e(f(\vec{x}))$

For $f \in \mathbb{Z}[\vec{x}]$, $g \in \mathbb{R}[\vec{x}]$ with degree $\leq k$, and P prime with $P^M \asymp B^k$ so that $M = \lfloor k \log B / \log P \rfloor$, such that $P^{1-M} \leq |f^{[k]}|_P \leq P^{1-k}$ we have

$$|S_w(B; f) S_w(B; g + \frac{1}{P^M} f)|^{2^{k-1}} \ll P^{-(k-1)s} \cdot \#\{(\vec{h}_1, \dots, \vec{h}_{k-1}) \in \mathbb{Z}^{(k-1)s} : |\vec{h}_i| \leq P, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k}|_P \geq P |f^{[k]}|_P\}.$$

For $f^{[k]}$ smooth this is $\ll P^{-s}$ ($\mathbb{F}_p$ -points on varieties).

Suppose that, whenever $P^{1-M} \leq |\vec{b} \cdot \vec{f}^{[k]}|_P \leq P^{1-k}$, we have

$$S_w(B; \vec{\alpha} \cdot \vec{f}) S_w\left(B; \left(\vec{\alpha} + \frac{\vec{b}}{P^M}\right) \cdot \vec{f}\right) \ll P^{-E(k)}.$$

Then $\text{meas}\{\vec{\alpha} \in [0, 1]^R : |S_w(B; \vec{\alpha} \cdot \vec{f})| > B^{s-K}\} \ll B^{\sum^R d_i(K/E(d_i)-1)}$.

If $\sum^R d_i/E(d_i) < 1$ then $N_{\vec{f}}(B) \sim cB^{s-\sum^R d_i}$. Smooth $\vec{f}^{[d]}$: $E(d) = \frac{n-R+1}{2^d}$. 11

Notation

- $e(t) = 2^{2\pi it}$, $\Delta_{\vec{h}} f(\vec{x}) = f(\vec{x} + \vec{h}) - f(\vec{x})$, $\Delta_{\vec{h}_1, \dots, \vec{h}_r} = \Delta_{\vec{h}_1} \cdots \Delta_{\vec{h}_r}$
- $f^{[k]}$ is the degree k homogeneous part of a polynomial f
- NB $\Delta_{\vec{h}_1, \dots, \vec{h}_k} f^{[k]}$ is a multilinear form in the $\vec{h}_i$, independent of $\vec{x}$

Proposition

Given $w \in C_c^\infty(\mathbb{R}^s)$ there is $\tilde{w} \in C_c^\infty(\mathbb{R}^{ks})$ as follows. For any $f \in \mathbb{R}[\vec{x}]$ of degree $\leq d$, and any $1 < k < d$,

$$\begin{aligned}
 & \left| \sum_{\vec{x} \in \mathbb{Z}^s} \frac{1}{B^s} w\left(\frac{\vec{x}}{B}\right) e(f(\vec{x})) \right|^{(2^{d-1} + 1) \cdots (2^k + 1) 2^{k-1} 3^{(k+1)(d-k-1)+1}} \\
 & \leq B^{-ks} \sum_{\vec{h}_1, \dots, \vec{h}_{k-1} \in \mathbb{Z}^s} \left| \sum_{\vec{h}_k \in \mathbb{Z}^s} \tilde{w}\left(\frac{\vec{h}_1}{B}, \dots, \frac{\vec{h}_k}{B}\right) e\left(\Delta_{\vec{h}_1, \dots, \vec{h}_k} f^{[k]}\right) \right| \\
 & \ll \frac{1}{B^{(k-1)s}} \#\{(\vec{h}_1, \dots, \vec{h}_{k-1}, \vec{u}) \in \mathbb{Z}^{ks} : |\vec{h}_i| \leq P, |\vec{m}_{\vec{h}_1, \dots, \vec{h}_{k-1}}^{f, k} - \vec{u}| \leq \frac{1}{B}\}
 \end{aligned}$$

Lemma

Given $w \in C_c^\infty(\mathbb{R}^s)$ there is $\tilde{w} \in C_c^\infty(\mathbb{R}^{(d+1)s})$ as follows. For any $f \in \mathbb{R}[\vec{x}]$ of degree $\leq d$,

$$\left| \sum_{\vec{x} \in \mathbb{Z}^s} \frac{1}{B^s} w\left(\frac{\vec{x}}{B}\right) e(f(\vec{x})) \right|^{2^{d-1} + 1} \\ \leq \left| \sum_{\vec{x}_1, \dots, \vec{x}_d} \sum_{\vec{x}} \tilde{w}\left(\frac{\vec{x}_d}{B}, \dots, \frac{\vec{x}_1}{B}, \frac{\vec{x}}{B}\right) e\left(f^{[<d]} + F(\vec{x}; \vec{x}_1, \dots, \vec{x}_d)\right) \right|$$

with $F = f^{[d]} - \Delta_{\vec{x} + \vec{x}_1, \dots, \vec{x} + \vec{x}_d} f^{[d]}$ of degree $< d$ in $\vec{x}$ and ≤ 1 in each $\vec{x}_i$.

Lemma

For $L \in \text{GL}_s(\mathbb{R})$, $A := \{\|\vec{x}\|_\infty \leq B : \|L\vec{x} - \vec{u}\|_\infty \leq 1/B \text{ some } \vec{u} \in \mathbb{Z}^s\}$. Given $w \in C_c^\infty(\mathbb{R}^{2s})$ there are $\tilde{w}_{\vec{c}}^{L,B} \in C_c^\infty(\mathbb{R}^s)$, whose values, support and derivatives are bounded in terms of w only, such that for all real g ,

$$\left| \sum_{\vec{x}, \vec{y}} w\left(\frac{\vec{x}}{B}, \frac{\vec{y}}{B}\right) e(\vec{y} \cdot L\vec{x} + g(\vec{x})) \right|^3 \leq \sum_{\vec{c} \in A-A} \sum_{\vec{x}} \tilde{w}_{\vec{c}}^{L,B}\left(\frac{\vec{x}}{B}\right) e(\Delta_{\vec{c}} g(\vec{x})).$$